

Marine Institute, Rinville, Oranmore, Galway, H91 R673, Ireland.

Bord Iascaigh Mhara (BIM) Crofton Rd, Dún Laoghaire, Dublin, A96 E5A0

Inland Fisheries Ireland (IFI) 3044 Lake Dr, Cheeverstown, Dublin

Regulation (EU) 2017/1004 of the European Parliament and of the Council of 17 May 2017

on the establishment of a Union framework for the collection, management and use of data in the fisheries sector and support for scientific advice regarding the common fisheries policy and repealing Council Regulation (EC) No 199/2008 (recast)

Commission Delegated Decision (EU) 2021/1167 of 27 April 2021

establishing the multiannual Union programme for the collection and management of biological, environmental, technical and socioeconomic data in the fisheries and aquaculture sectors from 2022

Commission Implementing Decision (EU) 2021/1168 of 27 April 2021

establishing the list of mandatory research surveys at sea and thresholds as part of the multiannual Union programme for the collection and management of data in the fisheries and aquaculture sectors from 2022

Commission Implementing Decision (EU) 2022/39 of 12 January 2022

laying down rules on the format and timetables for the submission of national work plans and annual reports for data collection in the fisheries and aquaculture sectors, and repealing Implementing Decisions (EU) 2016/1701 and (EU) 2018/1283

Ireland Annual Report on data collection in the fisheries and aquaculture sectors

2023

Version 3.0

Galway, 26/06/2024

Contents

SECTION 1: GENERAL INFORMATION	6
DATA COLLECTION FRAMEWORK AT NATIONAL LEVEL	6
TEXT BOX 1A: TEST STUDIES DESCRIPTION	9
TEXT BOX 1B: OTHER DATA COLLECTION ACTIVITIES.....	10
SECTION 2: BIOLOGICAL DATA	15
TEXT BOX 2.1: LIST OF REQUIRED SPECIES/STOCKS	15
<i>North-East Atlantic</i>	15
TEXT BOX 2.1: LIST OF REQUIRED SPECIES/STOCKS	16
<i>North Sea and Eastern Arctic</i>	16
TEXT BOX 2.1: LIST OF REQUIRED SPECIES/STOCKS	17
<i>Other Regions</i>	17
TEXT BOX 2.2: PLANNING OF SAMPLING FOR BIOLOGICAL VARIABLES	18
<i>North-East Atlantic</i>	18
TEXT BOX 2.2: PLANNING OF SAMPLING FOR BIOLOGICAL VARIABLES	21
<i>North Sea and Eastern Arctic</i>	21
TEXT BOX 2.2: PLANNING OF SAMPLING FOR BIOLOGICAL VARIABLES	21
<i>Other Regions</i>	21
TEXT BOX 2.3: DIADROMOUS SPECIES DATA COLLECTION IN FRESHWATER.....	22
TEXT BOX 2.4: RECREATIONAL FISHERIES.....	32
<i>North-East Atlantic</i>	32
TEXT BOX 2.5: SAMPLING PLAN DESCRIPTION FOR BIOLOGICAL DATA	34
<i>North-East Atlantic</i>	34
TEXT BOX 2.5: SAMPLING PLAN DESCRIPTION FOR BIOLOGICAL DATA	41
<i>North Sea and Eastern Arctic</i>	41
TEXT BOX 2.5: SAMPLING PLAN DESCRIPTION FOR BIOLOGICAL DATA	42
<i>Other Regions</i>	42
TEXT BOX 2.6: RESEARCH SURVEYS AT SEA	43
International Ecosystem Survey in the Nordic Seas (ASH)	43
Irish Groundfish Survey (IBTS_Q4).....	44
International Blue Whiting Spawning Survey (IBWSS)	46
International Mackerel and Horse Mackerel Egg Survey (MEGS)	49
Spawning/Pre – Spawning Herring/Boarfish Acoustic Survey (WESPAS_IRL)	50
Nephrops UWT Survey (UWV 16 – 17, UWTV 19, UWTV 20 – 22)	52
Celtic Sea Herring Acoustic Survey (CSHAS_IRL).....	54
Irish Anglerfish and Megrim Survey (IAMS_IRL).....	57
Cockle North Irish Sea Survey (CNIS)	59
Oyster West of Ireland (OWI).....	60
Razor Clam Irish Sea (RCIS)	62
Irish Coastal Ecosystem Survey	63
Scallop Survey (SS_IRL)	65
SECTION 3: FISHING ACTIVITY DATA	67
TEXT BOX 3.1: FISHING ACTIVITY VARIABLES DATA COLLECTION STRATEGY	67
TEXT BOX 3.2: FISHING ACTIVITY VARIABLES DATA COLLECTION STRATEGY (FOR INLAND EEL COMMERCIAL FISHERIES).....	68
SECTION 4: IMPACT OF FISHERIES ON MARINE BIOLOGICAL RESOURCES.....	69

TEXT BOX 4.2: INCIDENTAL CATCHES OF SENSITIVE SPECIES.....	69
(Region/RFMO/RFO/IO: Please indicate per text box and update the table of contents).....	69
TEXT BOX 4.3: FISHERIES IMPACT ON MARINE HABITATS	71
SECTION 5: ECONOMIC AND SOCIAL DATA IN FISHERIES.....	74
TEXT BOX 5.2: ECONOMIC AND SOCIAL VARIABLES FOR FISHERIES DATA COLLECTION.....	74
SECTION 6: ECONOMIC AND SOCIAL DATA IN AQUACULTURE	75
TEXT BOX 6.1: ECONOMIC AND SOCIAL VARIABLES FOR AQUACULTURE DATA COLLECTION.....	75
SECTION 7: ECONOMIC AND SOCIAL DATA IN FISH PROCESSING	77
TEXT BOX 7.1: ECONOMIC AND SOCIAL VARIABLES FOR FISH PROCESSING DATA COLLECTION	77
ANNEX 1.1 - QUALITY REPORT FOR BIOLOGICAL DATA SAMPLING SCHEME.....	79
QUALITY REPORTS RELATED TO TABLES 2.2 AND 2.5 AND TEXT BOXES 2.2 AND 2.5	79
SAMPLING SCHEME IDENTIFIER: DEMERSAL AT-SEA	79
SAMPLING SCHEME IDENTIFIER: DEMERSAL AT-SEA ENHANCED.....	83
SAMPLING SCHEME IDENTIFIER: DEMERSAL ON - SHORE.....	87
SAMPLING SCHEME IDENTIFIER: PELAGIC ON – SHORE OTHERS	91
SAMPLING SCHEME IDENTIFIER: PELAGIC ON – SHORE TUNA.....	96
SAMPLING SCHEME IDENTIFIER: PELAGIC ON – SHORE SPRAT.....	99
SAMPLING SCHEME IDENTIFIER: PELAGIC ON – SHORE BOARFISH	104
SAMPLING SCHEME IDENTIFIER: PELAGIC ON – SHORE NORWEGIAN SPRING SPAWNING HERRING	108
SAMPLING SCHEME IDENTIFIER: PELAGIC ON – SHORE HERRING.....	112
SAMPLING SCHEME IDENTIFIER: PELAGIC AT - SEA OTHERS.....	116
SAMPLING SCHEME IDENTIFIER: PELAGIC AT - SEA TUNA	120
SAMPLING SCHEME IDENTIFIER: PELAGIC AT - SEA HERRING.....	123
SAMPLING SCHEME IDENTIFIER: PELAGIC AT - SEA ENHANCED.....	127
SAMPLING SCHEME IDENTIFIER: <i>NEPHROPS</i> AT - SEA SELF-SAMPLING, <i>NEPHROPS</i> VESSELS IN FU16	132
SAMPLING SCHEME IDENTIFIER: <i>NEPHROPS</i> AT - SEA, <i>NEPHROPS</i> VESSELS IN FU16	136
SAMPLING SCHEME IDENTIFIER: <i>NEPHROPS</i> ON - SHORE, <i>NEPHROPS</i> VESSELS EXCLUDING FU16	140
SAMPLING SCHEME IDENTIFIER: CRUSTACEA AT-SEA	143
SAMPLING SCHEME IDENTIFIER: CRUSTACEA ON - SHORE	145
SAMPLING SCHEME IDENTIFIER: MOLLUSCA ON - SHORE	148
QUALITY REPORTS RELATED TO TABLE 2.3 AND TEXT BOX 2.3	150
SAMPLING SCHEME IDENTIFIER: DIAD_ESB_COUNTER (<i>SALMO SALAR</i>).....	150
SAMPLING SCHEME IDENTIFIER: DIAD_ESB_EEL (<i>ANGUILLA ANGUILLA</i>)	155
SAMPLING SCHEME IDENTIFIER: DIAD_MI_EEL (<i>ANGUILLA ANGUILLA</i>).....	164
SAMPLING SCHEME IDENTIFIER: DIAD_MI_SALMON_CWT (<i>SALMO SALAR</i>)	173
SAMPLING SCHEME IDENTIFIER: DIAD_MI_SALMON_SEATROUT (<i>SALMO SALAR</i> & <i>SALMO TRUTTA</i>).....	177
SAMPLING SCHEME IDENTIFIER: WILD SALMON AND SEA TROUT TAGGING SCHEME RECREATIONAL	185
SAMPLING SCHEME IDENTIFIER: WILD SALMON AND SEA TROUT TAGGING SCHEME COMMERCIAL.....	189
SAMPLING SCHEME IDENTIFIER: BIOLOGICAL SAMPLING NSIC	194
SAMPLING SCHEME IDENTIFIER: BIOLOGICAL SAMPLING SMOLTS NSIC	197
SAMPLING SCHEME IDENTIFIER: EEL_ELVER_TRAP_DIADROMOUS (SCIENTIFIC)	201
SAMPLING SCHEME IDENTIFIER: EEL_SILVER_DIADROMOUS (SCIENTIFIC).....	207
SAMPLING SCHEME IDENTIFIER: EEL_FYKENET_DIADROMOUS (SCIENTIFIC)	213
QUALITY REPORTS RELATED TO TABLE 2.4 AND TEXT BOX 2.4	219
SAMPLING SCHEME IDENTIFIER: IMREC_CREEL	219
SAMPLING SCHEME IDENTIFIER: IMREC_SKP_DI	223
SAMPLING SCHEME IDENTIFIER: IMREC_OB_CH_SURVEY	227
SAMPLING SCHEME IDENTIFIER: IMREC_ANG_DI.....	231
SAMPLING SCHEME IDENTIFIER: TUNA_CHART.....	235

QUALITY REPORTS RELATED TO TABLE 2.6 AND TEXT BOX 2.6.....	239
SAMPLING SCHEME IDENTIFIER: IBTS_Q4.....	239
SAMPLING SCHEME IDENTIFIER: IBWSS_IRL.....	243
SAMPLING SCHEME IDENTIFIER: MEGS_IRL.....	248
SAMPLING SCHEME IDENTIFIER: WESPAS_IRL.....	252
SAMPLING SCHEME IDENTIFIER: NEPHROS UWTW SURVEY (UWTW 16 – 17, UWTW 19, UWTW 20 – 22)..	256
SAMPLING SCHEME IDENTIFIER: CSHAS_IRL.....	262
SAMPLING SCHEME IDENTIFIER: IAMS_IRL.....	266
SAMPLING SCHEME IDENTIFIER: RCIS_IRL.....	270
SAMPLING SCHEME IDENTIFIER: CNIS_IRL.....	278
SAMPLING SCHEME IDENTIFIER: OWI_IRL.....	283
SAMPLING SCHEME IDENTIFIER: ICEos_IRL.....	289
SAMPLING SCHEME IDENTIFIER: SS_IRL.....	293
ANNEX 1.2 - QUALITY REPORT FOR SOCIOECONOMIC DATA SAMPLING SCHEME	301
QUALITY REPORT RELATED TO TABLE 5.2 AND TEXT BOX 5.2.....	301
<i>Sector Name(s): Fisheries - Non-Probability Sample Survey</i>	<i>301</i>
QUALITY REPORT RELATED TO TABLE 5.2 AND TEXT BOX 5.2.....	308
<i>Sector Name(s): Fisheries – Sentinel Vessel Programme</i>	<i>308</i>
QUALITY REPORT RELATED TO TABLE 6.1 AND TEXT BOX 6.1.....	313
<i>Sector Name(s): Aquaculture Census</i>	<i>313</i>
QUALITY REPORT RELATED TO TABLE 6.1 AND TEXT BOX 6.1.....	318
<i>Sector Name(s): Aquaculture NPS PSS.....</i>	<i>318</i>

SECTION 1: GENERAL INFORMATION

Data collection framework at national level

General comment: Use this text box to describe how data collection is organised in your Member State (institutions involved, contact information) and in which regional coordination groups (RCG) your Member State participates.

Outline the general framework of the national data collection programme in relation to the relevant sections of the EU MAP. If applicable, indicate major methodological changes in approach compared to previous year(s), and to which section(s) they apply.

The framework of Ireland's national workplan is in accordance with Regulation (EU) 2017/1004 of the European Parliament and of the Council of 17 May 2017, Commission Delegated Decision (EU) 2021/1167 of 16 July 2021 and Commission Implementing Decision (EU) 2021/1168 of 16 July 2021 and following the supplied guidance document for the layout of text and tables. The submission is for the full period 2022-2027.

2023 changes

- Updated the SecWeb Text box with the latest text available
- Inclusion of Irish Coastal Ecosystem Survey (ICEcoS) and Scallop survey with associated Annex 1.1 and updated Tables 2.2 and 2.5. These two surveys were mentioned in the 2022 program as new surveys.
- Updated nine Annex 1.1 with information on Pelagic on shore and at-sea sampling schemes
- Updated Table 2.2 with additional rows on the collection of maturity data from four species *Solea solea*, *Molva molva*, *Pollachius pollachius* and *Pollachius virens*. These were omitted from the original WP. Updated Annex 1.1 for sampling scheme Mollusca-on-shore
- WP Excel tables: Table 6.1. Economic and social variables for aquaculture data collection strategy. An error in the table format was identified in column B (i.e., implementation year), accordingly 'annual' was replaced with the correct variable '2022-2024' (edits throughout sheet marked in red Rows 3-331).
- In previous years, due to numbers, macroalgae and land-based shellfish were amalgamated under the all-catching segment 'multispecies' to enable the total national value of the sector to be reported, while at the same time protecting the financial data of individual micro-businesses. The national macro-algal segment has now developed to the point with sufficient new companies that data can be disaggregated into the stand-alone segment 'Longline Macroalgae'. Data will be collected for this new segment from 2023 onwards. In contrast, the land-based shellfish units have declined in output and unit number below the point enabling responsible reporting of them as a separate segment (i.e., multispecies) particularly when macroalgae has been removed from this segment. For data collection from 2023 onwards, land-based shellfish units will be incorporated into the most appropriate larger shellfish segment 'Oyster Other methods' (Oyster on trestles) as the units involved are mainly hatcheries and nurseries that supply this segment (i.e., Rows 250-290). As a result of these changes to macroalgae and land-based shellfish, the 'multispecies' segment is now defunct and will not be used for data collection from 2023 onwards. Accordingly, there are edits in multiple rows marked by red font (i.e., addition of new rows for macroalgae – Rows 168-208) and strikethrough to delete the multispecies segments (Rows 332-370). These edits are also carried through to the relevant Annex 1.2
- Minor edits to text on pg. 35 to explain discrepancies between EUMAP and EUROSTAT census data because of reporting time lags.
- There are additional edits in Column H throughout the sheet marked in red to reflect a

change from NPS to PSS in line with updates to the MS's data collection strategy for all species. Format changes to index the Quality Annexes'

Major changes:

Biological data on exploited biological resources caught by Union commercial fisheries: The major changes in the programme compared to previous submission are the inclusion of self-sampling on the commercial fisheries, demersal, pelagic, *Nephrops* and shellfish into the at sea sampling plan sampling frame. This work was initially driven by the Coronavirus pandemic but has proven to be a valuable data collection portal and is included as part of the at sea sampling frames.

Ireland also plans to commence the development of Digital Data Collection on our port sampling on commercial fish species in order to improve the quality of the data collected as it will not only save time in data entry but will provide instant feedback in data quality by performing live checks. Ireland intends to build capacity from 2023 and develop expertise in boarfish ageing to support development of age-based assessment models. We hope to develop bilateral agreements for sampling of Albacore Tuna and Boarfish, finalise 4S on our pelagic sampling, based on stock ID work to date and depending on the results of the benchmark process in 2021/2022, we are looking to separate mixed stock catches of herring in 6a using genetics. This work will commence in 2023.

In order to support the objectives of Natura, MSFD, OSPAR and especially the CFP to minimise the effects of by-catch of protected, endangered and/or threatened (PET) species, data collection under the DCF will be further developed and expanded with an enhanced sampling programme. These fisheries are the gillnet vessels >10m targeting demersal species and the OTM and PTM vessels >10m targeting *Trachurus trachurus*. Other data collection activities and dedicated projects to evaluate the impact of fisheries on by-catch will be conducted as part of the EMFAF Biodiversity /Marine Knowledge schemes and are described in Text box 4.2 Incidental catches of sensitive species.

Surveys at sea: Assessment of offshore scallop stocks in the Celtic Sea and south Irish Sea is limited by lack of survey data. It is proposed to initiate Annual scallop surveys in the Celtic and Irish Sea, and will be due to commence in 2023

Ireland also plans to commence an annual Young Fish Survey in 2023 in order to provide more information on recruitment.

The Scallop survey and the annual young fish survey will be included in subsequent updates.

Recreational Fisheries: Irelands plans to move to routine data collection programme for the recreational fisheries. This was developed by Inland Fisheries Ireland as a pilot study and will feed into assessment requirements for marine species.

VME/habitat impacts: Encounters of VME indicator species in the catch will be identified as required during surveys and on the at sea sampling frames. Other data collection activities and dedicated projects to evaluate the impact of fisheries on habitats will be conducted as part of the EMFAF Biodiversity /Marine Knowledge schemes. These Schemes are still under development for the Irish Operational Programme.

Stomach sampling: For the Celtic Seas, there are currently neither clearly defined end-user needs for stomach sampling nor regionally coordinated sampling programmes or pilot studies. Ireland therefore does not intend to commence a stomach sampling programme in 2022.

To evaluate future needs for this data collection activity, Ireland is taking the following steps:

- A concept study to evaluate the data requirements for predator prey interactions as inputs to a Celtic Seas multi-species VPA assessment.
- Further development on indicators and data needs for MSFD descriptor 4 including the use of DCF survey data to support trophic guild assessments.

- An increase in capacity in fisheries ecosystem modelling and advice from 2024 onwards once concept studies are concluded.
- Close monitoring of the North Sea regionally coordinated pilot study on IBTS stomach sampling and the interaction with and advisory outputs of WGSAM over the next two years.
- A financial placeholder for future funding of stomach sampling in the DCF budget estimations under EMFAF.

BIM: The MS wants to move away a paper-based system with manual data entry towards an online data entry system where data is stored in a database as there are still some paper based surveys circulated to the industry. The first phase of development in late 2021 will be for the fisheries industry whereby survey forms will be available online and these will connect with a database back end. This will enable the digital capturing, storing and reporting of data. The EU - MAP system will facilitate the entry of data through an online web portal and through a data entry and integration layer which will allow for access to data in current and future systems. The EU - MAP system will include the development of a scalable database storage and support reporting through a BI module. The main objective of this system to support the mandatory EU - MAP reporting.

RCG Secretariat and Webpage: The project SecWeb (MARE2020-08) was setup with the aim of developing mechanisms to support the planning and execution of administrative tasks and the branding and online visibility of the Regional Coordination Groups (RCGs), with the aim to establish a long-term supportive structure (the RCGs' Secretariat).

Give full name, acronym and contact details of all institutes that contribute to the data collection activities, and describe briefly their role in the work plan.

Marine Institute (MI) Rinville, Oranmore, Galway, H91R673, Ireland. +35391 387200

Responsible for the modules in relation to test studies, other data collection activities, biological data collection (excluding recreational fisheries), diadromous fisheries, surveys at sea, impact of fisheries on marine biological resources and relevant quality reports.

Bord Iascaigh Mhara (BIM) Crofton Rd, Dún Laoghaire, Dublin, A96 E5A0 +3531 2144100

Responsible for the data collection in relation to the economic and social data in fisheries, aquaculture and fish processing and the relevant quality reports for socioeconomic data

Inland Fisheries Ireland (IFI) 3044 Lake Dr, Cheeverstown, Dublin +3531 8842600

Responsible for the collection and collation of data from the Diadromous fisheries recreational fisheries in Ireland and associated quality reports.

Provide a link to the national data collection website, if there is one.

<https://www.demap-ireland.ie/>

(max. 1000 words)

Text Box 1a: Test studies description

General comment: This text box fulfils Chapter II, section 1.2 of the EU MAP Delegated Decision annex.

1. Aim of the test study- Digital Data collection

Digital data collection has been commonly used in a number of our sampling programmes and surveys, for example for measuring *Nephrops* lengths, or collecting data during the Irish Groundfish Survey. However, there are still a number of sampling programmes that rely on data being recorded on paper initially before being transcribed. The aim of this test study is to further roll-out digital data collection to onshore commercial sampling in ports.

Port sampling and the associated data entry and screening is very time consuming and is also prone to human error when samplers are transcribing biological parameters collected (i.e. length, weight, sex, maturity and age) from data sheets onto the Stockman database. The objective of transitioning to digital data collection in the ports, is to streamline the data collection process, facilitating quality checking of the data in situ, and in real time, as data is being collected in the sampling location e.g. length weight regressions. Digital data collection will eliminate the need to transcribe data from datasheets, thus reducing the potential for human error and also freeing up valuable time for busy samplers.

There is also a need to digitise the data collection of commercial data at sea, however the sampling on shore has been identified as the first priority with a focus on digitising the data collection on board commercial vessels to come in subsequent years.

2. Duration of the test study

The initial pilot will be performed during 2022, with a further roll-out of devices and refinement of methods during 2023 onwards.

3. Methodology and expected outcomes of the test study

Ruggedised tablets running the Windows operating system will be purchased and a version of our “Stockman” application will be written to run on them in an off-line capacity. Users will enter the data directly on the tablet – this will then be uploaded to the primary database once the user is back in the office.

An initial pilot of 3 tablets will be used - once the method is validated the aim is to have 10 tablets in the field. The functionality will initially be limited to the users manually measuring data and then entering the data into the tablet – it is planned that later versions will allow input directly from electronic measuring boards.

It is expected that time will be saved by avoiding the need to transcribe data. Quality checks can also be applied automatically whilst the data is being entered – this can allow measurements to be double-checked if needed whilst the fish is still in front of the sampler. These benefits should mean the data collection is more efficient and higher quality.

(max 900 words per study)

Brief description of the results (including deviations from the plan and justifications as to why if this was the case).

An initial, mobile version of the “Stockman” database application was developed so that it could be run on rugged tablets when they were off-line. The data would then be securely uploaded to the laboratory database once the tablet had a network connection. Feedback from users was gathered and further refinements were made to the application. This work has proceeded more slowly than anticipated.

Achievement of the original expected outcomes of the study and justification if this was not the case.

The application and tablets were tested during 2022 and 2023 but were not fully rolled out. The Marine Institute had a number of updates to its IT security protocols during this period. A number of these impacted the method that the tablets were using to upload the collected data back to the laboratory database. This meant that the application’s upload functions had to be re-written a number of times thus causing a delay in the roll-out.

Incorporation of study results into regular sampling by the Member State.

A tablet has been sent to a port laboratory on January 2024 to act as a pilot. Following any changes and adaptations suggested during the pilot it is planned that the application will be rolled out to more ports during the year.

(max. 900 words per study)

Text Box 1b: Other data collection activities

General comment: Use this text box to provide information on other data collection activities that relate to your EMFAF operational programme and need to be included in the work plan and the annual report. Describe activities that are funded by the DCF but fulfil objectives under other EMFAF priorities, like marine knowledge, or activities funded by the DCF, but without a direct link to the EU MAP specific requirements or WP template tables, like freshwater fisheries. You can also include one-off specific studies for a particular end-user need that do not enter the regular data collection.

1. Aim of the data collection activity

Herring Genetics

At least two stocks of herring are known to mix in ICES Division 6a during times of the year when they are acoustically surveyed and persecuted by the fishery. This issue has caused problems with the assessments for the 6a South7bc and 6a North stocks, leading to a combined assessment following the 2015 ICES benchmark. This combined 6a7bc assessment falls into a lower ICES assessment category than the previous separate assessments and has no biological reference points.

Following a series of projects with EU, national and industry funding, a genetic approach to differentiate the main herring stocks mixing in this area has successfully been developed. As a results of this significant progress, the stocks are due to be benchmarked

in Q1 2022. Genetic sample collection has been in place on the relevant acoustic surveys since 2014, allowing a split survey time series that will be examined at the benchmark meeting. Mixed catches of herring from the 6a fisheries have however not been genetically sampled to date. Should the benchmark mandate that the commercial catches of herring also be genetically split, then a new data collection and analysis activity will be necessary.

2. Duration of the data collection activity

If mandated by the benchmark, the collection and analysis of herring genetic samples in 6a would likely begin in 2023. The duration of the data collection activity (sampling and analysis of acoustic survey and commercial catches) would be indefinite i.e. as long as the two 6a stock assessments remain separate.

3. Methodology and expected outcomes of the data collection activity

The tissue sampling methodology has been well established on herring acoustic surveys since 2014 and a specialised tool has been developed to reduce processing time. Genetic markers to differentiate spawning locations have been developed and proven by external project partners. Applying these methods to samples taken from commercial catches should be straightforward once a statistically robust catch sampling regime is applied.

The expected outcomes of this data collection activity are: facilitating the separate assessment of the two herring stocks in 6a, improving the accuracy of said assessments, and improving the sustainable exploitation of a number of herring populations in the area.

(max 900 words per activity)

Brief description of the results (including deviations from the plan and justifications as to why if this was the case).

The expected outcome was achieved. The ICES benchmark in February 2022 approved a separate assessment for herring in 6a South, 7.bc and the ICES advice for fishing opportunities in 2023 was based on the Category 3 data limited (MSY) approach, removing the need for a monitoring TAC in the area. This change was mostly made possible by the genetically-split acoustic survey biomass index. Due to the need for a split biomass index each assessment year, the benchmark meeting recommended the continuation of genetic sampling on future Malin Shelf Herring Acoustic Surveys (MSHAS).

Achievement of the original expected outcomes of the study and justification if this was not the case.

A trial genetic sampling programme for the commercial catch herring in 6.a South, conducted in Q4 2022 and Q1 2023, was successful. The genotyping results of this trial are being used to inform an appropriate design for future long-term genetic sampling of catches from this stock.

Incorporation of study results into regular sampling by the Member State.

From 2023 onwards, regular genetic sampling of a certain proportion of herring in the acoustic survey and commercial catch have been and will continue to be incorporated into the sampling plan. A multiyear contract for genotyping is now in place.

(max. 900 words per study)

General comment: Use this text box to provide information on other data collection activities that relate to your EMFAF operational programme and need to be included in the work plan and the annual report. Describe activities that are funded by the DCF but fulfil objectives under other EMFAF priorities, like marine knowledge, or activities funded by the DCF, but without a direct link to the EU MAP specific requirements or WP template tables, like freshwater fisheries. You can also include one-off specific studies for a particular end-user need that do not enter the regular data collection.

By-catch

1. Aim of the data collection activity

The coastal tangle / trammel net fishery targeting Spiny Lobster and Turbot is known to pose a significant risk of by-catch to certain PET species. Data on by-catch from this metier has been increased since 2017 through a combination of scientific observer and self-reporting of catch and by-catch composition. The programme continues under a new framework where the majority of the vessels in this metier are now contracted to the MI to provide data on catch and by-catch and to engage in a suite of scientific projects with the objective of taking direct actions to mitigate by-catch. These projects are funded by the EMFF and EMFAF programmes and significantly enhance by-catch data for high risk metiers.

2. Duration of the data collection activity

Data will be collected over a four-year period as part of the project running from 2021 to 2025.

3. Methodology and expected outcomes of the data collection activity

The data collection methodology will follow the protocols of the MI at-sea catch sampling programme for these gears. The expected outcomes of this data collection activity are to provide more information on the rates of by-catch and actions to mitigate.

(max 900 words per activity)

Brief description of the results (including deviations from the plan and justifications as to why if this was the case).

PET species data on by-catch from this metier on the south west coast of Ireland was obtained annually from 2017-2021 (EMFF) and will continue to end of 2025 (EMFAF) through a combination of scientific observer and self-reporting of catch and by-catch composition. These projects significantly enhance by-catch data for high risk metiers.

Achievement of the original expected outcomes of the study and justification if this was not the case.

Species catch composition and catch rates have been estimated from observer and skipper sampling data between 2017-2023 in the crayfish tangle net fishery off the south west coast of Ireland. Between 300 and 800 nm of net hauls have been observed annually.

In 2023 crayfish were, numerically, the most common species caught in tangle nets off the Kerry coast followed by spider crab and brown crab. Significant numbers of lobsters are caught. Commercial fish caught in tangle nets include turbot, monkfish, pollack and spurdog and thornback rays along with low numbers of spotted and blonde rays. Other non-commercial species are caught including protected species such as grey seal and critically endangered species such as angel shark, flapper skate, blue or common skate and white skate. Single captures of Risso's dolphin, Harbour Porpoise and Common Dolphin were recorded in the by-catch in 2022 and 2023.

Acoustic arrays have been deployed in the area to detect acoustically tagged endangered species to identify patterns of movement of these fish in relation to the distribution of tangle net activity

Incorporation of study results into regular sampling by the Member State.

Data will be collected over a four-year period as part of the project running from 2021 to 2025. The expected outcomes of this data collection activity are to provide more information on the rates of by-catch and actions to mitigate.

(max. 900 words per study)

General comment: Use this text box to provide information on other data collection activities that relate to your EMFAF operational programme and need to be included in the work plan and the annual report. Describe activities that are funded by the DCF but fulfil objectives under other EMFAF priorities, like marine knowledge, or activities funded by the DCF, but without a direct link to the EU MAP specific requirements or WP template tables, like freshwater fisheries. You can also include one-off specific studies for a particular end-user need that do not enter the regular data collection.

Project SecWeb

1. Aim of the data collection activity

Support the operation and functioning of the RCG's Secretariat for a fluent regional coordination of data collection activities

2. Duration of the data collection activity

01/01/2023 – 31/12/2025

3. Methodology and expected outcomes of the data collection activity

The Secretariat's organizational structured has been set up and pilot tested throughout SecWeb project. The key functions of the RCG's Secretariat have been determined in close

collaboration with all RCGs, in particular with RCG and Intersessional Subgroups (ISSGs) chairs. A business model has been developed. In addition, good practices in communication within and among the RCGs have been promoted and installed. The overall capacity to reach out to a wider public and increase the visibility of the work and output of the RCGs has been boosted with the development of a dedicated website and the consolidation of a visual identity.

RCG chairs and the RCG's network in general have acknowledged the added value of having an RCG's Secretariat to the overall aim of improving data collection activities. Based on SecWeb project outputs the proposed data collection activity will connect the whole RCG network and stakeholders to work together on common goals. The Secretariat provides fluent administrative and coordination support for more efficient regional coordination liberating national experts involved in data collection activities from heavy burden administrative tasks.

Overall expected outcomes

- A full-time dedicated Secretariat support service for the RCGs enables a consistent approach to administering RCG activities, facilitates communication, and enhances the intersessional work, supporting also the work of sub-groups.
- A dynamic and permanently updated website will be kept available including as features:
 - o Integration – allowing seamless synchronization with third-party information needs and requests.
 - o Responsive display – to serve content across multiple devices, screens, and browsers.
 - o User experience- maintaining a satisfactory user experience throughout the website sections.
 - o Accessibility – To any interested visitor in a user-friendly way across the website sections.
 - o Retention- keeping visitors coming back to the website.
 - o Links to relevant restricted access sites and virtual environments.
- The Visual identity for the RCGs is increasingly consolidated and visibility and understanding of the work by the RCGs is enhanced for the relevant stakeholder groups.
- A regularly updated Stakeholders' database improves the communication function among the RCGs' experts and the stakeholders' community.
- Internal communication protocols and help-desk in place makes it easier for any new comer to efficiently join, adopt responsibilities, and contribute to the RCGs objectives and work commitments.
- The public description of the secretariat functions, operational working protocols and commitments will build trust and enhance the whole network transparency and accountability.

(max 900 words per activity)

Brief description of the results (including deviations from the plan and justifications as to why if this was the case).

During 2023, the RCGs Secretariat services and tools were financed by a short-term/low value contract by the individual Member States (23 out of 26 MS) together with a European Commission service contract running from the end of August 2023 to end of February 2024. These contracts enabled the full-scale operation of the RCGs Secretariat

along the year (from March 2023 until Feb 2024).

Achievement of the original expected outcomes of the study and justification if this was not the case.

The pan regional ISSG National Correspondents continued the discussions on how to implement the long-term stabilization of the services and tools and ran several consultations with the MS. A main conclusion from these, is the fact that the administrative procedures at national level are a continuous and complex constraint for some of the MS.

Incorporation of study results into regular sampling by the Member State.

Further discussions on the long-term funding scenarios are planned during 2024. The objective is to develop and implement the long-term funding solution for the RCGs Secretariat and the tools, while simultaneously, a suitable approach needs to be found to keep the services and tools working continuously to guarantee the support to the RCGs and the LM.

(max. 900 words per study)

SECTION 2: BIOLOGICAL DATA

Text Box 2.1: List of required species/stocks

North-East Atlantic

General comment: This text box fulfils Article 5(2)(a), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II point 2.1(a) of the EU MAP Delegated Decision annex. This text box applies to the annual report and complements Table 2.1.

Deviations from the work plan

List the deviations (if any) in the achieved data collection (lengths only) compared to what was planned.

The general reasons for deviations from the work plan in terms of planned vs. achieved data collection should be summarised in this section, while detailed comments on deviations on particular species/stocks should be included in the 'AR comments' column in Table 2.1.

Sample numbers across all stocks in the North East Atlantic are presented in Table 2.1

Clupea harengus 7aN: The very sporadic and extremely low landings from this fishery – just 4.3t landed in 2023, make it extremely difficult to sample. Attention is focused instead on adequately sampling herring fisheries in areas 7aS, 7gh, 7jk and 6a, 7bc where the vast majority of Ireland's herring landings occur.

Dasyatis pastinaca and *Hexanchus griseus* were not sampled, as they were not

encountered during 119 sampling events completed on the recreational fisheries sampling programme.

Nephrops norvegicus in 5b & 6 were highlighted to be sampled for length in Ireland's WP however

Faroes Grounds (Division 27.5.b): There are no designated *Nephrops* Functional Units in Division 27.5.b or *Nephrops* fishing grounds regularly visited by the Irish fleet. Hence no targeted Irish *Nephrops* fishing takes in Division 27.5.b from which to collect samples.

Rockall, Northwest Coast of Scotland and North Ireland (Subarea 27.6)

- Northwest Coast of Scotland and North Ireland or as the West of Scotland (Division 27.6.a)
- Rockall (Division 27.6.b)

FU11 - no Irish landings were recorded in 2022 or 2023

FU13 - no Irish landings were recorded in 2022 or 2023

FU12 - 34 tonnes landed in 2023. These Irish landings from FU12 were not sampled owing to their relatively small volumes and irregularity of landings and therefore sampling opportunities. Attention was focused on adequately sampling the important Functional Units in area 7 where the vast majority of Irish fishing effort is focused, and this can be seen in Table 2.1 with 128,072 individuals measured over 210 sampling events.

Raja undulata: It is also not possible to sample where there are no landings, which was the case for *Raja undulata* in area 7b, 7j.

Actions to avoid deviations

Describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

All stocks selected for sampling in 2023 were sampled – bar where there were no landings or where they were simply not encountered, in spite of extensive sampling as outlined above. In the case of herring in 7aN, sampling effort was redirected to ensure representative sampling of the bulk of the landings was achieved. No additional actions are needed.

(One text box of max. 1 000 words per region/RFMO/RFO/IO)

Text Box 2.1: List of required species/stocks

North Sea and Eastern Arctic

General comment: This text box fulfils Article 5(2)(a), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II point 2.1(a) of the EU MAP Delegated Decision annex. This text box applies to

<i>the annual report and complements Table 2.1.</i>
Deviations from the work plan <i>List the deviations (if any) in the achieved data collection (lengths only) compared to what was planned.</i> <i>The general reasons for deviations from the work plan in terms of planned vs. achieved data collection should be summarised in this section, while detailed comments on deviations on particular species/stocks should be included in the 'AR comments' column in Table 2.1.</i> NA Actions to avoid deviations <i>Describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.</i> All stocks selected for sampling in 2023 were sampled – bar where they were not encountered or there were no landings into Ireland, therefore no actions are needed. <i>(One text box of max. 1 000 words per region/RFMO/RFO/IO)</i>

Text Box 2.1: List of required species/stocks

Other Regions

<i>General comment: This text box fulfils Article 5(2)(a), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II point 2.1(a) of the EU MAP Delegated Decision annex. This text box applies to the annual report and complements Table 2.1.</i>
Deviations from the work plan <i>List the deviations (if any) in the achieved data collection (lengths only) compared to what was planned.</i> <i>The general reasons for deviations from the work plan in terms of planned vs. achieved data collection should be summarised in this section, while detailed comments on deviations on particular species/stocks should be included in the 'AR comments' column in Table 2.1.</i> NA Actions to avoid deviations <i>Describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.</i> NA <i>(One text box of max. 1 000 words per region/RFMO/RFO/IO)</i>

Text Box 2.2: Planning of sampling for biological variables

North-East Atlantic

General comment: This text box fulfils Article 5(2)(a), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II point 2.1(a) of the EU MAP Delegated Decision annex. This text box applies to the annual report and complements Table 2.2.

Deviations from the work plan

List the deviations (if any) in the achieved collection of biological data (other than lengths), compared to what was planned.

The general reasons for deviations from the work plan in terms of planned vs. achieved data collection should be summarised in this section, while detailed comments on deviations on particular species/stocks should be included in the 'AR comments' column in Table 2.2.

Demersal– at–sea:

Following the traumatic post Covid, post Brexit and fuel crisis years the Irish at sea sampling programme is starting to rebuild from a historic low. While 2023 showed a marked increase in achievement in the demersal at sea sampling programme, 100% of targets were not reached. In 2023 a significant proportion of the Irish fleet was decommissioned as part of the post Brexit restructuring which resulted in a lower number of potential participating vessels. The intention of removing these vessels was to increase the share of the available quota amongst those remaining in the fleet. Whilst there may be some increases, the overall feeling within the Industry, is that the trauma associated with the reduction of the fleet has yet to have the desired effect. The fleet is under enormous pressure from external factors associated with increased costs, tight margins and limited opportunities resulting in low morale. This low morale is having a knock on effect on our, at sea sampling, as many vessel owners are reluctant to engage with our programme.

Crustacea at-sea:

Cancer pagurus: No weights are collected at sea for brown crab, instead the focus is to collect weight data on – shore, where 3393 individuals were weighted.

Demersal – on – shore:

Gadus morhua 6b: Sampling Cod from this area was challenging, with just 14t landed in 2024.

Gadus morhua 7a: Sampling Cod from this area for maturity and sex was challenging,

with sporadic landings of just 32t across multiple ports in 2024

Leucoraja naevus 6, 7, 8ab: Just 16.7t of cuckoo ray were landed in very small quantities across multiple ports making it difficult to sample.

Molva molva are currently not aged by Ireland. 429 individuals were sampled for length and 424 for weight.

Raja brachyura: Only 2t of Blonde Ray were landed by Irish vessels from this area, and these landings were not encountered during on – shore sampling events.

Raja montagui: 6, 7b, 7j. Landings by the Irish fleet in 2023 were very low at just 2.6t, and were not encountered during on – shore sampling events.

Raja montagui: 7a, 7e-h Landings by the Irish fleet in 2023 were low at just 45t, and were not encountered during on – shore sampling events.

Pelagic - on shore.

Clupea harengus in area 7aN Extremely low landings from the Irish pelagic fleet in this area with just 4.3t landed in 2023, making it difficult to access samples.

Actions to avoid deviations.

Describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

Demersal– at–sea:

The Marine Institute has increased our stakeholder engagement via all avenues open to us. Through the Irish Fisheries Research Science Partnership (IFRSP) group and via direct meetings with fishermen pier side, at Co – Op and PO level we have emphasised the importance of ‘data from the deck’ as a critical input to ensure that the best available science is utilised in stock assessments. We have also worked in partnership with various Irish FIPS (Fishery Improvement Projects) where producers and processors work together to build on the sustainability of their operations. These stakeholder engagements have already had a positive impact on engagement with the observer programme and co-operation with the FIPs has also opened up access to new data sources.

The at sea sampling programme continues to be built on the multi-stream approach with both Sampler at Sea and At Sea Self Sampling data collection. The At Sea Self Sampling was originally designed as a Covid mitigation measure in the demersal fleet and has continued to be of great use, and is especially useful for vessels who cannot accommodate an observer due to space restrictions. In 2023 a tailored At Sea Self Sampling program for the pelagic fleet targeting Albacore Tuna fleet was launched. This programme facilitated increased coverage in a fishery where the logistics of getting samplers to sea are challenging and served to increase our sampling efforts especially in relation to PET

species. The offshore >10m static net fleet is still a challenge to sample due to limited accommodation. We are actively engaged with stakeholders in the fleet to design a suitable scheme which would help mitigate for this. As a separate mitigation we have undertaken additional day trips on static gear vessels which do not require overnight accommodation. Additional sampling at sea was achieved on the FU16 Porcupine Bank Nephrops directed fishery in 2023 as the state opened a limited fishery to a number of authorised vessels. This increased opportunity, resulted in additional sampling within this frame and has helped to improve the dataset for the summer months. The same is expected in 2024 under specific authorisation from the national authority.

As we have increased the number of projects we undertake aboard both nearshore and offshore fleets, we are careful to maximise the return for each engagement and collect the maximum amount of data from the deck. While 2023 was challenging already in 2024 an increased number of at sea sampling trips have been achieved in Q1 and the expectation is that this upward trajectory will continue throughout the year.

Crustacea at-sea:

Cancer pagurus: Irelands National Work Plan 2025 will be updated in 2024 to reflect that weight data is collected on shore and not at sea.

Demersal – on – shore:

Gadus morhua 6b: Cod 6b will continue to be sampled opportunistically as encountered during on – shore sampling events.

Gadus morhua 7a: will continue to be sampled when encountered through 4S sampling.

Leucoraja naevus 6, 7, 8ab: Cuckoo ray is always sampled when encountered during 4S sampling.

Molva molva are not aged by Ireland, and as the stock assessment is currently a length based assessment, we will continue to sample for other biological parameters but not age.

Raja brachyura: Blonde Ray landings are miniscule but will continue to be sampled on – shore when encountered.

Raja montagui: 6, 7b, 7j. will continue to be sampled opportunistically as encountered during on – shore sampling events.

Raja montagui: 7a, 7e-h will continue to be sampled opportunistically as encountered during on – shore sampling events.

Pelagic - on shore.

Clupea harengus in area 7aN will continue to be sampled opportunistically as encountered

during on – shore sampling events.

(One text box of max. 1 000 words per region/RFMO/RFO/IO)

Text Box 2.2: Planning of sampling for biological variables

North Sea and Eastern Arctic

General comment: This text box fulfils Article 5(2)(a), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II point 2.1(a) of the EU MAP Delegated Decision annex. This text box applies to the annual report and complements Table 2.2.

Deviations from the work plan

List the deviations (if any) in the achieved collection of biological data (other than lengths), compared to what was planned.

The general reasons for deviations from the work plan in terms of planned vs. achieved data collection should be summarised in this section, while detailed comments on deviations on particular species/stocks should be included in the ‘AR comments’ column in Table 2.2.

Clupea harengus areas I &II: Approx. 1,000t of herring were caught by the Irish fleet from this area and were landed into Norway.

Actions to avoid deviations.

Describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

Clupea harengus areas I &II: Landings from the herring fleet are monitored and landings into Irish ports are sampled.

(One text box of max. 1 000 words per region/RFMO/RFO/IO)

Text Box 2.2: Planning of sampling for biological variables

Other Regions

General comment: This text box fulfils Article 5(2)(a), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II point 2.1(a) of the EU MAP Delegated Decision annex. This text box applies to the annual report and complements Table 2.2.

Deviations from the work plan

List the deviations (if any) in the achieved collection of biological data (other than lengths), compared to what was planned.

The general reasons for deviations from the work plan in terms of planned vs. achieved data collection should be summarised in this section, while detailed comments on deviations on particular species/stocks should be included in the 'AR comments' column in Table 2.2.

Thunnus alalunga: The vast majority of landings from the Irish tuna fleet were made into France and Spain and are covered by existing bi – lateral agreements between Ireland and these two Member States.

Actions to avoid deviations.

Describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

NA

(One text box of max. 1 000 words per region/RFMO/RFO/IO)

Text Box 2.3: Diadromous species data collection in freshwater

General comment: This Textbox fulfils Article 5(2)(a), Article 6 (3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II, point 2.1(b) and point 2.3 of the EU MAP Delegated Decision annex. Use this text box to give an overview of the methodology used for the data collected from freshwater commercial fisheries for salmon, sea trout and eel, and from research surveys on salmon and sea trout in freshwater, and on eel in any relevant habitat including coastal waters.

MI/ESB Programmes: Text Box filled for Sampling Scheme Identifier: Diad_ESB_Eel, Diad_MI_Eel, Diad_MI_Salmon_SeaTrout, Diad_MI_Salmon_CWT and Diad_ESB_Counter

IFI Programmes: Text Box filled for Sampling Scheme Identifier: Wild Salmon and Sea Trout Tagging Scheme Recreational, Wild Salmon and Sea Trout Tagging Scheme commercial, Biological sampling NSIC, Biological sampling smolts NSIC

MI Programme (Including ESB)

Sampling Scheme Identifier: Diad_ESB_Counter

Counter Smolt Fixed permanent counter upstream monitor salmon and and kelt moving downstream, enabling full census on wild salmon and released reared salmon. These are situated at or closed to the tidal limit and are fixed to natural or hydropower barriers or weirs on the Shannon Ardnacrusha and Parteen (IE_Sha) and Erne (IE_NorW) and Clady. These counters provide a valuable time series of relative abundance of wild salmon smolts and released reared salmon smolts.

Use: Counters provide annual index recruitment abundance data for ICES WGNAS datacalls and in WGNAS assessment model.

Adult Fixed permanent counter downstream monitor adult salmon moving upstream, enabling full census on wild salmon and released reared salmon.

These are situated at or closed to the tidal limit and are fixed to natural or hydropower barriers or weirs on the Shannon Ardnacrusha and Parteen (IE_Sha) and Erne (IE_NorW) and Clady. These counters provide a valuable time series of relative abundance of wild salmon adults and released salmon adults.

Use: Counters provide annual index recruitment abundance data for ICES WGNAS datacalls and in WGNAS assessment model

Sampling Scheme Identifier: Diad_ESB_Eel

Glass Eel/Recruitment Traps Fixed permanent elver ladder traps monitor upstream recruiting juvenile eel. These are situated at or closed to the tidal limit and are fixed to natural or hydropower barriers or weirs on the Shannon Ardnacrusha and Parteen (IE_Sha) and Erne (IE_NorW). These traps provide a valuable time series of relative abundance of glass eel and young yellow eel recruits and are used in the annual ICES WGEEL stock assessment.

Use: Elver traps provide annual index recruitment abundance data for ICES WGEEL. Data are collected in biomass (kg) or where numbers are very low, counts are made and converted to biomass. For WGEEL datacalls, numbers are converted from biomass using site specific conversion factors related to size and age of recruits.

Silver Eel Escapement Methods: Data collected on silver eel conservation trap and Transport on the Erne (IE_NorW), Shannon (IE_Sha) and Liffey (IE_East). Eels are captured in the programme using location specific gear types, such as bridge mounted coghill nets, and or river anchored V-Wing Fykes. Programme to estimate silver eel production/escapement and to monitor downstream trap and transport of migrating silver eel using mark-recapture, DIDSON, hydrological profiles and assessment models. Additional sampling (length, silvering characteristics) undertaken at the points of capture.

Use: determines eel escapement. Used to set the levels of trap and transport of silver eel in the Erne, Shannon and Lee which is a management measure in the Irish EMP. Used in conjunction with T&T quantities to estimate silver eel production and escapement, Erne and Shannon are index rivers in the Irish model – IMESE)

The above described programmes contribute to the national eel monitoring programme (Eel: Council Regulation 1100/2007), which operate across different Irish agencies and

parent departments.

Sampling Scheme Identifier: Diad_MI_Eel

Glass Eel/Recruitment Traps Fixed permanent elver ladder traps monitor upstream recruiting juvenile eel. These are situated at or closed to the tidal limit and are fixed to natural or hydropower barriers or weirs on the Liffey (IE_East) and Burrishoole (IE_West). These traps provide a valuable time series of relative abundance of glass eel and young yellow eel recruits and are used in the annual ICES WGEEL stock assessment.

Use: Elver traps provide annual index recruitment abundance data for ICES WGEEL. Data are collected in biomass (kg) or where numbers are very low, counts are made and converted to biomass. For WGEEL datacalls, numbers are converted from biomass using site specific conversion factors related to size and age of recruits.

Yellow Eel Standing Stock Electrofishing and fyke net surveys Electrofishing river surveys and fyke net lake surveys on the Burrishoole Catchment (IE_West) target yellow eel in selected water bodies, all fish are identified; weight and length measurements taken.

Use: Estimating yellow eel (river, lake and transitional water) populations. Used in time series analysis in ICES WGEEL. May be used in future eel stock assessment modelling employing the French Eel Density Assessment (EDA) model. Outputs will be reported under EU Regulation metrics and also in ICES Datacalls

Silver Eel Production/Escapement Traps Permanent traps monitor downstream migrating silver eels on the Burrishoole River (IE_West) providing a full daily census.

Use: For estimating annual production and escapement of silver eel. Numbers of fish migrating downstream, daily number, size, weight and sex ratio of emigrating silver eels (used in the Irish model for estimating silver eel escapement - IMESE). Above described programmes contribute to the national eel monitoring programme (Eel: Council Regulation 1100/2007), which operates across different Irish agencies and parent departments.

Sampling Scheme Identifier: Diad_MI_Salmon_SeaTrout

Parr Electrofishing surveys Electrofishing (salmon, trout) surveys target juvenile salmon and trout in selected water bodies of the Burrishoole catchment (IE_West), all fish identified; weight and length measurements taken.

Use: Estimating juvenile salmon and trout (river) populations. Juvenile stocks linked to stock/recruitment data collected from the main census traps. Used in time series analysis in ICES WGTRUTTA.

Smolt Traps Permanent traps in Burrishoole (IE_West) monitor salmon and sea trout smolt (and kelt) moving downstream, enabling full census on wild salmon, released reared salmon and wild sea trout.

National Coded Wire Tagging Scheme Tags seaward migrating salmon smolts, detected upon river return as adults. Data include release and recovery locations, dates and sea age.

Use: Estimating survival/exploitation rates and straying of wild/hatchery salmon.

Adults Traps Permanent traps in Burrishoole (IE_West) monitor adult salmon and trout moving upstream from the sea, enabling full census on wild salmon, released reared salmon and wild sea trout.

Use: Estimating annual returns of adult salmon and recruitment of salmon and sea trout smolt. Numbers of fish upstream/downstream, daily number, size, weight and sex ratio of salmon, sea trout.

Data are provided in datacalls to ICES WGNAS and in annual data collation in ICES WGTRUTTA. Index data from Burrishoole are used to calibrate annual models and assessments, due to the long time series available.

Data also used by the National Technical Expert group on Salmon to calibrate the national assessments for setting Conservation Limits.

Sampling Scheme Identifier: Diad_MI_Salmon_CWT

National Coded Wire Tagging Scheme, Smolt and Adult life stages Tags seaward migrating salmon smolts, detected upon river return as adults. Data include release and recovery locations, length of tagged smolt, dates and sea age.

Tagging carried out on 7 rivers, Bundorragha river; Burrishoole river; Cong river ;Corrib river ;Erne river, Lee river and Shannon River.

Use: Estimating survival/exploitation rates and straying of wild/hatchery salmon.

Data are provided in data calls to ICES WGNAS. Index data from Burrishoole are used to calibrate annual models and assessments, due to the long time series available.

Data also used by the National Technical Expert group on Salmon to calibrate the national assessments for setting Conservation Limits.

Sampling Scheme Identifier: Eel_elver_trap_Diadromous (scientific)

Recruits: Time series data is required on number of recruiting eels (glass or elver) to Ireland. The elver traps are located at the high water mark capturing elvers as they migrate from transitional waters to freshwater. The trap is a unit of effort with catch per night recorded. Length data will be collected on specified dates to gather relevant biometry data for WGEEL data call and to supplement existing biometry data for the traps collected since 2009.

Sampling Scheme Identifier: Eel_Silver_Diadromous (scientific)

Silver Eel: Coghill nets set per night giving number of eels per survey. Number of nets are fixed for each site. Locations are former commercial fishing sites. Biometry data collected will include length, weight, eye measurements; where needed 50-100 eels will be taken back to laboratory for further analysis re sex determination, age, growth, parasite prevalence, intensity, swimbladder damage.

Sampling Scheme Identifier: Eel_Fykenet_Diadromous (scientific)

Yellow eel: Fyke Net Survey, nets set in standardised chains of 5 nets for 1 night giving a catch per unit of effort (net nights). Biometry data collected including length, weight, eye measurements. Standard Operating Procedure available on DCMAP Ireland websites. Biometry data collected will include length, weight, eye measurements; where needed 50-100 eels will be taken back to laboratory for further analysis re sex determination, age, growth, parasite prevalence, intensity, swimbladder damage.

**Sampling Scheme Identifier: - Wild Salmon and Sea Trout Tagging Scheme
Recreational**

Sampling scheme aiming at collecting annual catch quantities for *Salmo salar* in the freshwater part of their lifecycle as specified in Tables 1 and 4 of the EU MAP Delegated Decision annex and to provide data on fishing effort ;number and weight of all salmon caught separated by fisheries, location, age class with estimates also required for unreported catches; weight of ranched salmon caught; assignment to jurisdiction/region/river of origin of adult salmon; and sea age composition of returning adults; as specified by the RCG ISSG Diadromous Fishes.

The 5 Index rivers selected are Owenmore, Drowes, Mulkear, Laune, Slaney and Boyne.

Sea age will be determined based on date of capture or the relative proportion of stock that is one-sea-winter or multi-sea-winter proportion applied by the Technical Expert Group on Salmon. Unreported catch will be assumed be to 10% of reported raised recreational catch.

Sampling Scheme Identifier: Wild Salmon and Sea Trout Tagging Scheme commercial

These fisheries are primarily in single river estuaries (only three stocks i.e. Killary, Owenmore estuary and Castlemaine are mixed-stock estuary fisheries). As such this is considered to come under sampling scheme aiming at collecting annual catch quantities for *Salmo salar* in the freshwater part of their lifecycle as specified in Tables 1 and 4 of the EU MAP Delegated Decision annex and to provide data on fishing effort ;number and weight of all salmon caught separated by fisheries, location, age class with estimates also required for unreported catches; weight of ranched salmon caught; assignment to jurisdiction/region/river of origin of adult salmon; and sea age composition of returning adults; as specified by the RCG ISSG Diadromous Fishes.

There are no marine commercial or freshwater commercial fisheries for salmon in Ireland.

Locations include: Bandon; Barrow and Pollmounty; Belclare; Blackwater, Glenshelane, Finisk; Caragh; Castlemaine; Dawros; Eany; Feale, Galey and Brick; Glenamoy; Gweebarra; Ilen; Inny; Killary; Laune and Cottoners; Lower Lee (Cork); Maine; Moy; Newport; Nore; Owenduff; Owenea and Owentocker; Owenglin; Owenmore estuary; Roughty; Sheen; Sneem; Suir, Clodiagh, Lingaun, Blackwater; Waterville.

Sampling Scheme Identifier: Biological sampling NSIC

Length (cm), weight (kg) and age (scale sample as one-sea-winter or multi-sea-winter) are collected from a random sample of 100 adult Atlantic salmon per annum intercepted in the upstream fish trap at the National Salmonid Index Catchment River Erriff.

Sampling Scheme Identifier: Biological sampling smolts NSIC

Trapping facilities to capture and tag out-migrating smolts for DCMAP purposes at Tawnyard in the Erriff catchment during the smolt run in April to May each year. For the Erriff, total numbers can be determined, and age composition taken from scale samples of 100 smolts per species per annum.

(max 250 words per species and area)

Sampling Scheme Identifier: Diad_ESB_Counter

Were the planned numbers achieved? Yes/ No

Yes

If the answer is No, explain why not, and what measures were taken to avoid non-conformity.

(max 250 words per species and area)

NA

Sampling Scheme Identifier: Diad_ESB_Eel

Were the planned numbers achieved? Yes/ No

Diadromous Variable - Recruits: Yes, for all locations Erne, Shannon, and Lee

Diadromous Variable – Recruits Length: No, for Shannon, and Lee (Erne completed)

Diadromous Variable – Silver eel Escapement: No, for Erne and Shannon; only catch used, index sites not fished for assessment of escapement.

Diadromous Variable – Other: Yes, Shannon, 4 out of 5 for Erne

Diadromous Variable – Length: 3 out of 5 for Erne, No for Shannon

Diadromous Variable – Weight: No for Erne – only total weight, No for Shannon – only total weight

Diadromous Variable – Sex ratio: 3 out of 5 for Erne, No for Shannon

Diadromous Variable – Other: Yes, for Lee

Diadromous Variable – Length: No, for Lee

If the answer is No, explain why not, and what measures were taken to avoid non-conformity.

(max 250 words per species and area)

Recruits Length: Not sampled in Shannon. Measures taken: it is on the Agenda for the National Technical Eel Group

Silver eel Length/Weight/Sex ratio: Not sampled at 2 of the Erne locations and not sampled on the Shannon or Lee. Measures taken: The issue has been raised at the National Technical Eel Group and a letter went to the overall North South Standing Scientific Committee on Inland Fish (NSSSCIF) and to Inland Fisheries Ireland. No direct action has been taken yet, and for this reason, Ireland will not be in a position to determine silver eel production, escapement and mortality for 2023 when reporting to the EU in 2024.

Sampling Scheme Identifier: Diad_MI_Eel

Were the planned numbers achieved? Yes/ No

Diadromous Variable - Recruits: Yes, for all locations Burrishoole and Liffey

Diadromous Variable – Recruits Length: Yes, for all locations Burrishoole and Liffey

Diadromous Variables:

Standing stock	Fyke Net Survey	YES
Length	Fyke Net Survey	YES
Silver eel escapement	Permanent Trap	YES
Length	Permanent Trap	YES
Weight	Permanent Trap	YES
Sex ratio	Permanent Trap	YES
Standing stock	Electrofishing Survey	YES
Length	Electrofishing Survey	YES

If the answer is No, explain why not, and what measures were taken to avoid non-conformity.

(max 250 words per species and area)

NA

Diad_MI_Salmon_SeaTrout

Were the planned numbers achieved? Yes/ No

Diadromous Variables: Burrishoole Index – all YES

Standing stock	Electrofishing Survey	YES
Length	Electrofishing Survey	YES
Recruits	Permanent Trap	YES
Length	Permanent Trap	YES
Ascending individuals	Permanent Trap	YES
Length	Permanent Trap	YES

If the answer is No, explain why not, and what measures were taken to avoid non-conformity.

(max 250 words per species and area)

NA

Diad_MI_Salmon_CWT

Were the planned numbers achieved? Yes/ No

Yes 100% achieved

If the answer is No, explain why not, and what measures were taken to avoid non-conformity.

NA

(max 250 words per species and area)

Eel_elver_trap_Diadromous (scientific)

Were the planned numbers achieved? Yes/ No

100% achieved for the Corrib trap but under sampled for the Inagh trap.

If the answer is No, explain why not, and what measures were taken to avoid non-conformity.

(max 250 words per species and area)

For the Inagh trap excessive high water levels during the elver migration period meant the trap could not be serviced. The trap is located in the river and can only be monitored when conditions of low flow are experienced. Extensive rainfall for July and August impacted on the surveys. The Corrib elver trap is located in a fish ramp and is accessible under a wider range of conditions.

Eel_Silver_Diadromous (scientific)

Were the planned numbers achieved? Yes/ No

Yes for the Fane and mostly for the Barrow.

If the answer is No, explain why not, and what measures were taken to avoid non-conformity.

For the Barrow location 20 nights were fished, once consecutive zero catches are caught the fishing ceases for the month. Over the season only 20 nights were needed.

For the Fane 40 nights were fished due to the continuous low level of catches experienced during 2023 migration season.

(max 250 words per species and area)

Eel_Fykenet_Diadromous (scientific)

Were the planned numbers achieved? Yes/ No

No

If the answer is No, explain why not, and what measures were taken to avoid non-conformity.

The eel team in IFI consisted of only 1 full time person, therefore the extensive lake survey

could not be undertaken due to the lack of resources. Electrofishing of the River Erriff was carried out in 2023. A total of 28 sites were fished targeting European Eel. Only 8 eels were captured, with 11 eels observed but due to the rocky stream bed could not be retained. The average length of eel was 21.6cm with an average weight of 30g. Habitat assessments were carried out also in the catchment to investigate if the lack of eels was habitat related. The majority of sites were classified as high (50%) and Good (42%).

(max 250 words per species and area)

Wild Salmon and Sea Trout Tagging Scheme Recreational

Were the planned numbers achieved? Yes/ No

Mostly

If the answer is No, explain why not, and what measures were taken to avoid non-conformity.

Achievement of the planned numbers depend on: 1) how many salmon are caught by fisheries; and 2) whether the fishery is open for fishing or closed based on scientific and associated management advice for the particular stock. Therefore, in certain instances, it was not possible to achieve the minimum number of units in cases where the fishery was closed or less than the minimum number of units of fish were caught. Alternatively, when the catches were greater than the minimum number of units, the data is available and therefore, an excess of the minimum units is reported. In general, it should be noted that the units are identified as 'planned minimum number of units' and the WP comments in the excel file state "number will be dependent on returns from anglers".

(max 250 words per species and area)

Wild Salmon and Sea Trout Tagging Scheme commercial

Were the planned numbers achieved? Yes/ No

Mostly

If the answer is No, explain why not, and what measures were taken to avoid non-conformity.

Achievement of the planned numbers depend on: 1) how many salmon are caught by fisheries; and 2) whether the fishery is open for fishing or closed based on scientific and associated management advice for the particular stock. Therefore, in certain instances, it was not possible to achieve the minimum number of units in cases where the fishery was

closed or less than the minimum number of units of fish were caught. Alternatively, when the catches were greater than the minimum number of units, the data is available and therefore, an excess of the minimum units is reported. In general, it should be noted that the units are identified as ‘planned minimum number of units’ and the WP comments in the excel file state “number will be dependent on returns from fishers”.

(max 250 words per species and area)

Biological sampling NSIC

Were the planned numbers achieved? Yes/ No

Yes

If the answer is No, explain why not, and what measures were taken to avoid non-conformity.

NA

(max 250 words per species and area)

Biological sampling smolts NSIC

Were the planned numbers achieved? Yes/ No

Yes

If the answer is No, explain why not, and what measures were taken to avoid non-conformity.

NA

(max 250 words per species and area)

Text Box 2.4: Recreational Fisheries

North-East Atlantic

General comment: This text box fulfils Article 5(2)(a), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II point 2.2 of the EU MAP Delegated Decision annex. Use this text box to give an overview of the methodology used to collect data on marine and freshwater recreational catches. For freshwater diadromous species, use Table and Text Box 2.3.

Description of the sampling scheme/survey according to Table 2.4.

Fish catch rates from shore-based recreational angling (including beach, rock, pier and estuary) and inshore recreational angling (including private and rental boat, kayak and charter vessel) will be assessed as these components comprise virtually all marine recreational fishing (MRF) landings in Ireland.

Sampling methods, developed during the pilot study on MRF in Ireland (Ryan et al., 2021), have been refined and will be utilised to estimate area-specific catch rates on an annual basis. A multi-species sampling frame is being used which will account for all species listed in Table 4 of the EU MAP Delegated Decision annex, as well as all other angling species encountered in Ireland.

A combination of on-site (IMREC_CREEL, IMREC_OB_CH_SURVEY) and off-site (IMREC_ANG_DI, IMREC_SKP_DI) sampling programmes are being employed to estimate MRF catch rates in Ireland.

(max 900 words per region)

Deviations from the work plan

List the deviations (if any) in the achieved data collection, compared to what was planned in the work plan and explain the reasons for the deviations.

The angling diary (IMREC_ANG_DI) is active and diarists are submitting angling trips. However, potential bias is not well understood currently. Therefore, these data have not been employed to estimate catch yet. Catch estimates from the shore angling and small boat angling sectors were calculated from the on-site creel surveys (IMREC_CREEL).

Participation in the voluntary charter skipper diary (IMREC_SKP_DI) was low in 2023 and there were some concerns regarding data quality. Therefore, catch estimates from the charter sector were calculated from the onboard charter survey (IMREC_OB_CH_SURVEY) only in 2023.

Action to avoid deviations

Describe the actions that will be considered/have been taken to avoid the deviations in the future and when these actions are expected to produce results.

The angling diary (IMREC_ANG_DI) is a self-selecting (non-probabilistic) data collection programme. Therefore, care must be taken to ensure that catch rates reported are an appropriate reflection of the sector. During 2024 various statistical procedures, such as post-

stratification, are being assessed, to mitigate the effects of potential sampling bias in the diary panel. Also, a comparative analysis of on-site creel (IMREC_CREEL) versus off-site diary (IMREC_ANG_DI) sampling data is ongoing. The output of this work will provide insight about how best to combine the two data streams, so that precise estimates can be produced. The analysis is currently in progress. However, the timeframe for completion will be determined by the availability of resources. The outcome of the analysis will determine the level of confidence in the diary data for its potential use in catch estimates. It is anticipated that the diary data (IMREC_ANG_DI) will be comprehensively understood and ready for inclusion in catch estimates by 2025.

To encourage increased participation in the skipper diary in 2024 (IMREC_SKP_DI), the sampling team held regional workshops for charter skippers. These were held in several hubs around the country to these stakeholders, in March 2024. Participation was high and feedback was positive. Participation in the skipper diary is expected to increase considerably in 2024. Data collected through the skipper diary will be used in annual catch estimates for 2024.

(max 900 words per region)

Text Box 2.5: Sampling plan description for biological data

North-East Atlantic

General Comment: This text box fulfils Article 5(2)(a) and (b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter 2 point 2.1(a) of the EU MAP Delegated Decision annex. This text box complements Table 2.5.

Sampling scheme: Demersal at-sea, Demersal vessels – At-sea and self-sampling

The scheme covers all vessels >10m, all gears, that landed demersal species (including *Nephrops*) in the same quarter of the previous year. The scheme has two sampling components: At-sea sampling and self-sampling. At-sea sampling and self-sampling trips are based on the same sampling frame, which is a list of vessels, stratified into three geographic regions where the vessel mainly operates. PSUs are selected with unequal probability, based on past landings of demersal species. Currently Ireland has a limited at-sea sampling programme due to Covid 19 restrictions so where we are unable to get an on-board sampler, vessels are asked to undertake an at sea self-sampling trip. As restrictions ease and when safe to do so Ireland will endeavour to get more samplers to sea but will also continue with the at sea self-sampling scheme thus enabling two data streams to feed into the catch sampling programme. In future work programmes we will split out this scheme

into two separate schemes once a full at-sea programme resumes. Provision is made for monitoring bycatch, rare and VME species as part of this sampling scheme

Sampling scheme: Demersal at-sea Enhanced, Demersal vessels - Enhanced

In order to support the objectives of Natura, MSFD, OSPAR, and the CFP, further sampling of PET species bycatch will be undertaken as part of the ‘enhanced’ sampling programme to supplement bycatch sampling planned through the DCF sampling-at-sea scheme. The DCF demersal-at-sea sampling scheme covers all vessels >10m and all gears that land demersal species, and provision is made for monitoring bycatch, rare and VME species as part of this sampling scheme. The recommendations of STECF and WKPETSAMP included an increase in monitoring of metiers with a high risk of protected species bycatch. As such, based on a risk assessment of Irish fisheries, high risk metiers for the demersal fleet were identified and have been targeted by this enhanced sampling scheme. The enhanced scheme, in contrast to the regular DCF sampling scheme, targets demersal metiers deemed high risk to PETS following the results of this risk assessment. Specifically, the enhanced scheme covers all gillnet vessels >10m that landed demersal species. The enhanced sampling scheme will ensure sampling is undertaken on metiers deemed at high risk for bycatch of PETS, in addition to bycatch sampling concurrently carried out across all metiers. At-sea sampling are based on the same sampling frame, which is a list of gillnet vessels only, stratified into three geographic regions where the vessel mainly operates. PSUs are selected with unequal probability, based on past landings of demersal species. Currently Ireland has a limited at-sea sampling programme due to Covid 19 restrictions so where we are unable to get an on-board sampler, vessels are asked to undertake an at sea self-sampling trip. As restrictions ease and when safe to do so Ireland will endeavour to get more samplers to sea but will also continue with the at sea self-sampling scheme thus enabling two data streams to feed into the catch sampling programme.

Sampling scheme: Demersal on-shore, Main demersal ports

The scheme covers the top ports where 95% of the demersal landings (excluding *Nephrops*) take place. The scheme has a single sampling frame: Main Demersal Ports. The sampling frame is stratified into geographical areas for practical reasons. PSU targets are based on the demersal landings of the same quarter in the previous year. Provision is made for monitoring incidental bycatch and rare species as part of this sampling scheme.

Sampling scheme: Pelagic at-sea others, Pelagic vessels targeting Mackerel, horse mackerel, blue whiting and boarfish fisheries

All vessels >10m with average landings of Mackerel, Horse Mackerel, Blue Whiting and Boarfish combined of >10t in the most recent 3 years for the semester. PSUs are selected with unequal probability, based on average historic landings of the 4 species combined.

Provision is made for monitoring bycatch and rare species as part of this sampling scheme.

Sampling scheme: Pelagic at-sea Herring, Pelagic vessels targeting Herring

The target population is the group of vessels that are engaged in the Irish fisheries targeting the herring stocks of 6a.7bc (North West), Celtic Sea, and Irish Sea. All vessels that are authorised to fish for each herring stock in the sampling year are included in the sampling frame. The yearly lists of authorisations are compiled by the relevant section of the Department of Agriculture, Food and the Marine (DAFM) in conjunction with management advisory committees. The sampling frames are stratified by semester. The PSU is vessel*time. Vessels are sampled from the list with replacement. Sampling is carried out by a sampler on-board for the duration of the fishing trip. Provision is made for monitoring bycatch and rare species as part of this sampling scheme.

Sampling scheme: Pelagic at-sea enhanced, Pelagic vessels - Enhanced programme targeting horse mackerel, mackerel and blue whiting

In order to support the objectives of Natura, MSFD, OSPAR, and the CFP, further sampling of PET species bycatch will be undertaken as part of the ‘enhanced’ sampling programme to supplement bycatch sampling planned through the DCF sampling scheme. The DCF pelagic-at-sea sampling schemes cover all vessels >10m and all gears that land pelagic species such as tuna, mackerel, horse mackerel, herring, and blue whiting, and provision is made for monitoring bycatch, rare and VME species as part of this sampling scheme. The recommendations of STECF and WKPETSAMP included an increase in monitoring of metiers with a high risk of protected species bycatch. As such, based on a risk assessment of Irish fisheries, high risk metiers for the pelagic fleet were identified and have been targeted by this enhanced sampling scheme. The enhanced scheme, in contrast to the regular DCF sampling scheme, targets demersal metiers deemed high risk to PETS following results of this risk assessment. Specifically, the enhanced scheme covers all midwater otter trawl (OTM) and pelagic pair trawl (PTM) vessels >10m that landed horse mackerel, mackerel or blue whiting. The enhanced sampling scheme will ensure sampling is undertaken on metiers deemed at high risk for bycatch of PETS, in addition to generally bycatch sampling concurrently carried out across all metiers. At-sea sampling are based on the same sampling frame, which is a list of these high risk pelagic vessels only, stratified into three geographic regions where the vessel mainly operates. PSUs are selected with unequal probability, based on past landings of specific pelagic species. Currently Ireland has a limited at-sea sampling programme due to Covid 19 restrictions so where we are unable to get an on-board sampler, vessels are asked to undertake an at sea self-sampling trip. As restrictions ease and when safe to do so Ireland will endeavour to get more samplers to sea but will also continue with the at sea self-sampling scheme thus enabling two data streams to feed into the catch sampling programme.

Sampling scheme: Pelagic on-shore others, Pelagic ports landing Mackerel, Horse Mackerel, Blue Whiting

The sampling scheme covers landings and road transport of Mackerel, Horse Mackerel and Blue Whiting into the port of Killybegs for processing which accounts for over 95% of the total landings of these species into Ireland. The fisheries for these species commence at the start of the year and continue sequentially (Horse Mackerel, Mackerel and Blue Whiting) with some overlap for approximately 15 weeks. A second fishery for Mackerel and Horse Mackerel takes place in the fourth quarter for 6-8 weeks. The sampling frame is stratified by species, ICES division and vessel licence type (full pelagic or polyvalent). PSUs from vessels operating as pairs are combined. In the event the number of landings exceeds a weekly threshold, PSUs are selected at random.

Sampling scheme: Pelagic on-shore Boarfish, Boarfish Self sampling

A significant proportion of boarfish is landed into foreign ports and the fishery is prosecuted by a small number of vessels. A reference fleet (2 vessels) collect 2 samples per ICES division per week, freezes and stores for collection when returning to an Irish port.

Sampling scheme: Pelagic on-shore Herring, Pelagic vessels targeting Celtic Sea, NW, Irish Sea Herring

The sampling scheme covers all Irish pelagic and polyvalent vessels authorised to land herring in ICES areas 6 and 7. Both the Celtic Sea herring and 6a.7bc (North West) herring stocks are currently under monitoring TACs and sampling is therefore dictated by the needs to continue the respective scientific time-series. In the Celtic Sea, ICES has advised a target of 17 samples per year across the main fleet and sentinel fleet. To achieve this the PSU is haul * area or day where feasible. In 6a.7bc the PSU is fishing trip and all trips are to be sampled. The target of 25 samples has been set based on the number of trips in the preceding three years. Special arrangements are in place for both stocks to secure the necessary samples. The sampling scheme for Celtic Sea and 6a.7bc will need to be amended in order to be compliant with 4S if and when the stocks rebuild and support a full fishery. Pelagic on-shore sampling of Irish Sea herring is conducted on a random basis.

Sampling scheme: Pelagic on-shore Sprat, Pelagic vessels targeting Sprat

The targeted population is the commercial catch of Sprat by Irish pelagic vessels from all ICES areas. All vessels licenced for the fishery are included in the sampling frame, including a sizeable proportion of less than 10m vessels. The PSU is fishing trip*species. Irish sprat landings are highly variable year-to-year. The target number of PSUs therefore

needs to be flexible. In years with significantly higher sprat fishing effort, sampling intensity will be increased accordingly. In years with particularly low sprat fishing effort, it may not be possible to reach the sampling target.

Sampling scheme: *Nephrops* at sea self-sampling, *Nephrops* vessels in FU16

Further to the designated **Demersal at-sea** (detailed above) sampling targeting dedicated *Nephrops* fishing vessels takes place for Functional Unit (FU) 16 - Porcupine Bank *Nephrops* Ground. This is in order to ensure adequate sampling occurs to gather size measurements (carapace length) for the ICES FU16 assessment to be undertaken. The scheme covers all vessels >10m that target *Nephrops* in FU 16. These vessels pack and freeze catch on-board, reducing possibility for on-shore landings sampling. The scheme operates self-sampling programme. Vessels that have historically reported *Nephrops* landings are included in the quasi-reference fleet. Each undertakes on average 3 to 5 fishing trips per year to FU16. The quasi-reference fleets are not stratified owing to their low number and are sampled as and when availability allows according to reasonable logistics and constraints. The FU16, Porcupine Bank *Nephrops* Ground fishery is closed during summer months: by EU Regulation for three months (May 1st – July 31st) from 2010 to 2012, reducing to may since 2013, however Irish national restrictions continue to close the ground between the end of May and the end of September, this requires sampling to be aligned to available opportunities. PSUs are selected with unequal probability, based on past landings. Currently Ireland has a limited at-sea sampling programme due to Covid 19 restrictions so where we are unable to get an on-board sampler, vessels are asked to undertake an at sea self-sampling trip. As restrictions ease and when safe to do so Ireland will endeavour to get more samplers to sea but will also continue with the at sea self-sampling scheme thus enabling two data streams to feed into the catch sampling programme. Provision is made for monitoring bycatch, rare and VME species as part of this sampling scheme.

Sampling scheme: *Nephrops* at-sea, *Nephrops* vessels in FU16

Further to the designated **Demersal at-sea, Demersal vessels – At-sea and self-sampling** (detailed above) sampling targeting dedicated *Nephrops* fishing vessels takes place for Functional Unit (FU) 16 - Porcupine Bank *Nephrops* Ground. This is in order to ensure adequate sampling occurs to gather size measurements (carapace length) for the ICES FU16 assessment to be undertaken. The scheme covers all vessels >10m that target *Nephrops* in FU 16. These vessels pack and freeze catch on-board, reducing possibility for on-shore landings sampling. The scheme operates At-sea sampling. Vessels that have historically reported *Nephrops* landings are included in the quasi-reference fleet. Each undertakes on average 3 to 5 fishing trips per year to FU16. The quasi-reference fleets are not stratified owing to their low number and are sampled as and when availability allows according to reasonable logistics and constraints. The FU16, Porcupine Bank *Nephrops* Ground fishery is closed during summer months: by EU Regulation for three months (May 1st – July 31st) from 2010 to 2012, reducing to may since 2013, however Irish national restrictions continue

to close the ground between the end of May and the end of September, this requires sampling to be aligned to available opportunities. PSUs are selected with unequal probability, based on past landings. Provision is made for monitoring bycatch, rare and VME species as part of this sampling scheme.

Sampling scheme: *Nephrops* on shore, *Nephrops* vessels excluding FU16

All commercial catch fractions from the *Nephrops* fisheries landed into Ireland, from the primary *Nephrops* Functional Units (FU) fished by Irish registered vessels (or for fisheries where a bi-lateral agreement is in place).

Population sampled: 98% of the *Nephrops* landings are covered by the sampling program. All vessel classes > 10 metres and only *Nephrops norvegicus* are included in the sampling program. FUs 11 to 14 are not routinely sampled owing to low levels of national participation in these fisheries.

Stratification: Sampling events are stratified by FU/vessel/year/month. *Nephrops* grounds are geo-referenced by FU (FU15, FU17, FU19, FUs20 and 21 combined, and FU22).

Samples are brought ashore by fishers or by on-board samplers.

Sampling scheme: Crustacea at-sea, Potting Vessels

European lobster (*Homarus gammarus*) and Brown crab (*Cancer pagurus*) catches are sampled on board commercial vessels around the coast of Ireland in ICES areas 6 and 7 through an at sea sampling programme with sampling trips occurring on an ad-hoc basis during the 6-9 months that the fisheries take place.

Sampling scheme: Crustacea on-shore, Shellfish Co-ops and Processors

Landings, by Irish vessels, of the European lobster (*Homarus gammarus*) and Brown Crab (*Cancer pagurus*) are sampled monthly, where possible, at various processing facilities in the northwest, west and southwest of Ireland, during the 6-9 months of the fishing season.

Sampling scheme: Mollusc on-shore, Shellfish Processors

Landings, by Irish vessels, of the King Scallop (*Pecten maximus*) and Common whelk (*Buccinum undatum*) are sampled monthly, where possible. King Scallop are sampled at a processing facility in the southeast of Ireland, while Whelk are sampled at processing facilities in the northwest and southeast.

(One text box (max. 1 000 words) per region/RFMO/RFO/IO)

Deviations from the work plan

List deviations (if any) in the achieved data collection compared to what was planned in the work plan and

explain the reasons for the deviations.

Sampling scheme: Demersal at-sea, Demersal vessels - At-sea and self-sampling.

2023 remained challenging for the at sea and self-sampling programme mainly due to a variety of external extenuating circumstances affecting the availability of participating vessels. Morale in the fishing industry is at an all-time low following Brexit and the difficult trading conditions post Covid, particularly the high cost of fuel, and a decommissioning scheme. However, achievements in this frame increased by 50% compared with 2022 and it is hoped that this upward trajectory in achievement of targets will continue in 2024.

Sampling scheme: Demersal at-sea Enhanced, Demersal vessels – Enhanced

Sampling opportunities on the gillnet fleet were not achieved due to limited space aboard vessels to carry an observer on – board. The method of fishing does not lend itself to/effective self-sampling either, which has been a challenge. However, 10 vessels participated in the collection of catch and bycatch data relating to the crayfish fishery in the south west. These vessels completed a total of 310 trips, with 289 trips involving self-sampling and 21 with observers on board. The majority of these trips were on tangle net fishery which is a sub set of the gill net fishery.

Sampling scheme: Pelagic at-sea Others

Shortfall in achieved target was due to logistical challenges in 2023 in securing observer trips on the pelagic fleet.

Sampling scheme: Pelagic at-sea Herring, Pelagic vessels targeting Herring

Sampling opportunities were limited by the fact that the fishery is confined to a scientific fishery with limited number of authorised vessels/trips. Extra sampling via at sea self-sampling increases the achieved sampling by 6 trips.

Actions to avoid deviations

Describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

Sampling scheme: Demersal at-sea, Demersal vessels - At-sea and self-sampling

Ireland has continued to Increase engagement with key stakeholders such as Fish Producer Organisations, Fishermen's Co-ops and the Irish FIPs to increase engagement with the catch sampling programme to secure additional at sea sampling trips. The adoption off a twin

track approach with both the at sea observers and the at sea self-sampling programme has helped to mitigate when it is not possible for certain vessels to accommodate an observer on board, resulting in additional data collection opportunities. Two additional personnel have been recruited, and have been fully trained and will now have a strong focus on completing at sea observer work in 2024, which should increase achieved targets across 2024 and beyond.

Sampling scheme: Demersal at-sea Enhanced, Demersal vessels – Enhanced

The Marine Institute is working collaboratively with the fleet to develop a suitable self-sampling programme. Day trips on <10m vessels also target this fishery and a number of trips were achieved on the <10m fleet.

Sampling scheme: Pelagic at-sea Others

Additional observers have been recruited and are now fully trained. One of these observers is located in Ireland's largest pelagic port, which should aid in the achievement of at sea sampling targets. We will continue to engage with key stakeholders such as Fish Producers Organisations and Processors on a local level to secure sampling opportunities.

Sampling scheme: Pelagic at-sea Herring, Pelagic vessels targeting Herring

The Marine Institute is working closely with the Competent Authority responsible for regulating this scientific fishery in relation to getting observers on board, and will continue to operate a self-sampling scheme in parallel to augment the observer at sea trips.

(One text box of max. 1 000 words per region/RFMO/RFO/IO)

Text Box 2.5: Sampling plan description for biological data

North Sea and Eastern Arctic

General Comment: This text box fulfils Article 5(2)(a) and (b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter 2 point 2.1(a) of the EU MAP Delegated Decision annex. This text box complements Table 2.5.

Sampling scheme: Pelagic on-shore Norwegian Spring Spawning Herring, Pelagic vessels targeting NSSH self-sampling

A significant proportion of Norwegian Spring Spawning Herring is landed into foreign ports and the fishery is prosecuted by a small number of vessels (approx. 7). Arrangements are made with randomly selected vessels for the collection and storage of samples by the vessel

crew. Samples are collected by scientific staff when the vessel returns to an Irish port.

Deviations from the work plan

List deviations (if any) in the achieved data collection compared to what was planned in the work plan and explain the reasons for the deviations.

Sampling scheme: Pelagic on-shore Norwegian Spring Spawning Herring, Pelagic vessels targeting NSSH self-sampling.

Only 1 sampling event was achieved as the Approx. 1,000t of *Clupea harengus* caught by Irish vessels from this area - were landed into Norway

Actions to avoid deviations

Describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

Activity and landings from this sampling frame will continue to be monitored and additional resources will be put in place to ensure sampling targets are achieved in 2024.

(One text box of max. 1 000 words per region/RFM/RFO/IO)

Text Box 2.5: Sampling plan description for biological data

Other Regions

General Comment: This text box fulfils Article 5(2)(a) and (b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter 2 point 2.1(a) of the EU MAP Delegated Decision annex. This text box complements Table 2.5.

Sampling scheme: Pelagic at-sea Tuna, Pelagic vessels targeting Albacore Tuna

Tuna fisheries covered by a Joint MI- Irish Tuna FIP (Fisheries Improvement Project) providing an at-sea sampler who collects data according to MI protocols. This allows a trained sampler to be embedded with the fleet when they fish south in the Bay of Biscay and land the catch into Spanish & French Ports. In recent years the majority of Irish caught Albacore has been landed directly into Spain & France and are covered by Bilateral agreements to sample. Provision is made for monitoring bycatch, and rare species as part of this sampling scheme.

Sampling scheme: Pelagic on-shore Tuna, pelagic vessels landing Tuna

The sampling scheme covers the landings by Pelagic vessels of albacore tuna into Irish ports. Over the past few years' landings into foreign ports has increased significantly and

this is now covered in a bi-lateral agreement with France and Spain.

Deviations from the work plan

List deviations (if any) in the achieved data collection compared to what was planned in the work plan and explain the reasons for the deviations.

Sampling scheme: Pelagic on-shore Tuna, pelagic vessels landing Tuna.

Tuna landings fluctuate significantly year on year, making planning sampling difficult. There was less activity in this frame in 2023, compared to the reference years, which is why sampling is lower. The vast majority of Ireland's tuna catches were landed into France and Spain in 2023. These landings are covered by bilateral agreements between Ireland and France and Ireland and Spain, which were updated and signed in December 2021 and March 2022 respectively and remain in force.

Actions to avoid deviations

Describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

Ireland will continue to maintain close contacts with colleagues in France and Spain, helping to track Irish landings into both countries. Additional personnel will be based in the main port into which tuna are landed in 2024, and this will help to ensure that tuna catches landed into Ireland are sampled adequately.

(One text box of max. 1 000 words per region/RFMO/RFO/IO)

Text Box 2.6: Research surveys at sea

International Ecosystem Survey in the Nordic Seas (ASH)

General Comment: This text box fulfils Article 5(1)(b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapters I and II of the EU MAP Implementing Decision annex. It is intended to specify which research surveys at sea, as set out in Table 2 of the EU MAP Implementing Decision annex will be carried out. Member States shall specify whether the research survey is included in Table 2 of the EU MAP Implementing Decision annex or whether it is an additional survey.

International Ecosystem Survey in the Nordic Seas (ASH)

1. Objectives of the survey

See other MS WP (Denmark) for full details on this survey

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

See other MS WP (Denmark) for full details on this survey

3. For internationally coordinated surveys, describe the participating Member States/vessels.

See other MS WP (Denmark) for full details on this survey

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

IRL participates by sending personnel and cost sharing as per RCG 2021 agreement

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

A cost sharing agreement is in place, to reimburse Denmark for their ship time at the relative share of their TAC. IRL participates by sending personnel and cost sharing as per latest multilateral agreement. Please refer to the Danish Annual Report for all details relating to this survey.

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

Irish Groundfish Survey (IBTS_Q4)

General Comment: This text box fulfils Article 5(1)(b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapters I and II of the EU MAP Implementing Decision annex. It is intended to specify which research surveys at sea, as set out in Table 2 of the EU MAP Implementing Decision annex will be carried out. Member States shall specify whether the research survey is included in Table 2 of the EU MAP Implementing Decision annex or whether it is an additional survey.

Name of the research survey

Irish Groundfish Survey (IBTS_Q4)

1. Objectives of the survey

The main objective of the IBTS_Q4 is to collect data on the distribution, relative abundance and biological parameters of commercial commercially exploited demersal species in 6a south, 7b & 7g-j north. The indices currently utilised by assessment WG's are for haddock, whiting, plaice, cod, hake and sole. Survey data is also provided for white &

black anglerfish, megrim, pollack, ling, blue whiting and a number of elasmobranchs as well as several pelagics (herring, horse mackerel and mackerel). Occurrence of vulnerable or sentinel invertebrate species such as corals, sea pen, fan mussel and ocean quahog is also noted. Marine litter is also sorted and recorded. Oceanographic data are collected from CTD instrument on trawl door and occasional surface to sea bed CTD transects. Sediment grabs are carried out opportunistically using a Day grab.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

Stations are randomly selected within a stratified survey area based on depth and historic analysis of survey catch distribution rates. Fishing is conducted using a GOV 36/47 trawl (20mm liner) with 5.3m² (1450 Kg) Morgere otter doors, 16" hoppers (D-gear) in area 6a and 8" disks (A-gear) areas 7b, g and j. The gear is trawled at 4kn for 30min at each station. Sweeps are 55m up to 80m depth, extended to 110m in deeper water to minimise variable trawl geometry.

All fish and invertebrate species are sorted and weighed. Biological data are collected for selected commercial demersal species such as Cod, Haddock and Whiting etc. Occurrence of vulnerable or sentinel invertebrate species such as corals, sea pen, fan mussel and ocean quahog is also noted. Litter is sorted and recorded. Oceanographic data are collected from CTD instrument on trawl door and occasional surface to sea bed CTD transects. Sediment grabs are carried out opportunistically using Day grab.

3. For internationally coordinated surveys, describe the participating Member States/vessels.

The Irish IBTS_Q4 Survey is carried out in formal collaboration with the other IBTS surveys in the area run by Scotland, Spain, France and Northern Ireland to a lesser degree. IBTS_Q4 Survey data from France and Ireland for cod, haddock and whiting are aggregated by Ireland to produce combined indices for stock assessment and advice at the ICES Working Group for the Celtic Seas Ecoregion (WGCSE). Several other single survey indices are also provided for WGCSE, WGBIE, WGCEPH and WGEF.

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

Not applicable.

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

The Irish Groundfish Survey is co-ordinated internationally by the International Bottom Trawl Survey Working Group (IBTSWG): **International Bottom Trawl Survey Working Group (IBTSWG) (figshare.com)**.

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer*

clearly to the annex and map numbers.

The Q4 Irish Groundfish Survey (IGFS) collects data on the distribution, relative abundance and biological parameters of commercially exploited demersal species in VIa south, VIIb & VIIg,j north. The indices currently utilised by assessment WG' s are for haddock, whiting, plaice, cod, hake and sole. Survey data is also provided for white & black anglerfish, megrim, pollack, ling, blue whiting and a number of elasmobranchs as well as several pelagic species (herring, horse mackerel and mackerel).

A number of ancillary objectives are also undertaken:

- a. To collect hydrographical information on the water column and the seafloor;
- b. To collect information on the composition and quantities of litter items on the sea floor, as caught in the fish trawl;
- c. To collect data on biology and distribution of important elasmobranch species through opportunistic tagging during the course of the survey.

A Survey Map illustrating achieved stations in 2023 is available in Annex 1.1 'Quality Report for Biological Sampling Scheme IBTS_Q4' Figure 1

7. Extended comments

Extended AR comments can be placed under this section.

Current IBTS Term of Reference to develop a new sampling trawl is being carried out, including sea trials, with Marine Science Scotland and IFREMER, Lorient. Work is being absorbed within existing programs during roll out of the proposed trawl to IBTS participants during 2023-4.

Recent benchmarks have seen a shift towards modelled Spatio-temporal survey indices such as those for cod, haddock and whiting in the Celtic Sea. This has seen demonstrable improvements in precision of the indices so far. In addition, much of the routine diagnostics available as a result has proved valuable in managing issues such as loss of days at sea due to poor weather.

(max. 450 words per survey)

International Blue Whiting Spawning Survey (IBWSS)

General Comment: This text box fulfils Article 5(1)(b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapters I and II of the EU MAP Implementing Decision annex. It is intended to specify which research surveys at sea, as set out in Table 2 of the EU MAP Implementing Decision annex will be carried out. Member States shall specify whether the research survey is included in Table 2 of the EU MAP Implementing Decision annex or whether it is an additional survey.

International Blue Whiting Spawning Survey (IBWSS)

1. Objectives of the survey

- The primary aim of the International blue whiting spawning stock survey is to determine the age stratified abundance and distribution of blue whiting (*Micromesistius poutassou*) using acoustic survey techniques
- Collect hydrographic data by means of vertical CTD profiles
- Conduct directed trawl sampling using a pelagic trawl to determine the biological profile of target species
- Conduct directed trawl sampling using a pelagic trawl to determine the species composition of mesopelagic fish echotraces
- Conduct visual abundance surveys of marine mammals and seabirds

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

The Manual for International Pelagic Surveys (IPS) describes the methods used for survey design, analysis and reporting of survey data (SISP #9) and is described in detail in the latest IBWSS survey report (<http://hdl.handle.net/10793/1689>)

3. For internationally coordinated surveys, describe the participating Member States/vessels.

This survey acoustically measures the size of the spawning stock of blue whiting (*Micromesistius poutassou*) in western waters and is conducted by vessels from Ireland (RV *Celtic Explorer*), the Faroe Islands (RV *Jákup Sverri*), the Netherlands (RV *Tridens*), Norway (FV *Vendla*) and Spain (RV *Vizconde de Eza*).

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

A Danish scientist from DTUAqua participates in the IBWSS each year onboard the RV *Celtic Explorer* for the full duration of the survey (21 days).

A cost sharing agreement is in place, to reimburse Ireland and Netherlands for their ship time at the relative share of their TAC. Participating Member States for the blue whiting survey in 2022 are Denmark, Germany, Netherlands, Ireland, France and Sweden. Spain will provide ship time on its own vessel.

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

Working Group of International Pelagic Surveys (WGIPS) (figshare.com)

The International Blue Whiting Spawning Survey (IBWSS) is internationally coordinated

through
WGIPS:

National survey cruise report
<http://hdl.handle.net/10793/1862>

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

The International Blue Whiting Spawning stock survey was carried out over 17 days and thus within the recommended 21-day time window agreed by the group. Weather conditions were poor as compared to 2022, resulting in a delayed start of four days in the southern area by the Dutch and Irish vessels. The stock was regarded as suitably contained within the survey area. The total survey effort was comparable to survey effort in previous years. The survey in 2023 shows an 8% decrease in TSB and a corresponding 5% decrease in TSN when compared to the 2022 estimate for comparable survey effort and coverage. The estimated uncertainty around the total stock biomass was higher than in 2021, CV=0.16 compared to 0.19.

In terms of abundance, 3-year-olds (2020 year-class) were most abundant (49%), followed by the 2-year-olds (27%), 4-year-olds (9%) and 5-year-olds (5%), respectively. Immature fish represented 5% of TSB and 7% of TSN. Over 87% of the 2-year old fish were mature contributing to the SSB of the stock. The abundance of these two year classes, 2 & 3-year-old fish, in 2023 was the highest in the time series and above the numbers observed associated with the previous 2014 record year class.

Survey effort and coverage in the southernmost stratum remains an issue due to inconsistencies in the availability of the Spanish vessel from year to year. By not having a consistent time window of availability that fits with the other survey participants means the validity of the data is in question. This remains an ongoing issue that does not allow for Spanish survey data to become integrated into the time series.

A Survey Map illustrating achieved stations in 2023 is available in Annex 1.1 'Quality Report for Biological Sampling Scheme IBWSS' Figure 2.

7. Extended comments

Extended AR comments can be placed under this section.

Due to equipment failure on the R.V. Celtic Explorer, fishing trawls were not possible for the duration of the Irish leg of the International Blue Whiting Acoustic Survey 2023. Acoustic data from the survey was combined with acoustic data from the other nations taking part in this internationally coordinated survey as per standard protocols. Biological data from the nearest hauls taken by the other vessels were used for the calculation of the age-stratified blue whiting biomass for the entire area. There was therefore no impact on

the quality of the combined spawning stock biomass index.

(max. 450 words per survey)

International Mackerel and Horse Mackerel Egg Survey (MEGS)

General Comment: This text box fulfils Article 5(1)(b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapters I and II of the EU MAP Implementing Decision annex. It is intended to specify which research surveys at sea, as set out in Table 2 of the EU MAP Implementing Decision annex will be carried out. Member States shall specify whether the research survey is included in Table 2 of the EU MAP Implementing Decision annex or whether it is an additional survey.

Name of the research survey

International mackerel and horse mackerel egg survey (MEGS)

1. Objectives of the survey

The main objective of the survey is to extract, identify and stage the development of mackerel and horse mackerel eggs collected from plankton samples. Samples are collected every ICES half statistical rectangle. A CTD is attached to the plankton sampler and information on temperature, salinity and sample depth is collected at each station. Gonad samples are also collected from female fish which are analysed for fecundity, batch fecundity, atresia and POF stage. These data are used to provide WGWIDE, the assessment group for widely distributed pelagic fish, with a spawning stock biomass, SSB, estimate for mackerel, and an egg production estimate for horse mackerel.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

The survey protocols have been published by ICES and can be accessed at [https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%205%20-%20WGMEGS%20Manual%20for%20AEPM%20and%20DEPM.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%205%20-%20WGMEGS%20Manual%20for%20AEPM%20and%20DEPM.pdf) and [https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%206%20-%20MEGS%20V1.3.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%206%20-%20MEGS%20V1.3.pdf)

3. For internationally coordinated surveys, describe the participating Member States/vessels.

IPMA Portugal – Vizconde de Eza
IEO Spain - Vizconde de Eza
AZTI Spain – Ramon Margalef
TI Germany – Walther Herwig
WMR Netherlands - Tridens
DTU Aqua Denmark - Dana
IMR Norway - Charter
FAMRI Faroes – Jakup Sverri
MSS Scotland – Scotia + Charter
CEFAS England – Cefas Endeavour
MI Ireland – Celtic Explorer

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

No thresholds apply this is a mandatory survey for the MS

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

The International mackerel and horse mackerel egg survey (MEGS) survey is carried out every three years, and was completed in 2022 and is next scheduled for 2025.

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

NA

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

Spawning/Pre – Spawning Herring/Boarfish Acoustic Survey (WESPAS_ IRL)

General Comment: This text box fulfils Article 5(1)(b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapters I and II of the EU MAP Implementing Decision annex. It is intended to specify which research surveys at sea, as set out in Table 2 of the EU MAP Implementing Decision annex will be carried out. Member States shall specify whether the research survey is included in Table 2 of the EU MAP Implementing Decision annex or whether it is an additional survey.

Spawning/pre-spawning herring/boarfish acoustic survey (WESPAS_ IRL)

1. Objectives of the survey

- The primary aim of the WESPAS_ IRL survey is to determine the age stratified abundance and distribution of herring (*Clupea harengus*), boarfish (*Capros aper*) and horse mackerel (*Trachurus trachurus*) using acoustic survey techniques
- Collect biological samples from directed trawling on fish echotraces to determine age structure and maturity state of standing stocks
- Conduct genetic sampling of individual herring within ICES divisions 6a and 7b, c for stock identification analysis
- Use vertical CTD casts to determine hydrographic conditions and the extent of shelf front regions
- Collect zooplankton samples using dedicated vertical trawls to determine biomass of zooplankton and the spatial extent of areas of concentration
- Conduct visual abundance surveys of marine mammals and seabirds

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

The Manual for International Pelagic Surveys (IPS) describes the methods used for survey

design, analysis and reporting of survey data (SISP #9) and is described in detail in the latest WESPAS survey report (<http://hdl.handle.net/10793/1659>).

3. For internationally coordinated surveys, describe the participating Member States/vessels.

Not applicable

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

Not applicable

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

The WESPAS survey is coordinated through WGIPS:

<https://www.ices.dk/community/groups/pages/wgips.aspx>

The latest WESPAS Cruise Report is available here:

<https://oar.marine.ie/handle/10793/1871>

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

The objectives of the survey were carried out as planned and timing was consistent with previous years. Survey coverage (area) and acoustic sampling (transect miles) were comparable to 2022. The number of trawls carried out (n=53) was greater than in 2022 (n=40).

The Malin Shelf Herring total stock biomass (TSB) was 205,156 t and total stock numbers (TSN) was 1,719 million. The spawning stock biomass (SSB) was 154,776 t and spawning stock numbers (SSN) was 1,091 million. The CV for the survey was 0.40. The 2023 survey estimate for Malin Shelf herring was dominated by 2-wr (36% TSB and 44% TSN) and 4-wr (24% TSB and 18% TSN). The third most dominate age group was 3-wr herring contributing 14% to the TSB and 12% to TSN. Combined these three age classes represented 74% of TSB and 74% of TSN. Maturity analysis of herring samples in 2023 indicated overall 75% of herring (TSB) were mature. In 2023, 90% of herring (TSB) were mature.

Boarfish distribution was similar to previous years in terms of latitudinal range. Total

biomass (TSB) increased by 16% and total abundance (TSN) decreased by 14% compared to observations in 2022. The 4-year age class dominated the 2023 estimate contributing 25% of TSB and 29.4% of TSN (Table 6). Ranked second and third were the 3-year old (16% TSB & 24.9% TSN) and 7-year old fish (14.1% TSB & 11.5% TSN) respectively. Ranked fourth is the 6-year old fish (12.2% TSB & 11.4% TSN). Combined, these three age classes represented 67% of TSB and 77% of TSN. Maturity analysis indicated over 99.6% of TSB and 99% of TSN was represented by mature fish.

The estimate of Celtic Sea (CS) herring TSB (total stock biomass) and relative abundance (TSN) estimates were 29,267 t and 241,762,500 individuals (CV 0.93) respectively. Four winter ring fish dominated the total estimate, representing 36.8% of TSB and 35.1% TSN. Five winter ring fish ranked second contributing 32.5% and 28.6% respectively. Ranked third were 2 winter ring fish (11% TSB & 16.8% TSN). Ranked fourth were 6 winter ring fish (10.9% TSB & 9.7% TSN). Immature fish represented 11.4% of the total abundance (TSN) and 8% of the total biomass (TSB) observed.

Horse mackerel TSB (total stock biomass) and abundance (TSN) estimates were 94,852.1 t and 342,763,900 individuals (CV 0.22) respectively. The 2023 estimate is almost 4 times greater in terms of biomass and abundance compared to 2022. The 9-year-old fish dominated this year's survey estimate representing 29% of TSB and 26% of TSN. Six-year-old fish ranked second representing 17% of TSB and 20% of TSN and 15-year-old fish ranked third (12% to TSB & 9% TSN). Combined these three age classes represented 58% of TSB and 54% of TSN. Maturity analysis of horse mackerel samples indicated 100% maturity.

The same vessel and sampling equipment (transducers and trawl) were used.

A Survey Map illustrating achieved sampling in 2023 is available in Annex 1.1 'Quality Report for Biological Sampling Scheme WESPAS' Figure 3.

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

Nephrops UWTV Survey (UWV 16 – 17, UWTV 19, UWTV 20 – 22)

General Comment: This text box fulfils Article 5(1)(b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapters I and II of the EU MAP Implementing Decision annex. It is intended to specify which research surveys at sea, as set out in Table 2 of the EU MAP Implementing Decision annex will be carried out. Member States shall specify whether the research survey is included in Table 2 of the EU MAP Implementing Decision annex or whether it is an additional survey.

Name of the research survey

Nephrops UWTV Survey (UWTV 16-17, UWTV 19, UWTV 20-22)

MS undertakes three UWTV surveys in any one year but they are described here as one as the areas surveyed can interchange during the survey periods depending on weather conditions.

1. Objectives of the survey

The main objective of the Irish *Nephrops* Underwater TV surveys is to obtain quality assured estimates of *Nephrops* burrow densities for the following Functional Units (FU): 16-17, 19, 20-22. Occurrence of vulnerable or sentinel invertebrate species such as soft corals, and sea pens is also noted. Litter is recorded.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

Stations are randomly selected within a stratified survey area sufficient to cover adequately the known spatial and bathymetric distributions for each stock and ensure a CV of less than 20% for the total abundance estimate as recommended by SGNEPS. This data is submitted to WGCSE on an annual basis to contribute to stock assessment and management advice and also to WGNEPS annually, for survey coordination and quality control purposes.

At each station the UWTV sledge is deployed to capture time referenced high-definition image data with field of view or 'FOV' of 1.03 metre. Vessel position (DGPS), depth and position of sledge using a USBL transponder are recorded every 3 seconds.

Occurrence of vulnerable or sentinel invertebrate species such as corals and sea pen is also noted. Litter is recorded. Oceanographic data are collected from a sledge mounted CTD instrument. Sediment grabs are carried out opportunistically using Day grab.

When time allows beam trawling is carried out to opportunistically sample *Nephrops* and macro benthos, where the aim is to carry out approximately 7 beam trawls randomly on FU 17 and FU 22 only.

Documentation is listed in the quality report.

3. For internationally coordinated surveys, describe the participating Member States/vessels.

Not applicable.

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

Not applicable.

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

ICES (2023). Working Group on *Nephrops* Surveys (WGNEPS; outputs from 2022 meeting). ICES Scientific Reports. Report. <https://doi.org/10.17895/ices.pub.22211161.v1>

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

Primary result is to obtain quality assured estimates of *Nephrops* burrow densities and abundance estimates in area 7 which includes the *Nephrops* Functional Units (FU16, FU17, FU 19, FU 20-21 combined and FU22). This data is supplied to the relevant ICES stock assessment working group (WGCSE) to provide catch options for these commercial stocks. Secondary results include: collection of presence- absence data on sea-pen species where this data has been supplied to the relevant ICES working group on deep-water ecology (WGDEC) and uploaded to the Vulnerable Marine Ecosystem international database; presence- absence data of other macro benthos species, fish species and trawl marks on the sea-bed.

CTD data is collected and viewed as an emerging time series which will be used for looking at inter-annual and longer term variability of bottom sea temperature around the coast of Ireland.

100% coverage of all the *Nephrops* grounds was achieved in 2023 for stock assessment purposes and provision of catch advice for FU16, FU17, FU 19, FU 20-21 combined and FU22.

A Survey Map illustrating achieved sampling in 2023 is available in Annex 1.1 ‘Quality Report for Biological Sampling Scheme *Nephrops* UWTV Survey’ Figure 4.

7. Extended comments

Extended AR comments can be placed under this section.

In addition, 33% of the FU 15 survey grid was completed as a collaboration between AFBI – Northern Ireland and MI Ireland.

(max. 450 words per survey)

Celtic Sea Herring Acoustic Survey (CSHAS_ IRL)

General Comment: This text box fulfils Article 5(1)(b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapters I and II of the EU MAP Implementing Decision annex. It is intended to specify which research surveys at sea, as set out in Table 2 of the EU MAP Implementing Decision annex will be carried out. Member States shall specify whether the research survey is included in Table 2 of the EU MAP Implementing Decision annex or whether it is an additional survey.

Celtic Sea herring acoustic survey (CSHAS_ IRL)

1. Objectives of the survey

- The primary aim of the CSHAS_ IRL is to determine the age stratified abundance and distribution of herring (*Clupea harengus*) and sprat (*Sprattus sprattus*) using acoustic survey techniques
- Collect biological samples from directed trawling on insonified fish echotraces to determine age structure and maturity state of the herring stock
- Determine estimates of biomass and abundance for sprat within the survey area
- Collect physical oceanography data from vertical profiles from a deployed sensor array
- Conduct visual abundance surveys of marine mammals and seabirds

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

The Manual for International Pelagic Surveys (IPS) describes the methods used for survey design, analysis and reporting of survey data (SISP #9) and is described in detail in the latest CSHAS survey report (<http://hdl.handle.net/10793/1664>).

3. For internationally coordinated surveys, describe the participating Member States/vessels.

Not applicable

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

Not applicable

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

The 2023 Celtic Sea Herring cruise report is available here:

<https://oar.marine.ie/handle/10793/52>

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

Overall, geographical coverage and acoustic sampling effort was in line with previous years. Good trawl and hydrographic sampling was achieved. Core replicate surveys (Pass 1

& 2) were completed as planned. Adaptive surveys were carried out inshore (n=2) and offshore waters (n=2). Immature herring (0-wr) were well represented in the wider survey area (Pass 1 & Pass 2 core surveys), albeit in low numbers, as part of mixed species aggregations dominated by sprat and other small pelagics, namely sardine and anchovy.

The 2023 TSB estimate (Pass 1: core stratum) was 1,773.4 t and 109 million individuals (CV 0.61) and was composed entirely of immature (0-wr fish). Since 2016, the survey estimate is reported from core stratum effort only.

One adaptive survey was carried out on a discreet high density aggregation of herring. Herring in this area were in close proximity to the seabed and were typical of observations over recent years. This offshore adaptive survey yielded a TSB was 9,254.3 t and TSN of 70,633.5 ('000s) individuals (CV 0.54). The age composition was dominated by 4-wr, followed by 5-wr, 6-wr and 3-wr fish by weight. The dominant 4-wr fish contributed 45.9% to the TSB and 46.6% to TSN, followed by 5-wr fish (43.1% TSB & 41.5% TSN), then 6-wr fish (6.5% TSB & 5.7% TSN) and 3-wr fish (3.7% TSB & 4.6% TSN). All fish were found to be mature individuals. The age profile of mature herring sampled during the adaptive survey was representative of those in landings data and from observations during the summer WESPAS survey (2023).

The biomass of sprat was lower than observed in 2022 with the centre of gravity of distribution in the eastern Celtic Sea both inshore and offshore. Sprat The Pass 1 represented a total biomass of 11,342 t and a total abundance of 2,727,229 ('000s) individuals (CV 0.25).

Anchovy followed a similar pattern of distribution to sprat. Pass 1 represented a total biomass of 9,770 t and a total abundance of 769,823 ('000s) individuals (CV 0.73).

Sardine were found only in inshore waters, following a similar pattern to 2022. Pass 1 represented a total biomass of 22,069 t and a total abundance of 767,452 ('000s) individuals (CV 0.93).

A Survey Map illustrating achieved sampling in 2023 is available in Annex 1.1 'Quality Report for Biological Sampling Scheme CSHAS_IRL' Figure 5.

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

Irish Anglerfish and Megrin Survey (IAMS_IRL)

General Comment: This text box fulfils Article 5(1)(b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapters I and II of the EU MAP Implementing Decision annex. It is intended to specify which research surveys at sea, as set out in Table 2 of the EU MAP Implementing Decision annex will be carried out. Member States shall specify whether the research survey is included in Table 2 of the EU MAP Implementing Decision annex or whether it is an additional survey.

Name of the research survey

Irish Anglerfish and Megrin Survey (IAMS_IRL)

1. Objectives of the survey

The main objective of the survey is to obtain biomass and abundance indices for anglerfish (*Lophius piscatorius* and *L. budegassa*) and megrim (*Lepidorhombus whiffiagonis* and *L. boscii*) in areas 6a (south of 58°N) and 7 (west of 8°W). Secondary objectives are to collect data on the distribution, relative abundance and biology of other commercial demersal species (cod, haddock, ling, plaice, sole, pollack, saithe, whiting, brill, hake, john dory, lemon sole, turbot, brill, blonde ray, cuckoo ray, common skate/flapper skate, spotted ray, thornback ray, spurdog, *Nephrops*). Occurrence of vulnerable or sentinel invertebrate species such as corals, sea pen, fan mussel and ocean quahog is also noted. Litter is also sorted and recorded.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

Stations are randomly selected within a stratified survey area based on depth and commercial catch rates of the target species. Fishing is conducted using a Jackson trawl with 5.45m² Thyboron Type 16 otter doors, 16"oppers, 19mm tickler chain and 100mm cod end mesh. The gear is trawled at 3kn for one hour at each station. The warp to depth ratio is 3:1 for depths up to 200m, and 2:1 plus 200m in deeper water. All fish and invertebrate species are sorted and weighed. Biological data are collected for selected commercial demersal species such as Cod, Haddock and Whiting etc. Occurrence of vulnerable or sentinel invertebrate species such as corals, sea pen, fan mussel and ocean quahog is also noted. Litter is also sorted and recorded. Oceanographic data are collected from CTD instrument on trawl door and occasional surface to sea bed CTD transects. Sediment grabs are carried out opportunistically using Day grab.

Documentation is listed in the quality report

3. For internationally coordinated surveys, describe the participating Member States/vessels.

Irish Anglerfish and Megrin Survey is carried out in informal collaboration with Marine Scotland's Scottish Anglerfish and Megrin Survey (SIAMISS) and uses the same gear and fishing practices. IAMS is limited to south of 58° North while SIAMISS extends north of this line. Survey data are shared between Ireland and Scotland for the purpose of stock assessment and advice at ICES Working Group for the Celtic Seas Ecoregion (WGCSE) and Working Group for the Bay of Biscay and the Iberian Waters Ecoregion (WGBIE).

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

Not applicable.

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

IAMS is coordinated with the Scottish Anglerfish and Megrim Survey (SIAMISS) as part of ICES International Bottom Trawl Survey Working Group (IBTSWG) and uses the same gear and fishing practices.

Cruise Report for IAMS 2023 is available to download at
<https://oar.marine.ie/handle/10793/1864>

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

Survey indices, length, age and maturity data are routinely used in various ICES stock assessment working groups such as WGCSE and WGBIE.

https://ices-library.figshare.com/articles/report/Working_Group_for_the_Celtic_Seas_Ecoregion_WGCSE_/22268980

https://ices-library.figshare.com/articles/report/Working_Group_for_the_Bay_of_Biscay_and_the_Iberian_Waters_Ecoregion_WGBIE_/23541168

Environmental data such as temperature, salinity and underway water quality parameters are provided to ICES Oceanographic database.

A Survey Map illustrating achieved sampling in 2023 is available in Annex 1.1 'Quality Report for Biological Sampling Scheme IAMS_IRL' Figure 6.

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

Cockle North Irish Sea Survey (CNIS)

General Comment: This text box fulfils Article 5(1)(b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapters I and II of the EU MAP Implementing Decision annex. It is intended to specify which research surveys at sea, as set out in Table 2 of the EU MAP Implementing Decision annex will be carried out. Member States shall specify whether the research survey is included in Table 2 of the EU MAP Implementing Decision annex or whether it is an additional survey.

Name of the research survey: **Cockle North Irish Sea (CNIS)**

1. Objectives of the survey

Estimation of biomass to provide catch advice. Habitat assessment and impact of fishery on designated bird populations.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

Stratified random. Scientific quadrat and rake sampling. Enumeration of target species and other characterising species of benthic habitat.

3. For internationally coordinated surveys, describe the participating Member States/vessels.

N/A

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

N/A

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

The Cockle North Irish Sea (CNIS) survey is not coordinated internationally. The CNIS survey report is available in the Shellfish Stocks and Fisheries Review 2023 publication

Shellfish Stocks and Fisheries Review 2023: an assessment of selected stocks
(marine.ie)

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators).

Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

The main use of the results of the survey are for abundance estimates to advise on a TAC for the annual cockle fishery and to monitor the effect of the fishery on the surrounding habit and its associated species.

The Cockle Irish Sea survey is undertaken to provide size distribution and to estimate a biomass of cockles (*Cerastoderma edule*) in Dundalk Bay to advise on the status of the cockle stock. The Dundalk Bay cockle survey consists of a 500 m² survey grid mapped over the intertidal sand flat. Each grid cell was divided into 400 sub-cells 25m² in area. A quadrat and rake-over sample is collected from 3 randomly selected sub-cells (each of which was 25 x 25m) in each grid cell. The same grid cells are sampled each year where possible.

Survey data is submitted routinely to the government department responsible for the management of inshore fisheries in Ireland, i.e. The Department of Agriculture, Food and the Marine (DAFM). Data is also made available to the National and Regional Inshore Fisheries Forums (NIF and RIFF's), at which marine agencies, DAFM, the minister, representatives of the inshore industry and other stakeholders participate.

Size and weight distribution is calculated: reflecting exploitation rate, growth, mortality and recruitment history of the stock.

Survey Maps illustrating achieved sampling in 2023 are available in Annex 1.1 'Quality Report for Biological Sampling Scheme CNIS_IRL' Figures 8 and 8.1.

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

Oyster West of Ireland (OWI)

General Comment: This text box fulfils Article 5(1)(b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapters I and II of the EU MAP Implementing Decision annex. It is intended to specify which research surveys at sea, as set out in Table 2 of the EU MAP Implementing Decision annex will be carried out. Member States shall specify whether the research survey is included in Table 2 of the EU MAP Implementing Decision annex or whether it is an additional survey.

Name of the research survey: **Oyster West Ireland (OWI)**

1. Objectives of the survey

Estimation of biomass to provide catch advice on *Ostrea edulis* and *Magallana gigas* and habitat assessment.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

Regular grid or random, oyster dredge hauls. Enumeration of target and by-catch species. Size distribution data.

No map available as locations are very local and dispersed.

3. For internationally coordinated surveys, describe the participating Member States/vessels.

N/A

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

N/A

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

This survey is not coordinated internationally.

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

The main use of the results of the survey are for abundance estimates of oyster in order to give advice on a TAC for each oyster bed and to monitor the associated habitat. The results of the surveys undertaken in 2023 can be found here **Shellfish Stocks and Fisheries Review 2023: an assessment of selected stocks (marine.ie)**

Survey Maps illustrating achieved sampling in 2023 are available in Annex 1.1 'Quality Report for Biological Sampling Scheme OWI_IRL' Figures 9 – 9.8

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

Razor Clam Irish Sea (RCIS)

General Comment: This text box fulfils Article 5(1)(b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapters I and II of the EU MAP Implementing Decision annex. It is intended to specify which research surveys at sea, as set out in Table 2 of the EU MAP Implementing Decision annex will be carried out. Member States shall specify whether the research survey is included in Table 2 of the EU MAP Implementing Decision annex or whether it is an additional survey.

Name of the research survey: **Razor Clam Irish Sea (RCIS)**

1. Objectives of the survey

Estimation of biomass to provide catch advice on *Ensis siliqua* in the north and south Irish Sea by dredge haul. Monitoring of benthic habitats in the fished area.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

Stratified random based on high frequency iVMS data (Figure X and XX). Hydraulic dredge used for fish hauls. Enumeration of target species and other bivalves caught as by-catch. Size distribution target species. Marine community assessment.

3. For internationally coordinated surveys, describe the participating Member States/vessels.

N/A

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

N/A

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

This survey is not coordinated internationally

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

The main use of the results of the survey are for abundance estimates to advise on an annual TAC for the fishery and to monitor species caught as by-catch along with the

surrounding habitat. The results of the 2023 survey can be found here **[Shellfish Stocks and Fisheries Review 2023: an assessment of selected stocks \(marine.ie\)](#)**

Survey Maps illustrating achieved sampling in 2023 are available in Annex 1.1 ‘Quality Report for Biological Sampling Scheme RCIS_IRL’ Figures 7 and 7.1.

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

Irish Coastal Ecosystem Survey

General Comment: This text box fulfils Article 5(1)(b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapters I and II of the EU MAP Implementing Decision annex. It is intended to specify which research surveys at sea, as set out in Table 2 of the EU MAP Implementing Decision annex will be carried out. Member States shall specify whether the research survey is included in Table 2 of the EU MAP Implementing Decision annex or whether it is an additional survey.

Irish Coastal Ecosystem Survey

1. Objectives of the survey

The main objective of this survey will be to provide a new time series of abundance and distribution for key stocks within recruitment/nursery areas along the south and west coast of Ireland. Including but not limited to cod, whiting, haddock, plaice, sole, lemon sole, turbot, flounder, John dory, skates, rays and other elasmobranchs. This standardised inshore survey is timed to observe juvenile fish abundances, and associated ecosystem information. Occurrence of vulnerable or sentinel invertebrate species such as corals, sea pen, fan mussel and ocean quahog is also noted. Marine litter is also sorted and recorded. Oceanographic data are collected from CTD instrument on trawl door.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

Stations are randomly selected within a stratified survey area based on depth and historic analysis of survey catch distribution rates. Fishing is conducted using an otter trawl with a lined cod end.

All fish and invertebrate species are sorted and weighed. Biological data are collected for selected commercial demersal species. Occurrence of vulnerable or sentinel invertebrate species such as corals, sea pen, fan mussel and ocean quahog is also noted. Marine litter is also sorted and recorded. Oceanographic data are collected from CTD instrument on trawl door.

3. For internationally coordinated surveys, describe the participating Member States/vessels.

Not currently applicable

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

Not applicable

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

This survey is not part of any international coordination group. The national report is being written up at the moment and should be available later in 2024.

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

100 tows were completed in 2023. There is currently no baseline data for these coastal areas, therefore this survey is providing essential information to key stakeholders nationally on the structure and functioning of these important coastal areas.

Once a survey time series is established this data will provide:

- Indices of juvenile abundance for key demersal species within the Celtic Sea. Preliminary results indicate that the survey provides a stronger signal for the juvenile component of some commercial demersal species such as whiting, then the offshore Irish Ground fish survey (figure 2, Annex 1).

- Abundance estimates for a diversity of pelagic, demersal and elasmobranch species, and epi-benthic communities (figure 3, Annex 1). In total 260 species were identified on the survey, providing an invaluable baseline of the communities in this area.

A Survey Map illustrating achieved sampling in 2023 is available in Annex 1.1 'Quality Report for Biological Sampling Scheme ICEcoS' Figure 10.

7. Extended comments

No comment

(max. 450 words per survey)

Scallop Survey (SS_IRL)

General Comment: This text box fulfils Article 5(1)(b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapters I and II of the EU MAP Implementing Decision annex. It is intended to specify which research surveys at sea, as set out in Table 2 of the EU MAP Implementing Decision annex will be carried out. Member States shall specify whether the research survey is included in Table 2 of the EU MAP Implementing Decision annex or whether it is an additional survey.

Scallop Survey (SS_IRL)

1. Objectives of the survey

Estimation of relative abundance, distribution, size and age structure of scallop (*Pecten maximus*, *Aequipecten opercularis*) in western Irish sea and eastern Celtic Sea. The surveys will align with similar surveys undertaken by Northern Ireland in the north Channel, by the isle of Man in central Irish Sea and by England and Wales in Cardigan Bay and Channel. By-catch in scallop dredges will be estimated

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey. Dredge survey on commercial contracted vessels. Stratified random design based on Emodnet or other (acoustic backscatter) sediment layers. Survey effort will depend on charter costs.

3. For internationally coordinated surveys, describe the participating Member States/vessels.

N/A

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

N/A

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

N/A

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

The main use of the survey results will be to gain an estimation of relative abundance,

distribution, size and age structure of King scallop (*Pecten maximus*) in western Irish sea and eastern Celtic Sea. A survey was undertaken across three king scallop beds in the western Irish Sea (Bray Inshore, Bray Offshore and east of Dundalk Bay) in 2023.

A total of 74 tows were undertaken with 7 x 0.75 m wide spring-loaded scallop dredges. In one area 2 x 0.75 m wide queen scallop dredges were used to increase catchability of smaller scallops but due to uncertainty in the efficiency of the queen scallop dredges 2 x 0.75 m scallop dredges with more teeth and a smaller mesh liner were used during the survey east of Dundalk Bay.

The results of this survey can be found here **[Shellfish Stocks and Fisheries Review 2023: an assessment of selected stocks \(marine.ie\)](#)**

Survey Maps illustrating achieved sampling in 2023 are available in Annex 1.1 ‘Quality Report for Biological Sampling Scheme SS_IRL’ Figures 11 – 11.5.

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

SECTION 3: FISHING ACTIVITY DATA

Text Box 3.1: Fishing activity variables data collection strategy

General comment: This text box fulfils Article 5 (2)(c), Article 6 (3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II point 3.1 of the EU MAP Delegated Decision annex. It is intended to describe the method used to derive estimates on representative samples where data are not to be recorded under the Control Regulation (EC) No 1224/2009 or where data collected under Regulation (EC) No 1224/2009 are not at the right aggregation level for the intended scientific use. Text Box 3.1 should be filled only in case complementary data collection is planned

Explain the reasons for implementing complementary data collection.

Information on the sampling schemes is available in the quality document (Annex 1.2). However, the Member State is invited to highlight additional information here on sampling schemes and sampling frames deemed necessary to understand the actual sampling design planned for the implementation year(s).

The Member State will continue to collect transversal data, on a daily basis, from vessels < 12 meters in length (LOA) in a national, Sentinel Vessel Programme (SVP). These data report all transversal variables. However, it is not yet possible to define quantitative targets for a sampling programme for transversal parameters within metiers containing an inshore component; specifically for vessels <10 metres LOA and where official declarations of their landings are not required. More specifically the total number of active/inactive vessels in the inshore component is not accurately known and the SVP data are not raised to metier level. Also the SVP does not sample all possible metiers in the <10m fleet.

Further details are in Annex 1.2 Fisheries – Sentinel Vessel Programme

(max. 900 words)

Deviations from the work plan

List the changes from the work plan (if any) and explain the reasons.

Planned sample rates across the SSCF were reviewed resulting in an increase of 11 vessels (17%) participating in the Sentinel Vessel Programme in 2023, representing certain metiers of Ireland's SSCF.

Table 3.1 indicates that Purse Seiners 0-< 10m* are a segment in the IRL fleet (but this does not appear in Table 5.1 Fleet Population). Whilst this may have been the case when the WP was originally drafted, this segment is no longer a feature of the fleet.

Actions to avoid deviations

Briefly describe the actions that will be considered / have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

The increase in vessels participating in the Sentinel Vessel Programme was a positive

deviation which resulted in increased responses and data from certain metiers of Ireland's SSCF. Annual reviews are carried out routinely to ensure the MS is continually striving to plug data gaps and improve data quality.

In 2023, Purse Seiners 0-< 10m* did not feature as a segment in the IRL fleet. The MS will monitor this annually and report data for this segment if and when vessels in this length category begin using this gear again.

(max. 900 words)

Text Box 3.2: Fishing activity variables data collection strategy (for inland eel commercial fisheries)

General comment: This text box fulfils Article 5(2)(c), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II point 3.2 of the EU MAP Delegated Decision annex. It is intended to describe the methods and data sources used to estimate fishing capacity, effort and landings data.

Describe data collection of inland eel commercial fisheries landings, effort and capacity. List or describe, for instance, the number of fishing entities, fishing methods, and the associated units used.

As part of the management measures in the Ireland Eel Management Plan, commercial and recreational eel fisheries were ceased in 2009 (& 2010 in the UK part of IE_NorW) and this will be continued for the period 2022-2027. All bycaught eel must be released.

A full description of the historical fishery is given in the Irish EMP.

(max. 900 words)

Deviations from the work plan

List the changes from the work plan (if any) and explain the reasons.

There were no changes to the work plan. The fisheries remained closed in 2024

Actions to avoid deviations

Briefly describe the actions that will be considered / have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

This is kept under review by the National Technical Group on Eel (TEGE).

(max. 900 words)

SECTION 4: IMPACT OF FISHERIES ON MARINE BIOLOGICAL RESOURCES

Text Box 4.2: Incidental catches of sensitive species

(Region/RFMO/RFO/IO: Please indicate per text box and update the table of contents)

General Comment: This text box fulfils Article 5(2)(a) and (b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter 2 point 4.1 of the EU-MAP Delegated Decision annex. This text box complements Table 2.5.

This text box is complementary to information on the sampling schemes provided in the quality document (Annex 1.1). It serves to highlight information on sampling schemes and sampling frames related to incidental catches of sensitive species.

Additional information on planning the observation of incidental catches of sensitive species (if already filled in in Annex 1.1, please indicate where it can be found):

- Annex 1.1 Demersal at-sea enhanced
- Annex 1.1 Pelagic at-sea enhanced

- Has an assessment of the relative risk of bycatch for the different gear types/metiers taken place and been taken into account for the sampling design?

An assessment of high risk fishing metiers for cetacean, seabird and other PET species was carried out by the Marine Institute. Two metiers, (1) gillnet fisheries targeting fish and crustaceans and (2) pelagic trawl fisheries (OTM and PTM) targeting horse mackerel, mackerel and blue whiting, were identified as being high risk for bycatch for the Irish fleet. These fisheries were therefore specifically targeted for the enhanced bycatch sampling, with an additional 8 PSU for the gillnet fishery, and an additional 2 PSU for the pelagic fisheries targeting horse mackerel, mackerel or blue whiting. These sampling PSU are for 2022, with the intention of increasing these PSU incrementally over the coming years as capacity allows.

- What are the gear types/metiers that present the highest risk of bycatch per species/taxa of PETS in a given region?

For the Irish fleet, two metiers, (1) gillnet fisheries and (2) pelagic trawl fisheries (OTM and PTM) targeting horse mackerel, mackerel and blue whiting, were identified as high risk for cetacean and other PET species based on the available scientific literature and on a risk assessment carried out by the Marine Institute.

- What are the methods to calculate the observation effort?

The sampling design and protocol are described in detail in Annex1.1 for demersal at-sea enhanced, and Annex1.1 Pelagic at-sea enhanced sampling.

- Does the sampling design and protocol follow the recommendations from relevant expert groups? Provide appropriate references. If there are no relevant expert groups, the design and protocol have to be explained in the text.

The sampling design and protocol are described in detail in Annex1.1 for demersal at-sea

enhanced, and Annex 1.1 Pelagic at-sea enhanced sampling.

Additional information on observer protocols (if already filled in in Annex 1.1, indicate where it can be found):

- Annex 1.1 Demersal at-sea enhanced
- Annex 1.1 Pelagic at-sea enhanced

- Does the on-board observer protocol contain a check for rare specimens in the catch at opening of the cod-end? If YES is the observer instructed to indicate if the cod-end was NOT checked in a haul?

Not relevant for gill net fishery

- In gill nets - and hook-and-line fisheries: does the on-board observer protocol instruct the observer to indicate how much of the hauling process has been observed for (large) incidental bycatches that slip out of the net?

All interactions with the sampled hauled nets are observed and recorded. Some hauls are not observed due to H&S rest allowances/guidance and are recorded as not observed.

- In large catches: does the protocol instruct to check for rare specimens during sorting of the catch (i.e. at conveyor belt)? Is the observer instructed to indicate what percentage of the sorting or hauling process has been checked at "haul level"?

For the gillnet fishery it is possible to observe the total catch, for the pelagic fisheries due to the nature of the pumping operations it is only possible to observe the catch in the bag alongside the vessel and in the sample obtained. Samplers are instructed to note any incidental by-catch caught during the whole hauling process.

Additional information on sampling schemes

Member State may add specific contextual information related to a region and the implementation year(s), for instance highlighting new developments not yet detailed in the quality document, regional adaptation and/or perspectives for the future. Insert the information under the same sampling scheme identifier as in Table 2.5.

None

Additional description on sampling frames

Member State may add complementary description to what includes the 'Sampling frame description' column of Table 2.5. Insert the information under the same identifier and name as in columns 'Sampling frame identifier' and 'Sampling frame description' of Table 2.5, and in the same order (Sampling frame identifier + Sampling frame description).

None

(One text box (max. 1 000 words) per region/RFMO/RFO/IO)

Results

Provide additional information, if available, in this text box. For example, summary information on the number of individuals recorded as bycaught per species, gear group and monitoring method with information about the state of the animals (i.e. were they released alive, dead, or collected for sampling).

From the 56 trips of the Demersal at Sea Enhanced and Pelagic at Sea enhanced trips 6 birds and 1 mammal were encountered and all were released alive. The majority of the trips were in the tangle net fishery a subset of the gillnet metier.

The standard at sea sampling trips across the full range of fleets sampled (non-enhanced) recorded zero catch of PET species.

Deviations from the work plan

The Member State shall list the deviations (if any) in the achieved data collection compared to what was planned in the work plan and explain the reasons for the deviations.

Not all targets could be reached for the gillnet fleet, as it proved difficult to find a vessel with suitable accommodation to carry a sampler at sea. Space is limited on many of these vessels and the current at sea self-sampling protocol is not suitable for the gillnet fishery. However, a number of trips on the tangle net fishery were carried out in the south west which is a sub set of the gill net fishery.

Extra trips were carried out on pelagic vessels which increased the sampling in this metier.

A newly developed At Sea Self sampling programme for the Irish Albacore Fishery also increased the sampled number of trips by a further 39 PSUs which greatly increases the coverage achieved.

Actions to avoid deviations

The Member State shall describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

The Marine Institute is actively working with gillnet fishermen to develop a sampling strategy that could potentially facilitate at sea self-sampling. This will need to be augmented/validated with observer at sea data. We are looking into a new project which will hopefully help in this regard.

It is feared though that the introduction of the bycatch only fishery for Pollack will reduce the possible number of gill net trips available as many of the smaller vessels traditionally target Pollack during the summer season, but every effort is being made to work with industry to ensure trips are achieved.

The increase in at sea self-sampling via the albacore fishery will be further developed during the 2024 season in order to expand the number of at sea observations. The at sea self-sampling during the albacore season is an introduction to the pelagic fleet to this type of sampling and one which we expect to expand to other pelagic fisheries in the future.

(One text box of max. 1 000 words per region/RFMO/RFO/IO)

Text Box 4.3: Fisheries impact on marine habitats

General comment: This text box fulfils Article 5 paragraph 2(a) and 2(b), Article 6 paragraph 3(a), 3(b) and 3(c) of Regulation (EU) 2017/1004 and Chapter 2, section 4.2 of the EU MAP Delegated Decision annex. It contains information on additional studies on the fisheries impact on marine habitats.

The Marine Institute will develop specific projects targeting areas and metiers of potential high risk (VME indicator spp.) and this will be done outside the DCF through EMFAF and are currently in development.

As part of the Marine Institutes survey programme, data on VME species is routinely collected during multiple surveys such as the Ground Fish Survey, the UWTV surveys and DCF surveys. Where possible data are also collected on board the at-sea sampling programme.

1. Aim of the study

2. Duration of the study

3. Methodology and expected outcomes of the study

(max 900 words per study)

Brief description of the results (including deviations from the plan and justifications as to why if this was the case).

In 2023, the Marine Institute collected data on Vulnerable Marine Ecosystems (VME) during multiple fisheries assessment focused surveys, including IAMS, IGFS, and the standalone 2023 PORC survey. Surveys recorded benthos recovered in fishing nets to the highest possible taxonomic level along with weight, abundance, spatial and temporal location, depth and survey strata. Processing of UWTV survey tracks included presence/absence records of key benthos. 420 observations were recorded across the surveys: 192 from the UWTV surveys; 171 from the IGFS; 29 from the IAMS survey; and 28 from PORC.

A subset of 149 VME observations were submitted in response to an ICES data call in support of the Working Group on Deep-water Ecology (WGDEC), for their work estimating and spatially mapping VMEs across the Northeast Atlantic shelf and were incorporated into the ICES VME database. The Marine institute data submission included 78 records from the UWTV survey; 27 from IGFS; 16 records from IAMS; and 28 from PROC.

Achievement of the original expected outcomes and justification if this was not the case.

Data collection of these operations have, and continue to be functional components of fisheries surveys, recording the benthos caught as “value-added” information of Marine Institute fisheries survey operations and towards marine science. Outcomes have, and continue, to provide in part toward the ICES VME data-call and the assessments of WGDEC. Data are also available to third parties for research purposes.

Work undertaken in 2020 provided a scientific review of existing benthic data derived from fisheries surveys (2003 to 2018), establishing both a summary of the existing data and a

benchmark from which to monitor changes in benthic community assemblages.

Follow-up to the activities (what are the next steps, how the results will be used).

The complexities involved in defining, identifying and mapping VMEs has lead the Marine Institute to recruited a Scientific and Technical Officer (STO) to this area. The STO is collaborating with other Marine Institute teams, ICES and other international marine science organisations to strengthen Marine Institute VME research and contributions. Work focuses on monitoring FEAS benthos data, definition, mapping and changes to benthic assemblages, development of indicators, VMEs, national and international legislation.

(max. 900 words per study)

SECTION 5: **Economic** AND SOCIAL DATA IN FISHERIES

Text Box 5.2: Economic and social variables for fisheries data collection

General comment: This Text box fulfils Article 5(2)(d), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004, and Chapter II point 5 of the EU MAP Delegated Decision annex. It is intended to specify data to be collected under Tables 7, 8 and 9 of the EU MAP Delegated Decision annex.

1. Description of clustering

Clustering is used for a small number of segments for Irish vessels. Of the 35 unique segmentations in the Irish fleet, 14 are cluster into larger segments to maintaining confidentiality. Clustering of these 14 segments is done by combining vessels segment by vessel lengths with similar segments using the same gear. Clustering takes place when a segment has fewer than 10 vessels. The majority of these segments which are clustered are non-important and in total make up 3% of the total registered fleet. Refer to Table 5.1 for fleet clustering details.

2. Description of activity indicator

The activity of the Irish Fleet is described as “NA” as Ireland is not using activity indicators.

3. Deviation from the RCG ECON (ex. PGECON) definitions

The only deviation taken by Ireland is to not use the PIM. This was omitted due to the collection of tangible assets from financial accounts which are used to report capital values in line with the EU MAP Guidance which states that MS may use ‘Alternative methods based on company surveys. These alternative methods may be used if the derived estimates reflect the actual definition of net capital stock (depreciated replacement value of the vessel including on-board equipment with a useful lifetime of more than one year)’.

(max. 900 words)

Deviations from the work plan

List the changes from the work plan (if any) and explain the reasons.

In 2023, there were 32 unique segmentations in the Irish fleet and eight segments were clustered resulting in 21 segments for reporting purposes. Clustering was essential when a segment had fewer than 10 vessels (i.e. combining vessels segment by vessel lengths with similar segments using the same gear into larger segments) in order to maintain confidentiality. Most of these segments which are clustered (i.e. DRB1012 *, DRB2440 *, FPO1218 *, HOK1012*, TBB2440 *, TMVL0010*, TMVL1012*) are minor players in the

national fleet and in total make up 3% of the total registered fleet.

IRL achieved higher response rates than planned for many variables from 13 segments representing both the LSF and SSCF (i.e., DFN1012, DRB1012, DRB2440, DTS1012, DTS1218, DTS1824, DTS 2440, HOK1012, FPO0010, TBB2440, TM1218, TM2440, TM40XX) as detailed in Table 5.2, this has improved data quality. In particular, the response rates from FPO0010 (vessels using pots and traps 0-10m) were exceptional this year with 284 surveys completed from a total frame population of 531 representing a response rate of 54%. The overall success in terms of higher response rates across the 13 segments can be attributed to several factors: (i) enhanced outreach to vessel owners about the importance of completing the annual economic survey, (ii) increased understanding and buy-in to the MS's new online survey portal, and (iii) completion of the annual economic survey is now a condition of payment to vessel owners under EMFAF funded-grant aid schemes.

Whilst IRL had considerable success with the above segments, as detailed in Table 5.2, lower response rates were achieved for six segments, four from the SSCF and two from the LSF (i.e. DRB0010, DTS0010, DFN0010, FPO1012, FPO1218, DFN1824). These SSCF segments are traditionally challenging in terms of data collection for the MS, despite outreach efforts in recent years.

Actions to avoid deviations

Briefly describe the actions that will be considered / have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

In an effort to mitigate deviations (both positive and negative), the MS will review planned sample rates for future data collection in line with actual response rates. If certain trends emerge over the next few years, planned sample rates will be revised accordingly.

For the segments detailed above that achieved lower response rates than expected, the MS will continue to explore options to improve the response rates through pro-active engagement (e.g. targeting meetings through the inshore forums for the SSCF segments) with the sector and raising awareness about the importance of providing data on an annual basis to close the current data gaps.

(max. 900 words)

SECTION 6: ECONOMIC AND SOCIAL DATA IN AQUACULTURE

Text Box 6.1: Economic and social variables for aquaculture data collection

General comment: This text box fulfils Article 5(2)(e), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004, and Chapter II point 6 of the EU MAP Delegated Decision annex. It is intended to specify data to be collected under Tables 10 and 11 of the EU MAP Delegated Decision annex.

1. Description of the threshold application

Please provide a percentage for the Member State's production from the latest EU aquaculture production reported to Eurostat. Describe and justify the applied threshold(s).

All census data is used to generate total and segmented estimations for both EUMAP and EUROSTAT, therefore 100%. In both cases, data is generated from all commercially active and licenced production units. Any discrepancy between the estimations for EUMAP and EUROSTAT are due to two factors; the significant time-lag between the two data submissions and the variation that persists between the segmentations of the two regulations. Nevertheless, the estimation differences for equivalent segments are insignificant.

To protect business-level anonymity, there is a threshold of a minimum four business units applied to the reporting of any data by segment. Segment population below this number requires that segment data is amalgamated with that of the most appropriate related segment or both are amalgamated into a new segment and the historical data of the two old segments is incorporated accordingly into subsequent time series uploads.

2. Deviation from the RCG ECON (ex. PGECON) definitions

Describe and justify any deviations from variable definitions as listed in 'EU MAP Guidance Document' in the DCF website.

There are no deviations from variable definitions for aquaculture.

Land-based shellfish and marine macroalgal production have a collective output of less than 5% of total national production. Previously, these were amalgamated as an all-catching segment 'Other Shellfish other' so as to provide an accurate estimation of overall national output. The growing interest in Seaweed culture prompted Ireland to try separating Macro-algal production into a dedicated segment but the production output and the number of business involved to date are too few to allow this. A number of new companies have entered the population, since 2018, and output is expected to grow. As these numbers are too low to report on for confidentiality and would be under threshold they have not being included in table 6.1. If a minimum number of businesses are detected in the future threshold B should apply to Macro-Algal and land-based shellfish segment data provision. The latter segment may be incorporated into the most appropriate larger shellfish segment, probably 'Oyster Other methods' (Oyster on trestles) as the units involved are mainly hatcheries that supply this segment.

(max. 900 words)

Deviations from the work plan

List the changes from the work plan (if any) and explain the reasons.

The sample size of the population derived from the annual economic survey for certain costs variables had initially been set in the WP at 25% for all segments. As detailed in Table 6.1,

the MS achieved higher response rates leading to a greater sample size for five segments (i.e. macroalgae, mussel, oyster, salmon, trout) for most employment and operating costs variables from the annual economic survey. As a result, the larger sample sizes made it possible to raise the data to population estimates.

The MS achieved lower response rates than planned for two economic variable groups (i.e. operating costs and financial income) for five segments (salmon, trout, mussel, macroalgae, oyster). There can be challenges accessing this data for these segments in certain years.

Actions to avoid deviations

Briefly describe the actions that will be considered / have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

For segments that achieved higher response rates this year for certain variables, the sample size has been revised to 33% (from 25%) for future data collection. This reflects a positive deviation and the ability of the MS to gather more data, enabling better estimates for the total segment populations.

For segments that achieved lower response rates than expected, the MS will explore options to improve the response rates through pro-active engagement with the sector and raising awareness about the importance of providing data on an annual basis to close the current data gaps. In addition, similar to the fisheries data collection approach (Text Box 5.2), the MS is working towards adding completion of the annual economic survey as a condition of payment under EMFAF funded grant aid schemes to help improve response rates in the aquaculture sector.

(max. 900 words)

SECTION 7: ECONOMIC AND SOCIAL DATA IN FISH PROCESSING

Text Box 7.1: Economic and social variables for fish processing data collection

General comment: This text box fulfils Article 5(2)(f), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004, and Chapter II point 7 of the EU MAP Delegated Decision annex.

1. The Member State should provide justification for complementary data collection for fish processing in addition to Eurostat data.

2. Deviation from the RCG ECON (ex. PGECON) definitions

Describe and justify any deviations from variable definitions as listed in 'EU MAP Guidance Document' in the DCF website.

(max. 900 words)

No Programme Submitted.

Deviations from the work plan

List the changes from work plan (if any) and explain the reasons.

Actions to avoid deviations

Briefly describe the actions that will be considered / have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

(max. 900 words)

ANNEX 1.1 - QUALITY REPORT FOR BIOLOGICAL DATA SAMPLING SCHEME

The quality report fulfils Article 6(3)(d) of Regulation (EU) 2017/1004. This document is intended to specify data to be collected under Chapter II, point 2 of the EU MAP Delegated Decision annex: Biological data on exploited biological resources caught by Union commercial and recreational fisheries.

Use this document to state whether documentation in the data collection process (design, sampling implementation, data capture, data storage, sample storage and data processing) exists and identify where this documentation can be found. Provide short descriptions where indicated, even if the documentation can be found in English. Names of sampling schemes and strata shall be identical to those in Tables 2.2, 2.3, 2.4, 2.5, 2.6 and 4.1 of the WP/AR. For quality information on scientific surveys, use the survey acronym as a sampling scheme identifier. For mandatory surveys, refer to Table 1 of the EU MAP Implementing Decision annex, see also MasterCodeList 'Mandatory survey at sea'.

Quality Reports Related to Tables 2.2 and 2.5 and Text Boxes 2.2 and 2.5

Sampling scheme identifier: Demersal at-sea.

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: Demersal at-sea
Sampling scheme type: Commercial fishing trip
Observation type: SciObsAtSea
Time period of validity: 2018-2027
Description of the population
Population targeted: The target population is the group of vessels that are engaged in demersal fisheries (i.e. catching of demersal fish and <i>Nephrops</i> using bottom contacting gears)
Population sampled: All vessels that reported demersal landings in the same quarter of the previous year are included in the sampling frame.
Stratification: The sampling frames are stratified by year/quarter and geographic region (i.e., vessels that mainly operate in area 6, 7a or 7b-k).
Sampling design and protocols
Sampling design description: The PSU is vessel*time. The sampling frame is a quarterly list of vessels that were active in the same quarter of the previous year using the gear types OTB, SSC, GNS and TBB and the target assemblages DEF and CRU (demersal fish and crustaceans). Each vessel has a sampling probability based on the demersal landings in the relevant quarter of the previous year. Vessels are sampled with replacement. Any new vessels will not be included in the current year. Vessels catching demersal fish with

other gears are also not included. The vessels are sampled using two methods 1) At-Sea Sampling with a sampler aboard the vessel 2) At-Sea Self Sampling where the vessel skipper & crew collect samples and associated data which is brought ashore for further analysis. Rare/incidental bycatch of fish species are checked during each sampling event. Any Bird/Mammal/Reptile/PET/Decomposed organism that comes in contact with the gear during fishing operation is also recorded. VME indicator species are noted if present in the random box of discards Vessels targeting <i>Nephrops</i> also bring in a sample of catch and discards to augment the <i>Nephrops</i> on shore sampling Is the sampling design compliant with the 4S principle?: Y Regional coordination: N Link to sampling design documentation: https://www.dcmapi-ireland.ie/sites/default/files/DCF_Files/Trip%20selection%20for%20the%20demersal%20observer%20programme%20v2.docx Compliance with international recommendations: Y. The sampling program has been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints. Link to sampling protocol documentation: https://www.dcmapi-ireland.ie/sites/default/files/DCF_Files/docs/At-Sea%20Sampling%20Manual_2020.pdf Compliance with international recommendations: Y. The sampling protocols have been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints.
Sampling implementation Recording of refusal rate: Y – Refusals and analysis of refusal rates are in accordance with guidance from SGPIDS, 2013. Monitoring of sampling progress within the sampling year: Sampling progress against targets is tracked within the database and reports of sampling targets and achievements are available to samples.
Data capture Means of data capture: Data is initially recorded on paper and then transferred to a database application as soon as possible after sampling. Fish lengths are measured using 1 metre ruler; fish weights are only

recorded for samples that are brought back to the lab.

Data capture documentation:

SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and reviewed and updated regularly. A recent copy of the SOP for completing data sheets is available at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx

Quality checks documentation: Y

Data Validation occurs within the entry screens of the database. Data Checking occurs with a Voice Reports software package that reads what has been entered in the database and allows the data enterer to check the records that are being read out against the records written on the datasheets. For information the Operations Guide for the voice reports software is available at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/TextToSpeech%20Operations%20Guide.doc

Data storage

National database: FEAS_DemDiscards

International database: Detailed data is submitted to RDB / RDBES. Raised data is also currently submitted to the ICES InterCatch database.

Quality checks and data validation documentation: Y

Data Validation occurs within the entry screens of the database. Data Checking occurs with a Voice Reports software package that reads what has been entered in the database and allows the data enterer to check the records that are being read out against the records written on the datasheets.

Quality control occurs after each trip has been entered. These reports look at data anomalies, raising factors, tow and length data etc. This identifies any anomalies which can be checked against the paper data sheets or with the observer. The reports are run in R using imbedded SQL queries and the results are generated by LaTeX as PDF documents.

Length weight validation occurs in an excel spreadsheet. and serves as method for age and length checking also.

Data is also checked during extractions for end-users such as ICES / European Commission. The checks used will depend on the use of the data.

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

Fish samples are processed in Marine Institute laboratories for length, weight and maturity analysis. Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for processing and analysis. These age structures are stored at Marine Institute premises for a period of months before preparation and age reading is carried out.

The datasheets are scanned and stored electronically, the original raw datasheets are kept in the regional laboratories and then transferred to headquarters where they are filed. Raw data sheets, duplicate otoliths and read otoliths are archived offsite in secure storage areas after three years. Sectioned otoliths and illicia are stored on site in specialised file storage units.

Data processing

Evaluation of data accuracy (bias and precision): Y

There is not a stand-alone evaluation of bias and precision of the data collected by this scheme, but it has been evaluated in a recent ICES benchmark:

http://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/BSG/2017/WKIrish/wkirish3_2017.pdf

Editing and imputation methods: Y

Where errors are identified, the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during transcription.

Imputation of unsampled domains of interest often done at the international level (e.g. ICES), rather than at the national level. However, in some cases, it is appropriate to merge minor domains with larger ones to reduce the burden on the end-user to deal with data gaps. This can include merging ICES divisions or metiers with similar fishing patterns. At a lower level, imputation may also take place in applying age-length keys where length classes for which no age data exists are imputed following the approach described by Gerritsen et al (2006)

Gerritsen, Hans D., David McGrath, and Colm Lordan. "A simple method for comparing age-length keys reveals significant regional differences within a single stock of haddock (*Melanogrammus aeglefinus*).¹" ICES Journal of Marine Science 63.6 (2006): 1096-1100.

Quality document associated to a dataset:

No DOI is currently created for the dataset since it is not publicly available.

R markdown documents recording the estimation and imputation steps performed for ICES data calls are available on request. The estimation is based on the COST R package. An example of the scripts are available at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Discard%20Data%20Extraction_example.rmd

Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g., in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No Deviations

Sampling Design and Protocols – No Deviations

No Deviations

Sampling Implementation

The target number of PSU was not achieved; this is mainly due to high level of refusal rates.

The morale of the Irish fishing fleet is an all-time low and may be affecting the refusal rates. The refusals apportioned to non-availability of accommodation aboard vessels is mitigated for by the development and roll out of At Sea Self Sampling. While the At Sea Self Sampling provides a subset of the data that would be acquired with a sampler aboard the combination of both help achieve a greater number of observations

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: Demersal at-sea enhanced.

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: Demersal at-sea enhanced
Sampling scheme type: Commercial fishing trip
Observation type: SciObsAtSea
Time period of validity: 2022-2027
Description of the population
Population targeted: The enhanced sampling specifically targets fisheries determined to be higher risk to bycatch. The target population is the group of vessels that are engaged in demersal gillnet fisheries specifically (i.e. catching of demersal fish using gillnet gears). Population sampled: All vessels that reported demersal gillnet landings in the same quarter of the previous year are included in the sampling frame.
Stratification: The sampling frames are stratified by year/quarter and geographic region (i.e., vessels that mainly operate in area 6, 7a or 7b-k).
Sampling design and protocols
Sampling design description: The PSU is vessel*time. The sampling frame is a quarterly list of gillnet vessels only that were active in the same quarter of the previous year using the gear type GNS and the target assemblages DEF and CRU (demersal fish and crustaceans). Each vessel has a sampling probability based on the demersal landings in the relevant quarter of the previous year. Vessels are sampled with replacement. Any new vessels will not be included in the current year. Vessels catching demersal fish or crustaceans with other gears are also not included. The vessels are sampled using two

methods 1) At-Sea Sampling with a sampler aboard the vessel, and if necessary 2) At-Sea Self Sampling where the vessel skipper & crew collect samples and associated data which is brought ashore for further analysis.

Rare/incidental bycatch of fish species are checked during each sampling event.

Any Bird/Mammal/Reptile/PET/Decomposed organism that comes in contact with the gear during fishing operation is also recorded. In addition, associated information on each bycatch event are recorded, including haul number, gear used, whether the animal(s) is released alive or deceased, number of animals, condition of the animal(s), sex, and length.

Is the sampling design compliant with the 4S principle?: Y

Regional coordination: N

Link to sampling design documentation:

https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/Trip%20selection%20for%20the%20demersal%20observer%20programme%20v2.docx

Compliance with international recommendations: Y. The enhanced bycatch sampling programme is designed to supplement the regular bycatch sampling occurring under DCF. This sampling program has been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints. The recommendations of STECF included an increase in monitoring of métiers with a high risk of protected species bycatch. As such, based on a risk assessment of Irish fisheries, high risk fisheries for the fleet were identified and have been targeted by this enhanced sampling scheme.

Link to sampling protocol documentation:

https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/At-Sea%20Sampling%20Manual_2020.pdf

Compliance with international

recommendations: Y. The sampling protocols have been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints.

Sampling implementation

Recording of refusal rate: Y – Refusals and analysis of refusal rates are in accordance with guidance from SGPIDS, 2013.

Monitoring of sampling progress within the sampling year: Sampling progress against targets is tracked within the database and reports of sampling targets and achievements are available to samples.

Data capture

Means of data capture:

Data on bycatch is initially recorded on paper and then transferred to a database application as soon as possible after sampling. All bird, mammal, reptile (fresh or decomposed) that comes into contact with the gear during fishing operations (either collected on deck or falling out

during hauling) is recorded. Information on the bycaught species is recorded in the data sheets, if no bycatch is recorded this information is also noted to ensure all true zero samples are recorded and clear. Information recorded includes, haul number, gear used, whether the animal(s) is released alive or deceased, number of animals, condition of the animal(s), sex, and length.

Data capture documentation:

A recent copy of the SOP for completing data sheets is available at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx

Quality checks documentation: Y

Data Validation occurs within the entry screens of the database. Data Checking occurs with a Voice Reports software package that reads what has been entered in the database and allows the data enterer to check the records that are being read out against the records written on the datasheets. For information the Operations Guide for the voice reports software is available at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/TextToSpeech%20Operations%20Guide.doc

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Data storage

National database: FEAS_DemDiscards, bycatch data are also summarised and stored in a bycatch specific database.–

International database: Detailed data is submitted to RDB / RDBES. Raised data is also currently submitted to the ICES InterCatch database.

Bycatch data on seabirds, cetaceans, pinnipeds and PET species fish are submitted to ICES WGBYC data call annually, and to any additional ICES or OSPAR special request data calls relating to bycatch, such as WKMOMA.

Quality checks and data validation documentation: Y

Data Validation occurs within the entry screens of the database. Data Checking occurs with a Voice Reports software package that reads what has been entered in the database and allows the data enterer to check the records that are being read out against the records written on the datasheets.

Quality control occurs after each trip has been entered. These reports look at data anomalies, raising factors, tow and length data etc. This identifies any anomalies which can be checked against the paper data sheets or with the observer. The reports are run in R using imbedded SQL queries and the results are generated by LaTeX as PDF documents.

Length weight validation occurs in an excel spreadsheet and serves as method for age and length checking also.

Data is also checked during extractions for end-users such as ICES / European Commission. The checks used will depend on the use of the data.




Sample storage
The data sheets are scanned and stored electronically, the original raw datasets are kept in the regional laboratories and then transferred to headquarters where they are filed. Raw data sheets, duplicate otoliths and read otoliths are archived offsite in secure storage areas after three years. Sectioned otoliths and illicia are stored on site in specialised file storage units. Any retained fish samples are processed in Marine Institute laboratories for length, weight and maturity analysis. Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for processing and analysis. These age structures are stored at Marine Institute premises for a period of months before preparation and age reading is carried out.
Data processing
Evaluation of data accuracy (bias and precision): Y There is not a stand-alone evaluation of bias and precision of the data collected by this scheme, but it has been evaluated in a recent ICES benchmark: http://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/BSG/2017/WKIrish/wkirish3_2017.pdf Editing and imputation methods: Y Where errors are identified, the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during transcription. Imputation of unsampled domains of interest often done at the international level (e.g. ICES), rather than at the national level. However, in some cases, it is appropriate to merge minor domains with larger ones to reduce the burden on the end-user to deal with data gaps. This can include merging ICES divisions or métiers with similar fishing patterns. At a lower level, imputation may also take place in applying age-length keys where length classes for which no age data exists are imputed following the approach described by Gerritsen et al (2006) Gerritsen, Hans D., David McGrath, and Colm Lordan. "A simple method for comparing age-length keys reveals significant regional differences within a single stock of haddock (<i>Melanogrammus aeglefinus</i>). ICES Journal of Marine Science 63.6 (2006): 1096-1100. Quality document associated to a dataset: No DOI is currently created for the dataset since it is not publicly available. R markdown documents recording the estimation and imputation steps performed for ICES data calls are available on request. The estimation is based on the COST R package. An example of the scripts are available at https://www.dcmap-ireland.ie/sites/default/files/DCF_Files/docs/Discard%20Data%20Extraction_example.rmd Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g., in the case of input error) or excluded from that particular use.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.
<u>Description of the Population</u>

No Deviations
<u>Sampling Design and Protocols</u>
No Deviations
<u>Sampling Implementation</u>
The target number of PSUs was not achieved; this is mainly due to high refusal rates. There is great difficulty in getting samplers to sea on the gillnet fleet due to limited accommodation on the size of vessels targeting this fishery. Space is limited on most of these vessels and the current at sea self-sampling protocol is not suitable for the gillnet fishery. A number of trips on the tangle net fishery were carried out in the south west which is a sub set of the gill net fishery. This included 289 self-sampling and 21 observer at sea trips in 2023. We are actively working with gillnet fishermen to develop a sampling strategy that could potentially facilitate at sea self-sampling. This will need to be augmented/validated with observer at sea data. We are also looking into developing a new project which will hopefully help in this regard. It is feared that the introduction of a bycatch only fishery for Pollack may reduce the possible number of gill net trips available, as many of the smaller vessels would traditionally target Pollack during the summer season. However, Ireland will continue to proactively collaborate with the gillnet fleet to achieve DCF targets.
<u>Data Capture</u>
No Deviations
<u>Data Storage</u>
No Deviations
<u>Sample Storage</u>
No Deviations
<u>Data Processing</u>
No Deviations

Sampling scheme identifier: Demersal On - Shore.

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: Demersal On-Shore
Sampling scheme type: Commercial fishing trip
Observation type: SciObsOnShore
Time period of validity: 2014-2027
Description of the population
Population targeted: All commercial catch fractions from the demersal fisheries landed into Ireland from all ICES areas by Irish registered vessels (or for fisheries where a bi-lateral agreement is in place)

Population sampled: 95% of the demersal landings are covered by the sampling program. The top 21 ports are sampled – other ports are excluded. All vessel classes and all commercial species are included in the sampling program. Stratification: Sampling events are stratified by year/quarter and ports are grouped by geographic region (north-west / west / south-west / south-east / east).
Sampling design and protocols Sampling design description: The PSU is port*day. Targets for number of port-days to be sampled are set for each quarter / port – these are proportional to the landings from the relevant reference period. Expert judgement is used to decide which specific port-days are sampled. Vessels present during a sampling event are selected randomly for sampling - the sampler decides how to perform this selection. If there is not time to sample all species landed from a vessel then species selection is made on a priority basis (i.e. rarer species/stocks are prioritised over more commonly sampled species/stock). There is a length-stratified target number of otolith samples per sampling event. Rare/incidental bycatch of fish species are checked during each sampling event. Is the sampling design compliant with the 4S principle?: Y Regional coordination: N Link to sampling design documentation: https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf Compliance with international recommendations: Y. The sampling program has been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints. Link to sampling protocol documentation: https://www.dcmapp-ireland.ie/documents/methodologies Compliance with international recommendations: Y. The sampling protocols have been designed in accordance with guidance from the ICES WGBIOP working group. It has also been necessary to take into account national logistics and constraints.
Sampling implementation Recording of refusal rate: N – since samplers decide which port-days to sample using expert judgement the refusal rate is not relevant for this sampling program. Refusals would only occur extremely rarely in this sampling programme in any case. Monitoring of sampling progress within the sampling year: Sampling progress against targets is tracked within the database and reports of sampling targets and achievements are available to samples (and are automatically updated). If the targets are not being met, then targeted sampling may occur.

Data capture Means of data capture: Data is initially recorded on paper and then transferred to a database application as soon as possible after sampling. Data is measured using ruler and scales. Data capture documentation: SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and reviewed and updated regularly. A recent copy of the SOP for completing data sheets is available at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx Quality checks documentation: Y See “Data_checks_Demersal_Onshore.xlsx” at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Data_checks_Demersal_Onshore.xlsx for a summary of these checks.
Data storage National database: Stockman International database: Detailed data is submitted to RDB / RDBES. Raised data is also currently submitted to the ICES InterCatch database. Quality checks and data validation documentation: Y Data Quality QC functions are available within Stockman – these cover length-weight, age-length, length-frequency distributions e.g.    Data is checked during extractions for end-users such as ICES / European Commission. The checks used will depend on the use of the data. See “Data_checks_Demersal_Onshore.xlsx” at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Data_checks_Demersal_Onshore.xlsx for a summary of these checks. The data management of this data collection activity is incorporated into the Marine Institute’s IODE accredited Data Management – Quality Management Framework (DM-QMF).
Sample storage

Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for analysis. These age structures are stored at Marine Institute premises for a period of months before preparation and age reading is carried out.

Raw data sheets are kept in the regional laboratories and then transferred to headquarters where they are filed. Raw data sheets, duplicate otoliths and read otoliths are archived offsite in secure storage areas after three years. Sectioned otoliths and illicia are stored on site in specialised file storage units.

Data processing

Evaluation of data accuracy (bias and precision): Y

There is not a stand-alone evaluation of bias and precision of the data collected by this scheme but it has been evaluated in recent ICES benchmarks including:

WKIRISH3

2017 http://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/BSG/2017/WKIrish/wkIrish3_2017.pdf ,

WKAnglerfish 2018 http://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2018/WKAngler/WKAngler_2018.pdf

WKCeltic 2020

https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/Fisheries%20Resources%20Steering%20Group/2020/wkceltic_2020.pdf

WKFlatNSCS 2020

https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/Fisheries%20Resources%20Steering%20Group/2020/WKFlatNSCS_2020.pdf

WKDEM 2020 https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/Fisheries%20Resources%20Steering%20Group/2020/wkdem_2020.pdf

Editing and imputation methods: Y

Where errors are identified, the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during transcription.

Imputation of unsampled domains of interest often done at the international level (e.g. ICES), rather than at the national level. However, in some cases, it is appropriate to merge minor domains with larger ones to reduce the burden on the end-user to deal with data gaps. This can include merging ICES divisions or metiers with similar fishing patterns. At a lower level, imputation may also take place in applying age-length keys where length classes for which no age data exists are imputed following the approach described by Gerritsen et al (2006)

Gerritsen, Hans D., David McGrath, and Colm Lordan. "A simple method for comparing age-length keys reveals significant regional differences within a single stock of haddock (*Melanogrammus aeglefinus*).*" ICES Journal of Marine Science* 63.6 (2006): 1096-1100.

Quality document associated to a dataset:

No DOI is currently created for the dataset since it is not publicly available.

R markdown documents recording the estimation and imputation steps performed for ICES data calls are available on request. The estimation is based on the COST R package. An example of the scripts are available at https://www.dcmapireland.ie/sites/default/files/DCF_Files/docs/Cost%20Data%20Extraction_example.Rmd

Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g. in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No Deviations

Sampling Design and Protocols – No Deviations

No Deviations

Sampling Implementation

No Deviations

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: Pelagic On – Shore Others

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: Pelagic on-shore Others
Sampling scheme type: Commercial fishing trip
Observation type: SciObsOnShore
Time period of validity: 2021-2027
The sampling scheme “Pelagic on-shore” is aimed at collecting lengths, weights, age structure sex and is grouped for populations of Mackerel, Horse Mackerel and Blue Whiting landed in Ireland from all ICES areas
Description of the population
Population targeted: Commercial catch of Mackerel, Horse Mackerel and Blue Whiting landed into Ireland from all ICES area by Irish registered fishing vessels
Population sampled: All vessels landing catch for processing at facilities in Killybegs are included where over 95% of the catch of mackerel, horse mackerel and blue whiting landed in Ireland is processed (samples are secured upon arrival at the processing facility).
Stratification: Sampling events are stratified by year/quarter/week, species, ICES division

and fleet segment (RSW or Polyvalent).
Sampling design and protocols
Sampling design description: The PSU is fishing trip*species. Targets are set to maintain species level sampling levels achieved during the reference period. If the number of landings taking place during any week exceeds a threshold, a random selection if trip*species is made, after stratification by species, ICES division and fleet segment. Is the sampling design compliant with the 4S principle?: Y Regional Coordination: N Link to sampling design documentation: https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf Compliance with international recommendations: Y. The sampling program has been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints. Link to sampling protocol documentation: https://www.dcmapp-ireland.ie/documents/methodologies Compliance with international recommendations: Y. The sampling protocols have been designed in accordance with guidance from the ICES WGBIOP working group. It has also been necessary to take into account national logistics and constraints.
Sampling implementation
Recording of refusal rate: N – samplers have access to all potential sampling locations (processing facilities, reference fleet vessels or licenced vessels). Monitoring of sampling progress within the sampling year: Sampling progress against targets is tracked within the database and reports of sampling targets and achievements are available to samples (and are automatically updated). If the targets are not being met, then targeted sampling may occur.
Data capture
Means of data capture: After the sample is secured from the processing factory (or removed from freezer storage and defrosted) it is transported to the local Marine Institute laboratory. Length, weight, sex and maturity are recorded on paper for each specimen. Based on the (species specific) sampling protocol, otolith removal is also carried out. When the data has been recorded for all specimens in the sample, the data is manually input to the Stockman database. Otoliths are prepared for age reading immediately or are transported to another Marine Institute laboratory for preparation before forwarding to the appropriate laboratory for age reading. Ages are initially captured on paper before being entered into Stockman. Data capture documentation: SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and reviewed and updated regularly. A

recent copy of the SOP for completing data sheets is available at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx

Quality checks documentation: Y

See https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf and https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Pelagic_3_PortSampling.pdf for a summary of these checks.

Data capture

Means of data capture:

After the sample is secured from the processing factory (or removed from freezer storage and defrosted) it is transported to the local Marine Institute laboratory.

Length, weight, sex and maturity are recorded on paper for each specimen. Based on the (species specific) sampling protocol, otolith removal is also carried out. When the data has been recorded for all specimens in the sample, the data is manually input to the Stockman database.

Otoliths are prepared for age reading immediately or are transported to another Marine Institute laboratory for preparation before forwarding to the appropriate laboratory for age reading. Ages are initially captured on paper before being entered into Stockman.

Data capture documentation:

SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and reviewed and updated regularly. A recent copy of the SOP for completing data sheets is available at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx

Quality checks documentation: Y

See https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf and https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Pelagic_3_PortSampling.pdf for a summary of these checks.

Data storage


National database: Stockman

International database: Raised data is submitted to the ICES InterCatch database and to ICES accessions email in a proprietary data exchange format specified by end users.

Quality checks and data validation documentation: Y

Length, weight, maturity and age data are checked by comparing data both within an individual sample and between all samples with predefined spatial and temporal domains

Data Quality QC functions are available within Stockman – these cover distributions of lengths, weights, the length-weight regression, age-length, length-frequency distributions e.g.



Data is further checked during final extractions and raising of estimates in response to the needs of end-users such as ICES / European Commission. The specific checks used will depend on the purpose for which the data is requested *e.g.* stock assessment.

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for analysis. These age structures are stored at a Marine Institute premises for a period before further sample preparation and age reading is carried out.

Raw data sheets are kept in the regional laboratories and then transferred to headquarters where they are filed. Raw data sheets, duplicate otoliths and otoliths that have been read are archived offsite in secure storage areas after three years. Sectioned otoliths and illicia are stored on site in specialised file storage units.

Data processing

Evaluation of data accuracy (bias and precision): Y

Stand-alone evaluation of bias and precision of the data collected by this scheme is not conducted. Evaluation takes place at a stock level in assessment benchmark meetings and age reading workshops including:

WKARMAC2 – Report of the Workshop on Age Estimation of Atlantic Mackerel (<https://doi.org/10.17895/ices.pub.8171>)

WKARHOM3 – Workshop on Age Reading of Horse Mackerel, Mediterranean Horse Mackerel and Blue Jack Mackerel (<https://doi.org/10.17895/ices.pub.8170>)

WKARBLUE3 - Report of the Workshop on Age Estimation of Blue Whiting (in prep)

IBPNEAMAC – Interbenchmark Workshop on the Assessment of Northeast Atlantic Mackerel (<https://doi.org/10.17895/ices.pub.4985>)

WKWIDE - Report of the Benchmark Workshop on Widely Distributed Stocks (<https://doi.org/10.17895/ices.pub.5585>)

IBPher6a7bc 2019-Interbenchmark Protocol for Herring in 6.a, 7.b-c (http://doi.org/10.17895/ices.pub.5261) WKWEST 2015 Report of the Benchmark Workshop on West of Scotland Herring and Celtic Sea Herring https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2015/WKWEST/01%20WKWEST%20Report%20%20Final.pdf Editing and imputation methods: Y Where errors are identified, the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during transcription. Imputation of unsampled domains of interest is carried out by the stock coordinator upon receipt of the annual data submission. Quality document associated to a dataset: No DOI is currently created for the dataset since it is not publicly available. Documentation detailing the estimation and imputation steps performed for ICES data calls are available on request. Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g. in the case of input error) or excluded from that particular use.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan. <u>Description of the Population</u> No deviations <u>Sampling Design and Protocols</u> No deviations <u>Sampling Implementation</u> 96 PSUs were achieved, with a target of 107. The undershoot of the target is primarily due to the closure of the directed Western Horse Mackerel fishery, following scientific advice for zero catch in 2023. <u>Data Capture</u> No deviations <u>Data Storage</u> No deviations <u>Sample Storage</u> No deviations <u>Data Processing</u> No deviations

Sampling scheme identifier: Pelagic On – Shore Tuna

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: Pelagic on-shore Tuna
Sampling scheme type: Commercial fishing trip
Observation type: SciObsOnShore
Time period of validity: 2021-2027
Short description (max 100 words): Sampling scheme aiming at collecting lengths and weights from commercial landings on-shore for Tuna. The scheme covers all landings for Tuna in Ireland
Description of the population
Population targeted: Commercial catch of Tuna by Irish pelagic vessels from all ICES areas. Population sampled: All vessels licenced for the fishery are included in the sampling frame. Stratification: Sampling events are stratified by week and ICES division.
Sampling design and protocols
Sampling design description: The PSU is fishing trip*species. Irish tuna landings are highly variable year-to-year. The target number of PSUs therefore needs to be flexible. Targets are set to maintain species level sampling levels achieved during the reference period. Is the sampling design compliant with the 4S principle?: Y Regional Coordination: N Link to sampling design documentation: https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf Compliance with international recommendations: Y. The sampling program has been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints. Link to sampling protocol documentation: https://www.dcmapp-ireland.ie/documents/methodologies Compliance with international recommendations: Y. The sampling protocols have been designed in accordance with guidance from the ICES WGBIOP working group. It has also been necessary to take into account national logistics and constraints.
Sampling implementation
Recording of refusal rate: N – samplers have access to all potential sampling locations (processing facilities, reference fleet vessels or licenced vessels). Monitoring of sampling progress within the sampling year: Sampling progress

against targets is tracked within the database and reports of sampling targets and achievements are available to samples (and are automatically updated). If the targets are not being met, then targeted sampling may occur.

Data capture

Means of data capture:

After the sample is secured from the processing factory (or removed from freezer storage and defrosted) it is transported to the local Marine Institute laboratory.

Length, weight, sex and maturity are recorded on paper for each specimen. Based on the (species specific) sampling protocol, otolith removal is also carried out. When the data has been recorded for all specimens in the sample, the data is manually input to the Stockman database.

Otoliths are prepared for age reading immediately or are transported to another Marine Institute laboratory for preparation before forwarding to the appropriate laboratory for age reading. Ages are initially captured on paper before being entered into Stockman.

Data capture documentation:

SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and reviewed and updated regularly. A recent copy of the SOP for completing data sheets is available at

[https://www.dcmapp-](https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx)

[ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx](https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx)

Quality checks documentation: Y

See [https://www.dcmapp-](https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf)

[ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf](https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf)

and [https://www.dcmapp-](https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Pelagic_3_PortSampling.pdf)

[ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Pelagic_3_PortSampling.pdf](https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Pelagic_3_PortSampling.pdf) for a summary of these checks.

Data storage


National database: Stockman


International database: Raised data is submitted to the ICES InterCatch database and to ICES accessions email in a proprietary data exchange format specified by end users.

Quality checks and data validation documentation: Y

Length, weight, maturity and age data are checked by comparing data both within an individual sample and between all samples with predefined spatial and temporal domains

Data Quality QC functions are available within Stockman – these cover distributions of lengths, weights, the length-weight regression, age-length, length-frequency distributions e.g.





Data is further checked during final extractions and raising of estimates in response to the needs of end-users such as ICES / European Commission. The specific checks used will depend on the purpose for which the data is requested *e.g.* stock assessment.

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for analysis. These age structures are stored at a Marine Institute premises for a period before further sample preparation and age reading is carried out.

Raw data sheets are kept in the regional laboratories and then transferred to headquarters where they are filed. Raw data sheets, duplicate otoliths and otoliths that have been read are archived offsite in secure storage areas after three years. Sectioned otoliths and illicia are stored on site in specialised file storage units.

Data processing

Evaluation of data accuracy (bias and precision): Y

Stand-alone evaluation of bias and precision of the data collected by this scheme is not conducted. Evaluation takes place at a stock level in assessment benchmark meetings and age reading workshops including:

WKARMAC2 – Report of the Workshop on Age Estimation of Atlantic Mackerel (<https://doi.org/10.17895/ices.pub.8171>)

WKARHOM3 – Workshop on Age Reading of Horse Mackerel, Mediterranean Horse Mackerel and Blue Jack Mackerel (<https://doi.org/10.17895/ices.pub.8170>)

WKARBLUE3 - Report of the Workshop on Age Estimation of Blue Whiting (in prep)

IBPNEAMAC – Interbenchmark Workshop on the Assessment of Northeast Atlantic Mackerel (<https://doi.org/10.17895/ices.pub.4985>)

WKWIDE - Report of the Benchmark Workshop on Widely Distributed Stocks (<https://doi.org/10.17895/ices.pub.5585>)

IBPher6a7bc 2019-Interbenchmark Protocol for Herring in 6.a, 7.b-c (<http://doi.org/10.17895/ices.pub.5261>)

WKWEST 2015 Report of the Benchmark Workshop on West of Scotland Herring and Celtic Sea Herring

<https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2015/WKWEST/01%20WKWEST%20Report%20%20Final.pdf>

Editing and imputation methods: Y

Where errors are identified, the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during

transcription.

Imputation of unsampled domains of interest is carried out by the stock coordinator upon receipt of the annual data submission.

Quality document associated to a dataset:

No DOI is currently created for the dataset since it is not publicly available.

Documentation detailing the estimation and imputation steps performed for ICES data calls are available on request.

Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g. in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No deviations

Sampling Design and Protocols

No deviations

Sampling Implementation

2 samples were achieved, fewer than the target of 4. This is as a result of the low proportion of landings of tuna by Irish vessels into Irish ports, with the majority landed into French and Spanish ports. To ensure adequate sampling effort of the Irish fleet, bilateral agreements are in place with France and Spain to ensure landings by Irish vessels into foreign ports are sampled. Additional samples on landings are available from the at-sea sampling trips.

Data Capture

No deviations

Data Storage

No deviations

Sample Storage

No deviations

Data Processing

No deviations

Sampling scheme identifier: Pelagic On – Shore Sprat

MS : IRL

Region: North-East Atlantic

Sampling scheme identifier: Pelagic on-shore Sprat

Sampling scheme type: Commercial fishing trip

Observation type: SciObsOnShore
Time period of validity: 2021-2027
Sampling scheme aiming at collecting lengths and weights from commercial landings on-shore for Sprat. The scheme covers all landings for Sprat in Ireland
Description of the population
Population targeted: Commercial catch of Sprat by Irish pelagic vessels from all ICES areas. Population sampled: All vessels licenced for the fishery are included in the sampling frame. Stratification: Sampling events are stratified by week and ICES division.
Sampling design and protocols
Sampling design description: The PSU is fishing trip*species. Irish sprat landings are highly variable year-to-year. The target number of PSUs therefore needs to be flexible. Targets are set to maintain species level sampling levels achieved during the reference period. Is the sampling design compliant with the 4S principle?: Y Regional Coordination: N Link to sampling design documentation: https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf Compliance with international recommendations: Y. The sampling program has been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints. Link to sampling protocol documentation: https://www.dcmapp-ireland.ie/documents/methodologies Compliance with international recommendations: Y. The sampling protocols have been designed in accordance with guidance from the ICES WGBIOP working group. It has also been necessary to take into account national logistics and constraints.
Sampling implementation
Recording of refusal rate: N – samplers have access to all potential sampling locations (processing facilities, reference fleet vessels or licenced vessels). Monitoring of sampling progress within the sampling year: Sampling progress against targets is tracked within the database and reports of sampling targets and achievements are available to samples (and are automatically updated). If the targets are not being met, then targeted sampling may occur.
Data capture
Means of data capture: After the sample is secured from the processing factory (or removed from freezer storage and defrosted) it is transported to the local Marine Institute laboratory. Length, weight, sex and maturity are recorded on paper for each specimen. Based on the

(species specific) sampling protocol, otolith removal is also carried out. When the data has been recorded for all specimens in the sample, the data is manually input to the Stockman database.

Otoliths are prepared for age reading immediately or are transported to another Marine Institute laboratory for preparation before forwarding to the appropriate laboratory for age reading. Ages are initially captured on paper before being entered into Stockman.

Data capture documentation:

SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and reviewed and updated regularly. A recent copy of the SOP for completing data sheets is available at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx

Quality checks documentation: Y

See https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf and https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Pelagic_3_PortSampling.pdf for a summary of these checks.

Data capture

Means of data capture:

After the sample is secured from the processing factory (or removed from freezer storage and defrosted) it is transported to the local Marine Institute laboratory.

Length, weight, sex and maturity are recorded on paper for each specimen. Based on the (species specific) sampling protocol, otolith removal is also carried out. When the data has been recorded for all specimens in the sample, the data is manually input to the Stockman database.

Otoliths are prepared for age reading immediately or are transported to another Marine Institute laboratory for preparation before forwarding to the appropriate laboratory for age reading. Ages are initially captured on paper before being entered into Stockman.

Data capture documentation:

SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and reviewed and updated regularly. A recent copy of the SOP for completing data sheets is available at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx

Quality checks documentation: Y

See https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf and https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Pelagic_3_PortSampling.pdf for a summary of these checks.

Data storage


National database: Stockman

International database: Raised data is submitted to the ICES InterCatch database and to ICES accessions email in a proprietary data exchange format specified by end users.

Quality checks and data validation documentation: Y

Length, weight, maturity and age data are checked by comparing data both within an individual sample and between all samples with predefined spatial and temporal domains

Data Quality QC functions are available within Stockman – these cover distributions of lengths, weights, the length-weight regression, age-length, length-frequency distributions e.g.



Data is further checked during final extractions and raising of estimates in response to the needs of end-users such as ICES / European Commission. The specific checks used will depend on the purpose for which the data is requested *e.g.* stock assessment.

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for analysis. These age structures are stored at a Marine Institute premises for a period before further sample preparation and age reading is carried out.

Raw data sheets are kept in the regional laboratories and then transferred to headquarters where they are filed. Raw data sheets, duplicate otoliths and otoliths that have been read are archived offsite in secure storage areas after three years. Sectioned otoliths and illicia are stored on site in specialised file storage units.

Data processing

Evaluation of data accuracy (bias and precision): Y

Stand-alone evaluation of bias and precision of the data collected by this scheme is not conducted. Evaluation takes place at a stock level in assessment benchmark meetings and age reading workshops including:

WKARMAC2 – Report of the Workshop on Age Estimation of Atlantic Mackerel
(<https://doi.org/10.17895/ices.pub.8171>)

WKARHOM3 – Workshop on Age Reading of Horse Mackerel, Mediterranean Horse

Mackerel and Blue Jack Mackerel (https://doi.org/10.17895/ices.pub.8170) WKARBLUE3 - Report of the Workshop on Age Estimation of Blue Whiting (in prep) IBPNEAMAC – Interbenchmark Workshop on the Assessment of Northeast Atlantic Mackerel (https://doi.org/10.17895/ices.pub.4985) WKWIDE - Report of the Benchmark Workshop on Widely Distributed Stocks (https://doi.org/10.17895/ices.pub.5585) IBPher6a7bc 2019-Interbenchmark Protocol for Herring in 6.a, 7.b-c (http://doi.org/10.17895/ices.pub.5261) WKWEST 2015 Report of the Benchmark Workshop on West of Scotland Herring and Celtic Sea Herring https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2015/WKWEST/01%20WKWEST%20Report%20%20Final.pdf Editing and imputation methods: Y Where errors are identified, the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during transcription. Imputation of unsampled domains of interest is carried out by the stock coordinator upon receipt of the annual data submission. Quality document associated to a dataset: No DOI is currently created for the dataset since it is not publicly available. Documentation detailing the estimation and imputation steps performed for ICES data calls are available on request. Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g. in the case of input error) or excluded from that particular use.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan. <u>Description of the Population</u> No deviations <u>Sampling Design and Protocols</u> No deviations <u>Sampling Implementation</u> No deviations <u>Data Capture</u> No deviations <u>Data Storage</u> No deviations <u>Sample Storage</u> No deviations

Data Processing
No deviations

Sampling scheme identifier: Pelagic On – Shore Boarfish

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: Pelagic on-shore Boarfish
Sampling scheme type: Commercial fishing trip
Observation type: SciObsOnShore
Time period of validity: 2021-2027
Sampling scheme aiming at collecting lengths, weights, sex maturity and age structures from commercial landings on-shore for Boarfish. The scheme covers all landings for Boarfish in Ireland
Description of the population
Population targeted: Commercial catch of Boarfish by Irish pelagic vessels from all ICES areas. Population sampled: Catches of boarfish by reference fleet Stratification: Sampling events are stratified by week and ICES division
Sampling design and protocols
Sampling design description: The PSU is fishing trip*ICES division*week for vessels in the reference fleet. Targets (2 samples per week per ICES division) are set to maintain sampling levels achieved during the reference period. A sample is taken halfway through the pumping operation, bagged, labelled and frozen and delivered to a Marine Institute laboratory at the next available opportunity. Is the sampling design compliant with the 4S principle?: Y Regional Coordination: N Link to sampling design documentation: https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf Compliance with international recommendations: Y. The sampling program has been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints. Link to sampling protocol documentation: https://www.dcmapp-ireland.ie/documents/methodologies Compliance with international recommendations: Y. The sampling protocols have been designed in accordance with guidance from the ICES WGBIOP working group. It has also been necessary to take into account national logistics

and constraints.
Sampling implementation
Recording of refusal rate: N – samplers have access to all potential sampling locations (processing facilities, reference fleet vessels or licenced vessels). Monitoring of sampling progress within the sampling year: Sampling progress against targets is tracked within the database and reports of sampling targets and achievements are available to samples (and are automatically updated). If the targets are not being met, then targeted sampling may occur.
Data capture
Means of data capture: After the sample is secured from the processing factory (or removed from freezer storage and defrosted) it is transported to the local Marine Institute laboratory. Length, weight, sex and maturity are recorded on paper for each specimen. Based on the (species specific) sampling protocol, otolith removal is also carried out. When the data has been recorded for all specimens in the sample, the data is manually input to the Stockman database. Otoliths are prepared for age reading immediately or are transported to another Marine Institute laboratory for preparation before forwarding to the appropriate laboratory for age reading. Ages are initially captured on paper before being entered into Stockman. Data capture documentation: SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and reviewed and updated regularly. A recent copy of the SOP for completing data sheets is available at https://www.dcmapireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx Quality checks documentation: Y See https://www.dcmapireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf and https://www.dcmapireland.ie/sites/default/files/DCF_Files/docs/FEAS_Pelagic_3_PortSampling.pdf for a summary of these checks.
Data capture
Means of data capture: After the sample is secured from the processing factory (or removed from freezer storage and defrosted) it is transported to the local Marine Institute laboratory. Length, weight, sex and maturity are recorded on paper for each specimen. Based on the (species specific) sampling protocol, otolith removal is also carried out. When the data has been recorded for all specimens in the sample, the data is manually input to the Stockman database. Otoliths are prepared for age reading immediately or are transported to another Marine Institute laboratory for preparation before forwarding to the appropriate laboratory for age reading. Ages are initially captured on paper before being entered into Stockman. Data capture documentation: SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and reviewed and updated regularly.

A recent copy of the SOP for completing data sheets is available at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx

Quality checks documentation: Y

See https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf and https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Pelagic_3_PortSampling.pdf for a summary of these checks.


Data storage

National database: Stockman

International database: Raised data is submitted to the ICES InterCatch database and to ICES accessions email in a proprietary data exchange format specified by end users.

Quality checks and data validation documentation: Y

Length, weight, maturity and age data are checked by comparing data both within an individual sample and between all samples with predefined spatial and temporal domains. Data Quality QC functions are available within Stockman – these cover distributions of lengths, weights, the length-weight regression, age-length, length-frequency distributions e.g.



Data is further checked during final extractions and raising of estimates in response to the needs of end-users such as ICES / European Commission. The specific checks used will depend on the purpose for which the data is requested e.g. stock assessment.

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for analysis. These age structures are stored at a Marine Institute premises for a period before further sample preparation and age reading is carried out.

Raw data sheets are kept in the regional laboratories and then transferred to headquarters where they are filed. Raw data sheets, duplicate otoliths and otoliths that have been read are archived offsite in secure storage areas after three years. Sectioned otoliths and illicia are stored on site in specialised file storage units.

Data processing

Evaluation of data accuracy (bias and precision): Y

Stand-alone evaluation of bias and precision of the data collected by this scheme is not conducted. Evaluation takes place at a stock level in assessment benchmark meetings and age reading workshops including:

WKARMAC2 – Report of the Workshop on Age Estimation of Atlantic Mackerel (<https://doi.org/10.17895/ices.pub.8171>)

WKARHOM3 – Workshop on Age Reading of Horse Mackerel, Mediterranean Horse Mackerel and Blue Jack Mackerel (<https://doi.org/10.17895/ices.pub.8170>)

WKARBLUE3 - Report of the Workshop on Age Estimation of Blue Whiting (in prep)

IBPNEAMAC – Interbenchmark Workshop on the Assessment of Northeast Atlantic Mackerel (<https://doi.org/10.17895/ices.pub.4985>)

WKWIDE - Report of the Benchmark Workshop on Widely Distributed Stocks (<https://doi.org/10.17895/ices.pub.5585>)

IBPher6a7bc 2019-Interbenchmark Protocol for Herring in 6.a, 7.b-c (<http://doi.org/10.17895/ices.pub.5261>)

WKWEST 2015 Report of the Benchmark Workshop on West of Scotland Herring and Celtic Sea Herring

<https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2015/WKWEST/01%20WKWEST%20Report%20%20Final.pdf>

Editing and imputation methods: Y

Where errors are identified, the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during transcription.

Imputation of unsampled domains of interest is carried out by the stock coordinator upon receipt of the annual data submission.

Quality document associated to a dataset:

No DOI is currently created for the dataset since it is not publicly available.

Documentation detailing the estimation and imputation steps performed for ICES data calls are available on request.

Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g. in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No deviations

<u>Sampling Design and Protocols</u>
No deviations
<u>Sampling Implementation</u>
A fewer number of samples (2) than planned (15) were achieved for the 2 vessel reference fleet. However, a significant number of additional samples (19) were secured from other vessels (similar to those in the reference fleet) targeting boarfish such that the total number achieved exceeds the target. A greater proportion of landings have been made into Irish ports in recent years, allowing port based staff to secure these additional sampling opportunities.
<u>Data Capture</u>
No deviations
<u>Data Storage</u>
No deviations
<u>Sample Storage</u>
No deviations
<u>Data Processing</u>
No deviations

Sampling scheme identifier: Pelagic On – Shore Norwegian Spring Spawning Herring

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: Pelagic on-shore Norwegian Spring Spawning Herring
Sampling scheme type: Commercial fishing trip
Observation type: SciObsOnShore
Time period of validity: 2021-2027
Sampling scheme aiming at collecting lengths, weights, sex maturity and age structures from commercial landings on-shore for Norwegian Spring Spawning Herring The scheme covers all landings for Norwegian Spring Spawning Herring in Ireland
Description of the population
Population targeted: Commercial catch of Norwegian Spring Spawning Herring by Irish pelagic vessels from all ICES areas. Population sampled: All vessels licenced for the fishery are included in the sampling frame. Stratification: Vessels are selected at random from the sampling frame without stratification.
Sampling design and protocols
Sampling design description: The PSU is fishing trip. Vessels are selected at random from the list of licenced vessels participating in the fishery. Selected vessels secure a sample from a random haul which is frozen onboard and delivered to a Marine Institute laboratory at the

next available opportunity

Is the sampling design compliant with the 4S principle?: Y

Regional Coordination: N

Link to sampling design documentation:

https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf

Compliance with international recommendations: Y. The sampling program has been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints.

Link to sampling protocol documentation:

<https://www.dcmapp-ireland.ie/documents/methodologies>

Compliance with international recommendations: Y.

The sampling protocols have been designed in accordance with guidance from the ICES WGBIOP working group. It has also been necessary to take into account national logistics and constraints.

Sampling implementation

Recording of refusal rate: N – samplers have access to all potential sampling locations (processing facilities, reference fleet vessels or licenced vessels).

Monitoring of sampling progress within the sampling year: Sampling progress against targets is tracked within the database and reports of sampling targets and achievements are available to samples (and are automatically updated). If the targets are not being met, then targeted sampling may occur.

Data capture

Means of data capture:

After the sample is secured from the processing factory (or removed from freezer storage and defrosted) it is transported to the local Marine Institute laboratory.

Length, weight, sex and maturity are recorded on paper for each specimen. Based on the (species specific) sampling protocol, otolith removal is also carried out. When the data has been recorded for all specimens in the sample, the data is manually input to the Stockman database.

Otoliths are prepared for age reading immediately or are transported to another Marine Institute laboratory for preparation before forwarding to the appropriate laboratory for age reading. Ages are initially captured on paper before being entered into Stockman.

Data capture documentation:

SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and reviewed and updated regularly. A recent copy of the SOP for completing data sheets is available at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx

Quality checks documentation: Y

See https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf

and https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Pelagic_3_PortSampling.pdf for a summary of these checks.

Data capture

Means of data capture:

After the sample is secured from the processing factory (or removed from freezer storage and defrosted) it is transported to the local Marine Institute laboratory.

Length, weight, sex and maturity are recorded on paper for each specimen. Based on the (species specific) sampling protocol, otolith removal is also carried out. When the data has been recorded for all specimens in the sample, the data is manually input to the Stockman database.

Otoliths are prepared for age reading immediately or are transported to another Marine Institute laboratory for preparation before forwarding to the appropriate laboratory for age reading. Ages are initially captured on paper before being entered into Stockman.

Data capture documentation:

SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and reviewed and updated regularly. A recent copy of the SOP for completing data sheets is available at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx

Quality checks documentation: Y

See https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf and https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Pelagic_3_PortSampling.pdf for a summary of these checks.

Data storage




National database: Stockman


International database: Raised data is submitted to the ICES InterCatch database and to ICES accessions email in a proprietary data exchange format specified by end users.

Quality checks and data validation documentation: Y

Length, weight, maturity and age data are checked by comparing data both within an individual sample and between all samples with predefined spatial and temporal domains

Data Quality QC functions are available within Stockman – these cover distributions of lengths, weights, the length-weight regression, age-length, length-frequency distributions e.g.





Data is further checked during final extractions and raising of estimates in response to the needs of end-users such as ICES / European Commission. The specific checks used will depend on the purpose for which the data is requested *e.g.* stock assessment.

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for analysis. These age structures are stored at a Marine Institute premises for a period before further sample preparation and age reading is carried out.

Raw data sheets are kept in the regional laboratories and then transferred to headquarters where they are filed. Raw data sheets, duplicate otoliths and otoliths that have been read are archived offsite in secure storage areas after three years. Sectioned otoliths and illicia are stored on site in specialised file storage units.

Data processing

Evaluation of data accuracy (bias and precision): Y

Stand-alone evaluation of bias and precision of the data collected by this scheme is not conducted. Evaluation takes place at a stock level in assessment benchmark meetings and age reading workshops including:

WKARMAC2 – Report of the Workshop on Age Estimation of Atlantic Mackerel (<https://doi.org/10.17895/ices.pub.8171>)

WKARHOM3 – Workshop on Age Reading of Horse Mackerel, Mediterranean Horse Mackerel and Blue Jack Mackerel (<https://doi.org/10.17895/ices.pub.8170>)

WKARBLUE3 - Report of the Workshop on Age Estimation of Blue Whiting (in prep)

IBPNEAMAC – Interbenchmark Workshop on the Assessment of Northeast Atlantic Mackerel (<https://doi.org/10.17895/ices.pub.4985>)

WKWIDE - Report of the Benchmark Workshop on Widely Distributed Stocks (<https://doi.org/10.17895/ices.pub.5585>)

IBPher6a7bc 2019-Interbenchmark Protocol for Herring in 6.a, 7.b-c (<http://doi.org/10.17895/ices.pub.5261>)

WKWEST 2015 Report of the Benchmark Workshop on West of Scotland Herring and Celtic Sea Herring

<https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2015/WKWEST/01%20WKWEST%20Report%20%20Final.pdf>

Editing and imputation methods: Y

Where errors are identified, the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during transcription.

Imputation of unsampled domains of interest is carried out by the stock coordinator upon receipt of the annual data submission.

Quality document associated to a dataset:

No DOI is currently created for the dataset since it is not publicly available.

Documentation detailing the estimation and imputation steps performed for ICES data calls are available on request.

Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g. in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No deviations

Sampling Design and Protocols

No deviations

Sampling Implementation

50% of the target number of samples were achieved. This is due to a delay in the allocation of quota to vessels and a reduction in the number of fishing trips in 2023.

Data Capture

No deviations

Data Storage

No deviations

Sample Storage

No deviations

Data Processing

No deviations

Sampling scheme identifier: Pelagic On – Shore Herring

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: Pelagic on-shore Herring
Sampling scheme type: Commercial fishing trip
Observation type: SciObsOnShore
Time period of validity: 2021-2027
Sampling scheme aiming at collecting lengths, weights, sex maturity and age structures from commercial landings on-shore for Herring The scheme covers all landings for Herring in

Ireland
Description of the population
Population targeted: Commercial catch of Celtic Sea herring, Irish Sea herring and 6a (north West) herring by Irish pelagic vessels from ICES areas 6 and 7. Population sampled: All vessels licenced for the fishery are included in the sampling frame, including the main and sentinel fleet in the Celtic Sea and all authorised vessels for the monitoring TAC in 6a.7bc. Stratification: Vessels are selected at random from the sampling frame without stratification.
Sampling design and protocols
Sampling design description: Celtic Sea - The PSU is fishing haul * area or day. ICES has advised a total target PSU number of 17 across the main and sentinel fleets. All vessels from the authorised list of licenced vessels participating in the fishery are included. Special fishery arrangements and close liaison between Marine Institute scientists, fish producer organisations and vessel skippers is in place to secure the necessary samples. 6a.7bc (North West) - The PSU is fishing trip. All vessels from the authorised list of licenced vessels participating in the fishery are included. Special fishery arrangements and close liaison between Marine Institute scientists, fish producer organisations and vessel skippers is in place to secure the necessary samples. Irish Sea Herring - The PSU is fishing trip. Targets are set to maintain sampling levels achieved during the reference period. Is the sampling design compliant with the 4S principle?: Y* *Celtic Sea and 6a sampling designs are dictated by the monitoring TAC needs. Once these stocks have been rebuilt, further planning will be required to be compliant with 4S sampling design. Regional Coordination: N All landings of herring to be sampled in the Celtic Sea and 6a (North West) as per the monitoring TAC arrangements. PSU for Celtic Sea is haul*area or day. PSU for 6a (North West) is trip. Link to sampling design documentation: https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf Compliance with international recommendations: Y. The sampling program has been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints. Link to sampling protocol documentation: https://www.dcmapp-ireland.ie/documents/methodologies Compliance with international recommendations: Y. The sampling protocols have been designed in accordance with guidance from the ICES WGBIOP working group. It has also been necessary to take into account national logistics and constraints.
Sampling implementation

Recording of refusal rate: N – samplers have access to all potential sampling locations (processing facilities, reference fleet vessels or licenced vessels).

Monitoring of sampling progress within the sampling year: Sampling progress against targets is tracked within the database and reports of sampling targets and achievements are available to samples (and are automatically updated). If the targets are not being met, then targeted sampling may occur.

Data capture

Means of data capture:

After the sample is secured from the processing factory (or removed from freezer storage and defrosted) it is transported to the local Marine Institute laboratory.

Length, weight, sex and maturity are recorded on paper for each specimen. Based on the (species specific) sampling protocol, otolith removal is also carried out. When the data has been recorded for all specimens in the sample, the data is manually input to the Stockman database.

Otoliths are prepared for age reading immediately or are transported to another Marine Institute laboratory for preparation before forwarding to the appropriate laboratory for age reading. Ages are initially captured on paper before being entered into Stockman.

Data capture documentation:

SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and reviewed and updated regularly. A recent copy of the SOP for completing data sheets is available at https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling.docx

Quality checks documentation: Y

See https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/PortSamplingSummaryReport.pdf and https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Pelagic_3_PortSampling.pdf for a summary of these checks.

Data storage


National database: Stockman

International database: Raised data is submitted to the ICES InterCatch database and to ICES accessions email in a proprietary data exchange format specified by end users.

Quality checks and data validation documentation: Y

Length, weight, maturity and age data are checked by comparing data both within an individual sample and between all samples with predefined spatial and temporal domains

Data Quality QC functions are available within Stockman – these cover distributions of lengths, weights, the length-weight regression, age-length, length-frequency distributions e.g.



Data is further checked during final extractions and raising of estimates in response to the needs of end-users such as ICES / European Commission. The specific checks used will depend on the purpose for which the data is requested *e.g.* stock assessment.

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for analysis. These age structures are stored at a Marine Institute premises for a period before further sample preparation and age reading is carried out.

Raw data sheets are kept in the regional laboratories and then transferred to headquarters where they are filed. Raw data sheets, duplicate otoliths and otoliths that have been read are archived offsite in secure storage areas after three years. Sectioned otoliths and illicia are stored on site in specialised file storage units.

Data processing

Evaluation of data accuracy (bias and precision): Y

Stand-alone evaluation of bias and precision of the data collected by this scheme is not conducted. Evaluation takes place at a stock level in assessment benchmark meetings and age reading workshops including:

WKARMAC2 – Report of the Workshop on Age Estimation of Atlantic Mackerel (<https://doi.org/10.17895/ices.pub.8171>)

WKARHOM3 – Workshop on Age Reading of Horse Mackerel, Mediterranean Horse Mackerel and Blue Jack Mackerel (<https://doi.org/10.17895/ices.pub.8170>)

WKARBLUE3 - Report of the Workshop on Age Estimation of Blue Whiting (in prep)

IBPNEAMAC – Interbenchmark Workshop on the Assessment of Northeast Atlantic Mackerel (<https://doi.org/10.17895/ices.pub.4985>)

WKWIDE - Report of the Benchmark Workshop on Widely Distributed Stocks (<https://doi.org/10.17895/ices.pub.5585>)

IBPher6a7bc 2019-Interbenchmark Protocol for Herring in 6.a, 7.b-c (http://doi.org/10.17895/ices.pub.5261) WKWEST 2015 Report of the Benchmark Workshop on West of Scotland Herring and Celtic Sea Herring https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2015/WKWEST/01%20WKWEST%20Report%20%20Final.pdf Editing and imputation methods: Y Where errors are identified, the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during transcription. Imputation of unsampled domains of interest is carried out by the stock coordinator upon receipt of the annual data submission. Quality document associated to a dataset: No DOI is currently created for the dataset since it is not publicly available. Documentation detailing the estimation and imputation steps performed for ICES data calls are available on request. Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g. in the case of input error) or excluded from that particular use. AR comment: Indicate any deviations. Do not change the text already adopted in the work plan. <u>Description of the Population</u> No Deviations <u>Sampling Design and Protocols</u> No Deviations <u>Sampling Implementation</u> No Deviations <u>Data Capture</u> No Deviations <u>Data Storage</u> No Deviations <u>Sample Storage</u> No Deviations <u>Data Processing</u> No Deviations

Sampling scheme identifier: Pelagic at - sea Others

MS : IRL

Region: North-East Atlantic
Sampling scheme identifier: Pelagic at-sea Others
Sampling scheme type: Commercial fishing trip
Observation type: SciObsAtSea
Time period of validity: 2021-2027
Short description (max 100 words): A sampling scheme aiming at collecting length samples from commercial catches at sea targeting mackerel, horse mackerel, blue whiting, boarfish and Norwegian spring spawning herring The scheme covers all relevant pelagic fisheries in 27.2-8
Description of the population
Pelagic at-sea others Population targeted: The target population is the group of vessels that are engaged in the target mackerel, horse mackerel, blue whiting, boarfish and Norwegian spring spawning herring fisheries. Population sampled: All vessels that reported landings in the same semester of the previous year are included in the sampling frame. Stratification: The sampling frames are stratified by semester.
Sampling design and protocols
Sampling design description: The PSU is vessel*time. The sampling frame is a list of vessels active in the target fisheries for mackerel, horse mackerel, blue whiting, boarfish and Norwegian spring spawning herring during the reference period. Each vessel has a sampling probability based on average landings during the relevant semester over the reference period. Vessels are sampled from the list with replacement. Sampling is carried out by a sampler on-board for the duration of the fishing trip. Rare/incidental bycatch of fish species are checked during each sampling event. Any Bird/Mammal/Reptile/PET/Decomposed organism that comes in contact with the gear during fishing operation is also recorded. VME indicator species are noted if present in the random sample box
Is the sampling design compliant with the 4S principle?: Y
Regional coordination: N
Link to sampling design documentation: https://www.dcmapireland.ie/documents/methodologies
Compliance with international recommendations: Y. The sampling program has been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and

constraints. .

Link to sampling protocol documentation:

https://www.demap-ireland.ie/sites/default/files/DCF_Files/docs/At-Sea%20Sampling%20Manual_2020.pdf

Compliance with international recommendations: Y. The sampling protocols have been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints.

Sampling implementation

Recording of refusal rate: Y – Refusals and analysis of refusal rates are in accordance with guidance from SGPIDS, 2013.

Monitoring of sampling progress within the sampling year: : Sampling progress against targets is tracked within the database and reports of sampling targets and achievements are available to samples.

Data capture

Means of data capture: Data is initially recorded on paper and then transferred to a database application as soon as possible after sampling. Fish lengths are measured using 1 metre ruler; fish weights are only recorded for samples that are brought back to the lab.

Data capture documentation: SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and reviewed and updated regularly. A recent copy of the SOP for completing data sheets is available at https://www.demap-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling_0.docx

Quality checks documentation: Data Validation occurs within the entry screens of the database. Data Checking occurs with a Voice Reports software package that reads what has been entered in the database and allows the data enterer to check the records that are being read out against the records written on the datasheets.

Data storage

National database: Pelagic Discards Database (Access)

International database: Data (unraised) submitted to ICES in request to annual data call.

Quality checks and data validation documentation:

Data Validation occurs within the entry screens of the database. Input is constrained by the use of drop-down lists with further checks performed by the DB application code.

Data is further checked during extractions for end-users such as ICES / European

Commission. The checks used will depend on the use of the data. Migration to SQL Server will be undertaken following completion of the design and specification phases. The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF). .
Sample storage Fish samples are processed in Marine Institute laboratories for length, weight and maturity analysis. Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for processing and analysis. These age structures are stored at Marine Institute premises for a period of months before preparation and age reading is carried out. Raw data sheets are kept in the regional laboratories and then transferred to headquarters where they are filed. Raw data sheets, duplicate otoliths and read otoliths are archived offsite in secure storage areas after three years. Sectioned otoliths and illicia are stored on site in specialised file storage units.
Data processing Evaluation of data accuracy (bias and precision): No stand-alone evaluation of bias and precision of the data collected by this scheme. Any data collected is submitted to end users at which point accuracy will be assessed. Editing and imputation methods: Y Where errors are identified, the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during transcription. Imputation of unsampled domains of interest often done at the international level (e.g. ICES), rather than at the national level. Quality document associated to a dataset: No DOI is currently created for the dataset since it is not publicly available. Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g., in the case of input error) or excluded from that particular use.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan. <u>Description of the Population</u> No Deviations <u>Sampling Design and Protocols</u> No Deviations

<u>Sampling Implementation</u> Shortfall in achieved target was due to logistical challenges in 2023 in securing observer trips on the pelagic fleet. Additional observers have been recruited and are now fully trained. One of these observers is located in Irelands largest pelagic port, which should aid in the achievement of at sea sampling targets. We will continue to engage with key stakeholders such as Fish Producers Organisations and Processors on a local level to secure sampling opportunities.
<u>Data Capture</u> No Deviations
<u>Data Storage</u> No Deviations
<u>Sample Storage</u> No Deviations
<u>Data Processing</u> No Deviations

Sampling scheme identifier: Pelagic at - sea Tuna

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: Pelagic at-sea Tuna
Sampling scheme type: Commercial fishing trip
Observation type: SciObsAtSea
Time period of validity: 2021-2027
Short description (max 100 words): A sampling scheme aiming at collecting length samples from commercial catches at sea targeting Tuna in 27. 7-8
Description of the population
Pelagic at-sea Tuna Population targeted: The target population is the group of vessels that are engaged in the target Tuna fisheries. Population sampled: All vessels that are authorised to fish for each tuna stock in the sampling year are included in the sampling frame. The yearly lists of authorisations are compiled by the relevant section of the Department of Agriculture, Food and the Marine (DAFM) in conjunction with management advisory committees. Stratification: The sampling frames are stratified by semester
Sampling design and protocols
Sampling design description:

The PSU is vessel*time. The sampling frame is a list of vessels active in the target fisheries for Tuna during the reference period. Sampling is carried out by a sampler on board the vessels is assigned by the industry for the duration of the fishing trip.

Rare/incidental bycatch of fish species are checked during each sampling event.

Any Bird/Mammal/Reptile/PET/Decomposed organism that comes in contact with the gear during fishing operation is also recorded.

VME indicator species are noted if present in the random sample box

Is the sampling design compliant with the 4S principle?: Y

Regional coordination: N

Link to sampling design documentation: <https://www.dcmapp-ireland.ie/documents/methodologies>

Compliance with international recommendations: Y. The sampling program has been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints. .

Link to sampling protocol documentation:

https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/At-Sea%20Sampling%20Manual_2020.pdf

Compliance with international recommendations: Y. The sampling protocols have been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints.

Sampling implementation

Recording of refusal rate: Y – Refusals and analysis of refusal rates are in accordance with guidance from SGPIDS, 2013.

Monitoring of sampling progress within the sampling year: Sampling progress against targets is tracked within the database and reports of sampling targets and achievements are available to samples.

Data capture

Means of data capture: Data is initially recorded on paper and then transferred to a database application as soon as possible after sampling. Fish lengths are measured using 1 metre ruler; fish weights are only recorded for samples that are brought back to the lab.

Data capture documentation: SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and

reviewed and updated regularly. A recent copy of the SOP for completing data sheets is available at https://www.dcmapireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling_0.docx

Quality checks documentation: Data Validation occurs within the entry screens of the database. Data Checking occurs with a Voice Reports software package that reads what has been entered in the database and allows the data enterer to check the records that are being read out against the records written on the datasheets.

Data storage

National database: Pelagic Discards Database (Access)

International database: Data (unraised) submitted to ICES in request to annual data call.

Quality checks and data validation documentation:

Data Validation occurs within the entry screens of the database. Input is constrained by the use of drop-down lists with further checks performed by the DB application code.

Data is further checked during extractions for end-users such as ICES / European Commission. The checks used will depend on the use of the data.

Migration to SQL Server will be undertaken following completion of the design and specification phases.

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

Fish samples are processed in Marine Institute laboratories for length, weight and maturity analysis. Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for processing and analysis. These age structures are stored at Marine Institute premises for a period of months before preparation and age reading is carried out.

Raw data sheets are kept in the regional laboratories and then transferred to headquarters where they are filed. Raw data sheets, duplicate otoliths and read otoliths are archived offsite in secure storage areas after three years. Sectioned otoliths and illicia are stored on site in specialised file storage units.

Data processing

Evaluation of data accuracy (bias and precision):

No stand-alone evaluation of bias and precision of the data collected by this scheme. Any data collected is submitted to end users at which point accuracy will be assessed.

Editing and imputation methods: Y Where errors are identified, the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during transcription.

Imputation of unsampled domains of interest often done at the international level (e.g. ICES), rather than at the national level.

Quality document associated to a dataset:

No DOI is currently created for the dataset since it is not publicly available.

Validation of the final dataset:

Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g., in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No Deviations

Sampling Design and Protocols

The target PSU were reached in 2023. Sampling was increased by the introduction of an Albacore Fishery-At Sea Self Sampling program where skippers collected data according to the appointed SOP on a daily basis and delivered this data to the MI. The data was quality controlled live on a daily basis via WhatsApp communication with the fleet. The sea self-sampling augments the sampler at sea data collected under this program and is especially relevant in increasing the observations in relation to potential PET bycatch.

Sampling Implementation

No Deviations

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: Pelagic at - sea Herring

MS : IRL

Region: North-East Atlantic
Sampling scheme identifier: Pelagic at-sea Herring
Sampling scheme type: Commercial fishing trip
Observation type: SciObsAtSea
Time period of validity: 2021-2027
Short description (max 100 words): A sampling scheme aiming at collecting length samples from commercial catches at sea targeting Herring in 6a.7bc, Celtic Sea, Irish Sea
Description of the population
Population targeted: The target population is the group of vessels that are engaged in the Irish fisheries targeting the herring stocks of 6a.7bc (NW), Celtic Sea, and Irish Sea. Population sampled: All vessels that are authorised to fish for each herring stock in the sampling year are included in the sampling frame. The yearly lists of authorisations are compiled by the relevant section of the Department of Agriculture, Food and the Marine (DAFM) in conjunction with management advisory committees. Stratification: The sampling frames are stratified by semester.
Sampling design and protocols
Sampling design description: The PSU is vessel*time. The three sampling frames are the authorised list of Irish vessels licenced to target herring in the three stocks (6a.7bc, Celtic Sea, Irish Sea). Vessels are sampled from the list with replacement. Sampling is carried out by a sampler on-board for the duration of the fishing trip. Rare/incidental bycatch of fish species are checked during each sampling event. Any Bird/Mammal/Reptile/PET/Decomposed organism that comes in contact with the gear during fishing operation is also recorded. VME indicator species are noted if present in the random sample box Is the sampling design compliant with the 4S principle?: Y Regional coordination: N Link to sampling design documentation: https://www.dcmaphireland.ie/documents/methodologies Compliance with international recommendations: Y. The sampling program has been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints. Link to sampling protocol documentation: https://www.dcmaphireland.ie/sites/default/files/DCF_Files/docs/At-Sea%20Sampling%20Manual_2020.pdf

Compliance with international recommendations: Y. The sampling protocols have been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints.
Sampling implementation
Recording of refusal rate: Y – Refusals and analysis of refusal rates are in accordance with guidance from SGPIDS, 2013. Monitoring of sampling progress within the sampling year: Sampling progress against targets is tracked within the database and reports of sampling targets and achievements are available to samples.
Data capture
Means of data capture: Data is initially recorded on paper and then transferred to a database application as soon as possible after sampling. Fish lengths are measured using 1 metre ruler; fish weights are only recorded for samples that are brought back to the lab. Data capture documentation: SOPs for sampling (weighing / measuring / completing data sheets / otolith storage etc.) are held in Paradigm3 (a document management system) and reviewed and updated regularly. A recent copy of the SOP for completing data sheets is available at https://www.dcmapi-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling_0.docx Quality checks documentation: Data Validation occurs within the entry screens of the database. Data Checking occurs with a Voice Reports software package that reads what has been entered in the database and allows the data enterer to check the records that are being read out against the records written on the datasheets.
Data storage
National database: Pelagic Discards Database (Access) International database: Data (unraised) submitted to ICES in request to annual data call. Quality checks and data validation documentation: Data Validation occurs within the entry screens of the database. Input is constrained by the use of drop-down lists with further checks performed by the DB application code. Data is further checked during extractions for end-users such as ICES / European Commission. The checks used will depend on the use of the data. Migration to SQL Server will be undertaken following completion of the design and specification phases. The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage Fish samples are processed in Marine Institute laboratories for length, weight and maturity analysis. Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for processing and analysis. These age structures are stored at Marine Institute premises for a period of months before preparation and age reading is carried out. Raw data sheets are kept in the regional laboratories and then transferred to headquarters where they are filed. Raw data sheets, duplicate otoliths and read otoliths are archived offsite in secure storage areas after three years. Sectioned otoliths and illicia are stored on site in specialised file storage units.
Data processing Evaluation of data accuracy (bias and precision): No stand-alone evaluation of bias and precision of the data collected by this scheme. Any data collected is submitted to end users at which point accuracy will be assessed. Editing and imputation methods: Y Where errors are identified, the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during transcription. Imputation of unsampled domains of interest often done at the international level (e.g. ICES), rather than at the national level. Quality document associated to a dataset: No DOI is currently created for the dataset since it is not publicly available. Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g., in the case of input error) or excluded from that particular use.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan. <u>Description of the Population</u> No Deviations <u>Sampling Design and Protocols</u> Sampling was increased by the introduction of -At Sea Self Sampling program where skippers collected data according to the appointed SOP on a daily basis and delivered to the MI. This data was quality controlled live on a daily basis via WhatsApp communication with the fleet. The sea self-sampling augments the sampler at sea data collected <u>Sampling Implementation</u>

The PSUs achieved were 50%. The opportunity to sample was limited due to low TAC and tight management of the season. Extra sampling where accommodation was limited was achieved via the at sea self-sampling in 2023.

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: Pelagic at - sea Enhanced

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: Pelagic at-sea Enhanced
Sampling scheme type: Commercial fishing trip
Observation type: SciObsAtSea
Time period of validity: 2021-2027
Description of the population
Population targeted: The enhanced sampling specifically targets fisheries determined to be higher risk to bycatch. The target population is the group of vessels that are engaged in the target mackerel, horse mackerel, and blue whiting fisheries. Population sampled: All vessels that reported landings in the same semester of the previous year are included in the sampling frame. Stratification: The sampling frames are stratified by semester.
Sampling design and protocols

Sampling design description:

The PSU is vessel*time. The sampling frame is a list of vessels active in the target fisheries for mackerel, horse mackerel, and blue whiting, during the reference period. Each pelagic vessel has a sampling probability based on average landings during the relevant semester over the reference period. Vessels are sampled from the list with replacement. Sampling is carried out by a sampler on-board for the duration of the fishing trip.

Rare/incidental bycatch of fish species are checked during each sampling event.

Any Bird/Mammal/Reptile/PET/Decomposed organism that comes in contact with the gear during fishing operation is also recorded. In addition, associated information on each bycatch event are recorded, including haul number, gear used, whether the animal(s) is released alive or deceased, number of animals, condition of the animal(s), sex, and length.

VME indicator species are noted if present in the random sample box

Is the sampling design compliant with the 4S principle? Y

Regional coordination: N

Link to sampling design documentation:

<https://www.dcmapp-ireland.ie/documents/methodologies>

Compliance with international recommendations: Y. The enhanced bycatch sampling programme is designed to supplement the regular bycatch sampling occurring under DCF. This sampling program has been designed in accordance with guidance from the ICES WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints. The recommendations of STECF included an increase in monitoring of métiers with a high risk of protected species bycatch. As such, based on a risk assessment of Irish fisheries, high risk fisheries for the fleet were identified and have been targeted by this enhanced sampling scheme.

Link to sampling protocol documentation:

https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/At-Sea%20Sampling%20Manual_2020.pdf

Compliance with international recommendations: Y.

The sampling protocols have been designed in accordance with guidance from the ICES

WKPICS workshops and WGCATCH working group. It has also been necessary to take into account national logistics and constraints.
Sampling implementation
Recording of refusal rate: Y – Refusals and analysis of refusal rates are in accordance with guidance from SGPIDS, 2013. Monitoring of sampling progress within the sampling year: Sampling progress against targets is tracked within the database and reports of sampling targets and achievements are available to samples.
Data capture
Means of data capture: Data on bycatch is initially recorded on paper and then transferred to a database application as soon as possible after sampling. All bird, mammal, reptile (fresh or decomposed) that comes into contact with the gear during fishing operations (either collected on deck or falling out during hauling) is recorded. Information on the bycaught species is recorded in the data sheets, if no bycatch is recorded this information is also noted to ensure all true zero samples are recorded and clear. Information recorded includes, haul number, gear used, whether the animal(s) is released alive or deceased, number of animals, condition of the animal(s), sex, and length. Data capture documentation: A recent copy of the SOP for completing data sheets is available at https://www.dcmapi-ireland.ie/sites/default/files/DCF_Files/docs/SOP_Sampling%20Demersal%20and%20Pelagic%20Form%20filling_0.docx Quality checks documentation: Y Data Validation occurs within the entry screens of the database. Data Checking occurs with a Voice Reports software package that reads what has been entered in the database and allows the data enterer to check the records that are being read out against the records written on the datasheets.
Data storage
National database: Pelagic Discards Database (Access), bycatch data are also summarised

and stored in a bycatch specific database. International database: Detailed data is submitted to RDB / RDBES. Raised data is also currently submitted to the ICES InterCatch database. Bycatch data on seabirds, cetaceans, pinnipeds and PET species fish are submitted to ICES WGBYC data call annually, and to any additional ICES or OSPAR special request data calls relating to bycatch, such as WKMOMA. Quality checks and data validation documentation: Y Data Validation occurs within the entry screens of the database. Input is constrained by the use of drop-down lists with further checks performed by the DB application code. Data is further checked during extractions for end-users such as ICES / European Commission. The checks used will depend on the use of the data. Migration to SQL Server will be undertaken following completion of the design and specification phases. The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).
Sample storage Raw data sheets are kept in the regional laboratories and then transferred to headquarters where they are filed. Raw data sheets, duplicate otoliths and read otoliths are archived offsite in secure storage areas after three years. Sectioned otoliths and illicia are stored on site in specialised file storage units. Any retained fish samples are processed in Marine Institute laboratories for length, weight and maturity analysis. Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for processing and analysis. These age structures are stored at Marine Institute premises for a period of months before preparation and age reading is carried out.
Data processing
Evaluation of data accuracy (bias and precision): Y No a stand-alone evaluation of bias and precision of the data collected by this scheme. Any data collected is submitted to end users at which point accuracy will be assessed.

Editing and imputation methods: Y

Where errors are identified, the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during transcription.

Imputation of unsampled domains of interest often done at the international level (e.g. ICES), rather than at the national level.

Quality document associated to a dataset:

No DOI is currently created for the dataset since it is not publicly available.

Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g., in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No Deviations

Sampling Design and Protocols

No Deviations

Sampling Implementation

No Deviations

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: *Nephrops* at - sea self-sampling, *Nephrops* vessels in FU16

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: <i>Nephrops</i> at-sea self-sampling, <i>Nephrops</i> vessels in FU16
Sampling scheme type: Commercial fishing trip
Observation type: SciObsOnShore
Time period of validity: 2020-ongoing
Description of the population
Population targeted: All commercial catch fractions from the <i>Nephrops</i> fisheries landed into Ireland, from the primary <i>Nephrops</i> Functional Units (FU16) fished by Irish registered vessels. Population sampled: 100% of the <i>Nephrops</i> landings are covered by the sampling program. All vessel classes > 10 metres and only <i>Nephrops norvegicus</i> are included in the sampling program. Stratification: Sampling events are stratified by FU/vessel/year/month. <i>Nephrops</i> grounds are geo-referenced by FU16.
Sampling design and protocols
Sampling design description: The Primary Sampling Unit (PSU) is vessel*month. Targets for number of vessels to be sampled are set for each FU/month – these are proportional to the landings from the relevant reference period. Samples consist of one box of unsorted catch and one box of discards taken from three random valid fishing tows and brought ashore for subsequent work up from one PSU. A valid fishing tow is when no fishing gear problems encountered during fishing tow. Fishing vessels are contacted and asked to bring the samples ashore. Three samples brought ashore from each fishing trip sampled. Is the sampling design compliant with the 4S principle?: Y Regional coordination: N Link to sampling design documentation: https://www.dcmapp-

ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Nephrops_SOP_2016.pdf

Compliance with international recommendations: Y. The sampling programme is designed to gather samples covering *Nephrops* FU16 from the fleets fishing each. These quasi-reference fleets are sampled to the extent of one sample per 50 tonnes total landings, directed according to the recent three-year average. It has also been necessary to take into account national logistics and constraints.

Link to sampling protocol documentation:

https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/At-Sea%20Sampling%20Manual_2020.pdf

Documentation is stored on national server and will be transferred to the Marine Institutes “Paradigm3”, SOP document management repository in 2022.

Sampling implementation

Recording of refusal rate: N – since samplers decide which vessels to sample using expert judgement the refusal rate is not relevant for this sampling program. Refusals would only occur extremely rarely in this sampling programme in any case.

Monitoring of sampling progress within the sampling year: Sampling planning and progress against targets is tracked and sampling achievements are available to samplers and are automatically updated. If the targets are not being met, then targeted sampling may occur.

Data capture

Means of data capture:

Nephrops catch weights, length frequency distributions and biological data are captured using digital electronic callipers and marine scales connected to a tablet and stored locally on the tablet and then uploaded to a SQL server database “Nemesys” as soon as possible after sampling.

Data capture documentation:

SOPs for sampling (weighing / measuring etc.) are held locally on the network and will be transferred to the Marine Institutes “Paradigm3” document management repository, to be reviewed and updated as necessary, in early 2022.

Quality checks documentation: Y

Data are QC'd during the “Nemesys” database collection process.

Data storage


National database: FEAS Nemesys

Stored in secure database on Marine Institute IT servers with regular backups following the Institutes IT protocols.

International database: Detailed data will be submitted to ICES RDB / RDBES. Raised data is currently submitted to the ICES InterCatch database.

Quality checks and data validation documentation: Y

Data Quality QC functions are available within “Nemesys” – these cover for example, length-weight, length-frequency distributions.



Data are also checked during extractions for end-users such as ICES / European Commission. The checks used depend on the intended use of the data.

The data management of this data collection activity is incorporated into the Marine Institute’s IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

Samples are routinely labelled and frozen in freezers in port laboratories before work-up.

Data are initially saved to Local “Nemesys” databases on tablets, which are subsequently uploaded to the master Nemesys SQL server soon after processing of a sample.

Additionally, Local Nemesys database on tablets are backed up to the Marine Institute network.

Data processing

Evaluation of data accuracy (bias and precision): Y

There is not a stand-alone evaluation of bias and precision of the data collected by this

scheme but it has been evaluated in recent ICES benchmarks including:

Report of the Benchmark Workshop on *Nephrops* Stocks (WKNEPH)2013
https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2013/WKNEPH%202013/wkneph_2013.pdf

Report of the Benchmark Workshop on Celtic Sea stocks (WKCELT) 2014
<https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2014/WKCELT/WKCELT%20Final%20Report.pdf>

Hans Gerritsen, Jennifer Doyle and Colm Lordan, 2006. An Evaluation of the Precision of length-frequency samples of *Nephrops* from the Western Irish Sea (FU 15), Working Document to the Workshop on *Nephrops* Stocks (WKNEPH), 24–27 January 2006, ICES Headquarters. ICES CM 2006/ACFM:12. 85 pp.
<https://www.ices.dk/sites/pub/CM%20Ddocuments/2006/ACFM/ACFM1206.pdf>

Editing and imputation methods: Y

Where errors are identified, the preferred action is to correct the errors in the database.

Imputation of unsampled domains of interest are often done at the international level (e.g. ICES), rather than at the national level.

Quality document associated to a dataset:

No DOI is currently created for the dataset since it is not publicly available.

R markdown documents recording the estimation and imputation steps performed for ICES data calls are available on request.

Validation of the final dataset: Data are checked during extractions for end-users such as ICES / European Commission - the checks used depend on the type and intended use of the data. If errors or anomalies are observed, then data are either corrected by reference to the database (e.g. in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No Deviations

Sampling Design and Protocols

No Deviations

Sampling Implementation

No Deviations

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

Uploader data facility from sampling tablet devices to the SQL server database has been updated to concur with stringent IT policies.

Data Processing

No Deviations

Sampling scheme identifier: *Nephrops* at - sea, *Nephrops* vessels in FU16

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: <i>Nephrops</i> at-sea, <i>Nephrops</i> vessels in FU16
Sampling scheme type: Commercial fishing trip
Observation type: SciObsAtSea
Time period of validity: 2011 – 2027
Description of the population
Population targeted: The target population is the group of vessels that are engaged in demersal fisheries (i.e. catching of demersal fish and <i>Nephrops</i> using bottom contacting gears) of the FU16, Porcupine Bank <i>Nephrops</i> Grounds only.
Population sampled: Primary vessels that have historically reported <i>Nephrops</i> landings are included in the quasi-reference fleet. Each undertakes on average 3 to 5 fishing trips per year to FU16.
Stratification: The quasi-reference fleets are not stratified owing to their low number, and are sampled as and when availability allows according to reasonable logistics and constraints.
Sampling design and protocols
Sampling design description: The Primary Sample Unit (PSU) is vessel*time. The sampling frame is a quarterly list of vessels that were active in the same quarter of the

previous year. A sampling target of 3 trips is defined per quarter that the FU is open (historically FU16 has been closed to fishing from the end of June to the start of October). Rare/incidental bycatch of fish species are checked during each sampling event. Any Bird/Mammal/Reptile/PET/Decomposed organism that comes in contact with the gear during fishing operation is also recorded. VME indicator species are noted if present in the random box of discards

Is the sampling design compliant with the 4S principle?: Y

Regional coordination: N

Link to sampling design documentation:

https://www.dcmapi-ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Nephrops_SOP_2016.pdf

Compliance with international recommendations: Y. The sampling programme is designed to gather samples from the fleet fishing *Nephrops* FU16. It has been necessary to take into account national logistics and constraints.

Link to sampling protocol documentation:

Documentation is stored on national server and will be transferred to the Marine Institutes “Paradigm3”, SOP document management repository in 2022.

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: N – Since samplers decide which vessels to sample using expert judgement the refusal rate is not relevant for this sampling program. Refusals would only occur extremely rarely in this sampling programme in any case.-

Monitoring of sampling progress within the sampling year: Sampling planning and progress against targets is tracked and sampling achievements are available to samplers. If the targets are not being met, then targeted sampling may occur.

Data capture

Means of data capture:

Nephrops catch weights, length frequency distributions and biological data are captured using digital electronic callipers and marine scales connected to a tablet and stored locally in wet laboratory and then uploaded to a SQL server database “Nemesys” as soon as possible after sampling.

Data capture documentation:

SOPs for sampling (weighing / measuring etc.) are held locally on the network and will be transferred to the Marine Institutes “Paradigm3” document management repository, to be reviewed and updated as necessary, in early 2022.

Quality checks documentation: Y

Data are QC’d during the “Nemesys” database collection process.

Data storage


National database: FEAS Nemesys

Stored in secure database on Marine Institute IT servers with regular backups following the Institutes IT protocols.

International database: Detailed data will be submitted to ICES RDB / RDBES. Raised data is currently submitted to the ICES InterCatch database.

Quality checks and data validation documentation: Y

Data Quality QC functions are available within “Nemesys” – these cover length-weight, length-frequency distributions e.g.



Data are also checked during extractions for end-users such as ICES / European Commission. The checks used will depend on the intended use of the data.

The data management of this data collection activity is incorporated into the Marine Institute’s IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

Samples are routinely labelled and frozen in freezers in port laboratories before work-up.

Data are initially saved to Local “Nemesys” databases on tablets, which are subsequently uploaded to the master Nemesys SQL server soon after processing of a sample.

Additionally, Local Nemesys database on tablets are backed up to the MI network.

Data processing

Evaluation of data accuracy (bias and precision): Y

There is not a stand-alone evaluation of bias and precision of the data collected by this scheme but it has been evaluated in recent ICES benchmarks including:

Report of the Benchmark Workshop on *Nephrops* Stocks (WKNEPH)2013

https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2013/WKNEPH%202013/wkneph_2013.pdf

Hans Gerritsen, Jennifer Doyle and Colm Lordan, 2006. An Evaluation of the Precision of length-frequency samples of *Nephrops* from the Western Irish Sea (FU 15), Working Document to the Workshop on *Nephrops* Stocks (WKNEPH), 24–27 January 2006, ICES Headquarters. ICES CM 2006/ACFM:12. 85 pp.
<https://www.ices.dk/sites/pub/CM%20Documents/2006/ACFM/ACFM1206.pdf>

Editing and imputation methods: Y

Where errors are identified, the preferred action is to correct the errors in the database. Imputation of unsampled domains of interest are often done at the international level (e.g. ICES), rather than at the national level.

Quality document associated to a dataset:

No DOI is currently created for the dataset since it is not publicly available.

R markdown documents recording the estimation and imputation steps performed for ICES data calls are available on request.

Validation of the final dataset: Data are checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the intended use of the data. If errors or anomalies are observed, then data are either corrected by reference to the original data sheets (e.g. in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No Deviations

Sampling Design and Protocols

No Deviations

Sampling Implementation

No Deviations

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

Uploader data facility from sampling tablet devices to the SQL server database has been updated to concur with stringent IT policies.

Data Processing

No Deviations

Sampling scheme identifier: *Nephrops* On - Shore, *Nephrops* vessels excluding FU16

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: <i>Nephrops</i> On-Shore, <i>Nephrops</i> vessels excluding FU16
Sampling scheme type: Commercial fishing trip
Observation type: SciObsOnShore
Time period of validity: 2001-ongoing
Description of the population
Population targeted: All commercial catch fractions from the <i>Nephrops</i> fisheries landed into Ireland, from the primary <i>Nephrops</i> Functional Units (FU) fished by Irish registered vessels (or for fisheries where a bi-lateral agreement is in place). Population sampled: 98% of the <i>Nephrops</i> landings are covered by the sampling program. All vessel classes > 10 metres and only <i>Nephrops norvegicus</i> are included in the sampling program. FUs 11 to 14 are not routinely sampled owing to low levels of national participation in these fisheries. Stratification: Sampling events are stratified by FU/vessel/year/month. <i>Nephrops</i> grounds are geo-referenced by FU (FU15, FU17, FU19, FUs20 and 21 combined, and FU22).
Sampling design and protocols
Sampling design description: The Primary Sampling Unit (PSU) is vessel*month. Targets for number of vessels to be sampled are set for each FU/month – these are proportional to the landings from the relevant reference period. Samples consist of one box of catch and one box of discards taken from any one haul and brought ashore for subsequent work up. Is the sampling design compliant with the 4S principle?: Y Regional coordination: N Link to sampling design documentation: https://www.dcmaph-ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Nephrops_SOP_2016.pdf Compliance with international recommendations: Y. The sampling programme is designed to gather samples covering all <i>Nephrops</i> FUs from the fleets fishing each. These quasi-reference fleets are sampled to the extent of one sample per 50 tonnes total landings, directed according to the recent three-year average. It has also been necessary to take into account national logistics and constraints. Link to sampling protocol documentation: https://www.dcmaph-ireland.ie/sites/default/files/DCF_Files/docs/At-Sea%20Sampling%20Manual_2020.pdf Documentation is stored on national server and will be transferred to the Marine Institutes “Paradigm3”, SOP document management repository in 2022.
Sampling implementation

Recording of refusal rate: N – since samplers decide which vessels to sample using expert judgement the refusal rate is not relevant for this sampling program. Refusals would only occur extremely rarely in this sampling programme in any case.

Monitoring of sampling progress within the sampling year: Sampling planning and progress against targets is tracked and sampling achievements are available to samplers and are automatically updated. If the targets are not being met, then targeted sampling may occur.

Data capture

Means of data capture:

Nephrops catch weights, length frequency distributions and biological data are captured using digital electronic callipers and marine scales connected to a tablet and stored locally on the tablet and then uploaded to a SQL server database “Nemesys” as soon as possible after sampling.

Data capture documentation:

SOPs for sampling (weighing / measuring etc.) are held locally on the network and will be transferred to the Marine Institutes “Paradigm3” document management repository, to be reviewed and updated as necessary, in early 2022.

Quality checks documentation: Y

Data are QC'd during the “Nemesys” database collection process.

Data storage



National database: FEAS Nemesys

Stored in secure database on Marine Institute IT servers with regular backups following the Institutes IT protocols.

International database: Detailed data will be submitted to ICES RDB / RDBES. Raised data is currently submitted to the ICES InterCatch database.

Quality checks and data validation documentation: Y

Data Quality QC functions are available within “Nemesys” – these cover for example, length-weight, length-frequency distributions.



Data are also checked during extractions for end-users such as ICES / European Commission. The checks used depend on the intended use of the data.

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

Samples are routinely labelled and frozen in freezers in port laboratories before work-up.

Data are initially saved to Local "Nemesys" databases on tablets, which are subsequently uploaded to the master Nemesys SQL server soon after processing of a sample.

Additionally, Local Nemesys database on tablets are backed up to the Marine Institute network.

Data processing

Evaluation of data accuracy (bias and precision): Y

There is not a stand-alone evaluation of bias and precision of the data collected by this scheme but it has been evaluated in recent ICES benchmarks including:

WKNEPH 2013

https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2013/WKNEPH%202013/wkneph_2013.pdf

WKCELT 2014

<https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2014/WKCELT/WKCELT%20Final%20Report.pdf>

IBPNeph 2015 https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2015/IBP%20Neph/IBPNeph_2015.pdf

Hans Gerritsen, Jennifer Doyle and Colm Lordan, 2006. An Evaluation of the Precision of length-frequency samples of *Nephrops* from the Western Irish Sea (FU 15), Working Document to the Workshop on *Nephrops* Stocks (WKNEPH), 24–27 January 2006, ICES Headquarters. ICES CM 2006/ACFM:12. 85 pp.

Editing and imputation methods: Y

Where errors are identified, the preferred action is to correct the errors in the database.

Imputation of unsampled domains of interest are often done at the international level (e.g. ICES), rather than at the national level.

Quality document associated to a dataset:

No DOI is currently created for the dataset since it is not publicly available.

R markdown documents recording the estimation and imputation steps performed for ICES data calls are available on request.

Validation of the final dataset: Data are checked during extractions for end-users such as ICES / European Commission - the checks used depend on the type and intended use of the data. If errors or anomalies are observed, then data are either corrected by reference to the database (e.g. in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No Deviations

Sampling Design and Protocols

No Deviations

Sampling Implementation

No Deviations

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

Uploader data facility from sampling tablet devices to the SQL server database has been updated to concur with stringent IT policies.

Data Processing

No Deviations

Sampling scheme identifier: Crustacea at-sea

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: Crustacea at-sea
Sampling scheme type: Commercial fishing trip
Observation type: SciObsAtSea
Time period of validity: 2013-ongoing
European lobster (<i>Homarus gammarus</i>) and Brown crab (<i>Cancer pagurus</i>) catches are sampled on board commercial vessels around the coast of Ireland through an at sea observer programme with sampling trips occurring on an ad-hoc basis during the 6-9 months that the fisheries take place. A Sentinel Vessel Programme and a skipper self-sampling programme, where skippers keep records of daily catches of lobster and brown crab runs parallel to the observer programme.
Description of the population
Population targeted: All commercial catch fractions from the European lobster (<i>Homarus gammarus</i>) and Brown crab (<i>Cancer pagurus</i>) fisheries around in the Irish coast in ICES Areas 6 and 7.
Population sampled: <1% of the total number of fishing trips of vessels >12 meters are sampled.
Stratification: Sampling events are stratified by geographic region (north-west / west / south-

west / south east / east).
Sampling design and protocols
Sampling design description: The PSU is vessel-trip. Targets for number of trips undertaken are set for each region. Is the sampling design compliant with the 4S principle?: N Regional coordination: N Link to sampling design documentation: N Sampling of lobster and brown crab catches from ICES Areas 6 and 7 is carried out opportunistically for the 6-9 months of the season. Compliance with international recommendations: N Link to sampling protocol documentation: An SOP for the at sea sampling (measuring, sexing and recording data) are held in Paradigm 3 (a document management system) and are reviewed and updated regularly. Compliance with international recommendations: N
Sampling implementation
Recording of refusal rate: N Monitoring of sampling progress within the sampling year: Sampling progress against targets is tracked throughout the fishing season.
Data capture
Means of data capture: Data is initially recorded on paper and is then input into an excel template prior to being transferred to a database application as soon as possible after sampling. Size of lobster and crab is measured using callipers and measuring boards. Data capture documentation: SOPs for sampling (measuring /sexing/recording data) are held in Paradigm3 (a document management system) and reviewed and updated regularly. Quality checks documentation: Y Data validation checks and database validation are carried out along with visual inspection of outliers.
Data storage
National database: FEAS_InshoreFisheries International database: NA Quality checks and data validation documentation: Y Data Quality QC functions are available within an excel spreadsheet that is used to input the data. Data is also checked during extractions for end-users. The checks used will depend on the use of the data. The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).
Sample storage

Storage description: NA
Data processing
Evaluation of data accuracy (bias and precision): Y The precision of the data is reported in the catch index of our assessments. Editing and imputation methods: Y Where errors are identified then the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during transcription. Quality document associated to a dataset: No DOI is currently created for the dataset since it is not publicly available. Validation of the final dataset: Data is checked during extractions for end-users - the checks used will depend on the use of the data. If errors or anomalies are observed then data is either corrected by reference to the original data sheets (e.g. in the case of input error) or excluded from that particular use.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan. <u>Description of the Population</u> No Deviations <u>Sampling Design and Protocols</u> No Deviations <u>Sampling Implementation</u> No Deviations <u>Data Capture</u> No Deviations <u>Data Storage</u> No Deviations <u>Sample Storage</u> No Deviations <u>Data Processing</u> No Deviations

Sampling scheme identifier: Crustacea On - Shore

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: Crustacea On-Shore

Sampling scheme type: Commercial fishing trip
Observation type: SciObsOnShore
Time period of validity: 2013-ongoing
Landings, by Irish vessels, of the European lobster (<i>Homarus gammarus</i>) and Brown Crab (<i>Cancer pagurus</i>) are sampled on a monthly basis, where possible, at various processing facilities in the northwest, west and southwest of Ireland, during the 6-9 months of the fishing season. Overall <1% of vessel trips are sampled.
Description of the population
Population targeted: The PSU is port*day. All commercial landings (HUC) from the European lobster (<i>Homarus gammarus</i>) and Brown crab (<i>Cancer pagurus</i>) fisheries landed into Ireland from ICES Areas 6 and 7.
Population sampled: <1% of the lobster and brown crab landings from ICES Areas 6 and 7 are covered by the sampling program.
Stratification: Sampling events will occur on a monthly basis in 3 regions undertaken in the north-west, west and south-west.
Sampling design and protocols
Sampling design description: The PSU is port-day. The PSU is port-day. Ports/processing facilities are targeted based on location of where the majority of landings occur. Approximately 20% of <i>Homarus gammarus</i> and <i>Cancer pagurus</i> landings will be sampled.
Is the sampling design compliant with the 4S principle?: N
Regional coordination: N
Link to sampling design documentation: N
Sampling of lobster and brown crab landings from ICES Areas 6 and 7 is carried out on a monthly basis, where possible. The sampling is stratified into 3 regions and sampling occurs at processing facilities in the northwest, west and southwest of Ireland.
Compliance with international recommendations: N
Link to sampling protocol documentation:
An SOP for sampling (collecting, measuring, weighing and recording data) are held in Paradigm 3 (a document management system) and are reviewed and updated regularly. https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/1.0%20FEAS%20Inshore%20Sampling%20Overview.docx
Compliance with international recommendations: N
Sampling implementation
Recording of refusal rate: N
Samplers decide which port-days to sample by liaising with the processing facility and using expert judgement the refusal rate is not relevant for this sampling program.
Monitoring of sampling progress within the sampling year: Sampling progress against targets is tracked throughout the year.
Data capture
Means of data capture:
Data is initially recorded on paper and is then input into an excel template to be transferred to a database application as soon as possible after sampling. Data is measured using callipers or measuring boards and scales.
Data capture documentation:
SOPs for sampling (weighing / measuring /) are held in Paradigm3 (a document management system) and reviewed and updated regularly. https://www.dcmapp-

ireland.ie/sites/default/files/DCF_Files/docs/FEAS_Inshore_3_PortSampling.pdf Quality checks documentation: Y Data validation checks and database validation are carried out along with visual inspection of outliers.
Data storage
National database: FEAS_Inshore International database: NA Quality checks and data validation documentation: Y Data Quality QC functions are available within an excel spreadsheet that is used to input the data. Data is also checked during extractions for end-users. The checks used will depend on the use of the data. The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).
Sample storage
Storage description: N/A
Data processing
Evaluation of data accuracy (bias and precision): N There is no stand-alone evaluation of bias and precision of the data collected by this scheme. Editing and imputation methods: Y Where errors are identified then the preferred action is to correct the errors in the database – this might involve reference to the original data sheets if the data has occurred during transcription. Quality document associated to a dataset: No DOI is currently created for the dataset since it is not publicly available. Validation of the final dataset: Data is checked during extractions for end-users - the checks used will depend on the use of the data. If errors or anomalies are observed then data is either corrected by reference to the original data sheets (e.g. in the case of input error) or excluded from that particular use.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.
<u>Description of the Population</u> No Deviations
<u>Sampling Design and Protocols</u> No Deviations
<u>Sampling Implementation</u> No Deviations
<u>Data Capture</u> No Deviations

<u>Data Storage</u>
No Deviations
<u>Sample Storage</u>
No Deviations
<u>Data Processing</u>
No Deviations

Sampling scheme identifier: Mollusca On - Shore

MS: IRL
Region: North-East Atlantic
Sampling scheme identifier: Mollusca On-Shore
Sampling scheme type: Commercial fishing trip
Observation type: SciObsOnShore
Time period of validity: 2002-ongoing
Landings, by Irish vessels, of the King Scallop (<i>Pecten maximus</i>) and Common whelk (<i>Buccinum undatum</i>) are sampled on a monthly basis, where possible. King Scallop are sampled at a processing facility in the southeast of Ireland, while Whelk are sampled at processing facilities in the northwest and southeast. Overall <1% of vessel trips are sampled.
Description of the population
Population targeted: The PSU is port*day. All commercial landings (HUC) from the King Scallop (<i>Pecten maximus</i>) and Common whelk (<i>Buccinum undatum</i>) fisheries landed into Ireland from ICES Areas 6 and 7. Population sampled: <1% of the scallop and whelk landings from ICES Areas 6 and 7 are covered by the sampling program. Stratification: Sampling events will occur on a monthly basis on the south east coast where over 90% of landings occur and where possible on the north west coast for whelk.
Sampling design and protocols
Sampling design description: The PSU is port-day. Ports/processing facilities are targeted based on location of where the majority of landings occur and >80% of <i>Pecten maximus</i> and <i>Buccinum undatum</i> landings will be sampled. Is the sampling design compliant with the 4S principle?: N

Regional coordination: N Link to sampling design documentation: N Sampling of scallop and whelk landings from ICES Areas 6 and 7 is carried out on a monthly basis, where possible. The sampling is stratified into 2 regions and sampling occurs at processing facilities in the northwest and southwest of Ireland. Frequency of sampling also depends on the pattern of commercial landings. Compliance with international recommendations: N Link to sampling protocol documentation: An SOP for sampling (collecting, measuring, weighing and recording data) are held in Paradigm 3 (a document management system) and are reviewed and updated regularly. https://www.dcmmap-ireland.ie Compliance with international recommendations: N
Sampling implementation
Recording of refusal rate: N Samplers decide which port-days to sample by liaising with the processing facility and using expert judgement the refusal rate is not relevant for this sampling program. Monitoring of sampling progress within the sampling year: Sampling progress against targets is tracked throughout the year.
Data capture
Means of data capture: Data is initially recorded on paper and is then input into an excel template to be transferred to a database application as soon as possible after sampling. Data is measured using callipers or measuring boards and scales. Data capture documentation: SOPs for sampling (weighing / measuring /) are held in Paradigm3 (a document management system) and reviewed and updated regularly. https://www.dcmmap-ireland.ie

Quality checks documentation: Y Data validation checks and database validation are carried out along with visual inspection of outliers.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan. <u>Description of the Population</u> No Deviations <u>Sampling Design and Protocols</u> Whelk (<i>Buccinum undatum</i>) have only been sampled in the southeast since 2022, however whelk from the northwest can be sampled at the facility along with whelk caught off the east coast of Ireland. <u>Sampling Implementation</u> No Deviations <u>Data Capture</u> No Deviations <u>Data Storage</u> No Deviations <u>Sample Storage</u> No Deviations <u>Data Processing</u> No Deviations

Quality Reports Related to Table 2.3 and Text Box 2.3

Sampling scheme identifier: Diad_ESB_Counter (*Salmo salar*)

MS : Ireland
Region: North East Atlantic
Sampling scheme identifier: Diad_ESB_Counter (<i>Salmo salar</i>)

Sampling scheme type: Diadromous (Scientific)
Observation type: EMA water body
Time period of validity: from when until when 2022-2027
Short description (max 100 words): Species: <i>Salmo salar</i> Sampling Scheme Type: Diadromous (Scientific) Sampling Scheme Identifier: Diad_ESB_Counter Fixed permanent counter upstream and downstream (n=2) monitor salmon and and kelt moving downstream, as well as adult salmon moving upstream enabling full census on wild salmon and released reared salmon. These are situated at or closed to the tidal limit and are fixed to natural or hydropower barriers or weirs on the Shannon Ardnacrusha and Parteen (IE_Sha) and Erne (IE_NorW) and Clady. These counters provide a valuable time series of relative abundance of wild salmon smolts/adults and released reared salmon smolts/adults. Counters provide annual index recruitment abundance data for ICES WGNAS datacalls and in WGNAS assessment model.
Description of the population
Population targeted: Target Species: <i>Salmo salar</i> Stocks targeted are the total stock (downstream migrating recruits and upstream migrating returning adults) for both species in the Erne, Clady, Shannon and Lee river. Population sampled: Specify which part of the target population will be sampled and specify which part of the target population is unreachable for sampling or excluded for some reason to explain, <i>e.g. major ports being listed as auctions excluding all minor ports and no sampling during the week-ends</i>. For research surveys at sea describe target species in single-species surveys or ecosystem component (<i>e.g. demersal, pelagic</i>) in multispecies surveys. Smolts and Adults. Stratification: These rivers (Erne IE_NorW, Shannon IE_Sha, Lee IE_SouW) were identified as having

major hydropower stations at the tidal limit.
Rivers were ESB operate hydropower stations.
Sampling design and protocols
Sampling design description: Describe how the sampling allocation is defined; how PSU and SSU are selected for sampling; indicate for which catch fraction the sampling scheme applies. Recruitment data are continual census data Recruitment data (smolt) are continual census data at fixed counter locations at the tidal limit in the Erne, Clady, Shannon and Lee Rivers. Adult upstream count data are continual census data at fixed counter locations at the tidal limit in the Erne, Clady, Shannon and Lee Rivers. Is the sampling design compliant with the 4S principle?: Y/N/NA (NA for e.g. surveys and diadromous and recreational sampling schemes) NA Regional coordination: Indicate if the sampling design and protocols were developed as part of a regional or multi-lateral agreement, and if yes, refer to the agreement (table 1.3) and list all MS participating. No regional co-ordination. Link to sampling design documentation: It is managed by the Irish Electricity Supply Board (ESB). https://www.esb.ie/docs/default-source/investor-relations-documents/esb-annual-report-2013 Compliance with international recommendations: Member State shall state ‘Y’ (yes) if the sampling protocol is in line with international recommendations, and ‘N’ if not. If no relevant expert or coordination groups exist, the sampling protocol should be shortly explained in the text, and should be available upon request for the evaluators. N, currently in discussion in WGNAS. Has been ongoing discussion for 4 years now.
Sampling implementation
Recording of refusal rate: NA Monitoring of sampling progress within the sampling year: Indicate how sampling allocations are adjusted (if needed) and followed-up, what are the mechanisms in place to

resolve issues and adopt mitigation measures during the sampling year? Oversight by the Technical Expert Group on Salmon (TEGOS). The total catches are what is caught – there is no sub-sampling – its census data or total catch data. Sampling is monitored at end of season and adjusted for next season Biological sampling rates can be adjusted as well
Data capture
Means of data capture: short description (+ photo optionally). Indicate what are the means for collecting the data, e.g. scales, measuring board, dedicated software, ... Field data are initially recorded by hand on field sheets or fisher's logs, and subsequently transcribed to excel spreadsheets. Data capture documentation: https://www.esb.ie/docs/default-source/investor-relations-documents/esb-annual-report-2013 https://www.fisheriesireland.ie/sites/default/files/migrated/docman/The%20Status%20of%20Irish%20Salmon%20Stocks%20in%202020%20with%20Catch%20Advice%20for%202021.pdf Quality checks documentation: Indicate with 'Y' (yes) or 'N' (no). If 'N' (no), indicate when (year) documentation will be available. Provide a link to a webpage where the documentation can be found. If no link is available, but documentation exists, provide a literature reference (author(s), year and type of publication - e.g. internal report). If no documentation on the quality checks exists, provide some details in the text box. Visual checks are applied to the data at each stage. Once data is on spreadsheet, graphical checks are used to identify outliers. These are either changed with cross checks with the paper records and data providers, or deleted where verification is not possible. Fishers logs and reported catch data are followed up if data is not consistent with expectations by area managers.

Data storage
National database: Recruitment data are uploaded annually with TEGOS. International database: Provide the name of international database(s) and the organisation hosting the database, if applicable. Otherwise, insert 'NA' (not applicable). Provide a link if the database is accessible through a website. ICES WGNAS database. Quality checks and data validation documentation: Not currently available. WGNAS on going work on this: https://www.ices.dk/sites/pub/Publication%20Reports/Forms/DispForm.aspx?ID=37558
Sample storage
Storage description: N/A – no biological samples taken from Adults or Smolts.
Data processing
Evaluation of data accuracy (bias and precision): No formal evaluation, data used by WGNAS for years (since early 1980s). No issues have been identified. Editing and imputation methods: N Quality document associated to a dataset: No. There isn't a DoI publication of all the data.

Data are summarised and published in the Annual Technical Expert Group on Salmon Reports

<https://www.fisheriesireland.ie/sites/default/files/migrated/docman/The%20Status%20of%20Irish%20Salmon%20Stocks%20in%202020%20with%20Catch%20Advice%20for%202021.pdf>

Validation of the final dataset:

Data are quality checked during capture and transcription, the annual dataset is then manually / graphically checked for correctness and consistency, the data undergoes further checks upon upload to the WGNAS database, during WGNAS meeting. Spurious unverified data is discarded and not used in any stock assessments.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No Deviations

Sampling Design and Protocols

No Deviations

Sampling Implementation

No Deviations

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: Diad_ESB_Eel (*Anguilla anguilla*)

MS : Ireland

Region: North East Atlantic

Sampling scheme identifier: Diad_ESB_Eel (<i>Anguilla anguilla</i>)
Sampling scheme type: Diadromous (Scientific)
Observation type: SciObs water body
Time period of validity: from when until when 2022-2027
Short description (max 100 words): Sampling Scheme Identifier: Diad_ESB_Eel Commercial and recreational fisheries are closed in Ireland, so all data collection is Fishery Independent and Scientific Programme of data collection for eel, glass (recruitment), and silver eel stages. a/ Data collected on silver eel conservation trap and Transport on the Erne (IE_NorW), Shannon (IE_Sha) and Liffey (IE_East). Eels are captured in the programme using location specific gear types, such as bridge mounted coghill nets, and or river anchored V-Wing Fykes. Data collected as part of the ESB Silver Eel Trap and Transport programme, biomass in kg and biometry, length (cm) and weight (gm). Sex is imputed from the length frequency distributions. Programme to estimate silver eel production/escapement and to monitor downstream trap and transport of migrating silver eel using mark-recapture, DIDSON, hydrological profiles and assessment models (described in the Irish TEGE annual assessment reports https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/TEGE_Report_2019_final.pdf , and the Reports to the EU – eg https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/ROI%20Eel%20Management%20Actions%202015-2017_6%20July%202018.pdf) Data collection is used for reporting under obligation of the EU Eel Stock Recovery Regulation, and the ICES Data calls for eel stock assessment in ICES WGEEL. b/ Data collected on recruiting glass eel and young yellow eels. Fixed permanent elver ladder traps monitor upstream recruiting juvenile eel. These are situated at or closed to the tidal limit and are fixed to natural or hydropower barriers or weirs on the Shannon Ardnacrusha and Parteen (IE_Sha) and Erne (IE_NorW). These traps provide a valuable time series of relative abundance of glass eel and young yellow eel

recruits and are used in the annual ICES WGEEL stock assessment. Data type is census data and measured in kgs.
Description of the population
Population targeted: Target Species: <i>Anguilla anguilla</i> National, using index rivers and reporting by Eel Management Unit (EMU) The silver eel data are collected from the total emigrating stock for three individual rivers, Erne, Shannon and Lee, which cover approx. 50% of the Irish wetted area. Population sampled: Specify which part of the target population will be sampled and specify which part of the target population is unreachable for sampling or excluded for some reason to explain, <i>e.g. major ports being listed as auctions excluding all minor ports and no sampling during the week-ends</i>. For research surveys at sea describe target species in single-species surveys or ecosystem component (<i>e.g. demersal, pelagic</i>) in multispecies surveys. Recruits (glass eel and young yellow eel): census data at whole river impassable barriers Silver Eel production (B_{best}) and escapement ($B_{current}$) and mortality ΣA determined on a whole river basis Stratification: EU Regulation (1100/2007) requires reporting by EMU. These rivers (Erne IE_NorW, Shannon IE_Sha, Lee IE_SouW) were identified in the National Eel Management Plan as having major hydropower stations at the tidal limit. They are also key index rivers in the Irish modelling for estimating silver eel output. https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/Ireland%20NationalManagementPlan191208v%5B1%5D.pdf
Sampling design and protocols
Sampling design description: Describe how the sampling allocation is defined; how PSU and SSU are selected for sampling; indicate for which catch fraction the sampling scheme

applies.

Recruitment data are continual census data

Silver Eel data are collected from the Trap and Transport Fisheries. Total Catches.

Is the sampling design compliant with the 4S principle?: Y/N/NA (NA for e.g. surveys and diadromous and recreational sampling schemes) NA

Regional coordination: Indicate if the sampling design and protocols were developed as part of a regional or multi-lateral agreement, and if yes, refer to the agreement (table 1.3) and list all MS participating.

No regional co-ordination. Sampling has been reported since 2009 for silver eel, and since the 1980s for the recruitment data. No issues have been raised in ICES WGEel or in WKEPEMP 2013 or WKEMP 2018

https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2013/WKEPEMP/wkepemp_2013.pdf

https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2018/WKEMP/wkemp_2018.pdf

Link to sampling design documentation:

The overall framework for the silver eel trap and transport programme was laid out in the Irish Eel Management Plan (EMP) (https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/Ireland%20NationalManagementPlan191208v%5B1%5D.pdf). It is managed by the Irish Electricity Supply Board (ESB) and its use as index sites in the silver eel stock assessments is also described in the Irish EMP.

The Recruitment monitoring elver traps are located at historically chosen sites laid down in legislation governing the construction of the hydropower stations.

Compliance with international recommendations:

There are no specific international recommendations.

This programme follows any recommendations as laid out in the EU Regulation for the Recovery of the Eel Stock (EU 1100/2007).

Also follows recommendations from ICES WGEEL

(<https://www.ices.dk/community/groups/Pages/WGEEL.aspx>) and ICES Eel Datacalls (<https://www.ices.dk/sites/pub/Publication%20Reports/Data%20calls/data.call.WGEEL.2021.zip>)

Link to sampling protocol documentation:

Refer to the Irish Eel Management Plan and subsequent compliance reports in 2012, 2015, 2018 and 2021 (not published yet) for details (https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/Ireland%20NationalManagementPlan191208v%5B1%5D.pdf https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/ROI%20Eel%20Management%20Actions%202015-2017_6%20July%202018.pdf)

Compliance with international recommendations: Member State shall state ‘Y’ (yes) if the sampling protocol is in line with international recommendations, and ‘N’ if not. If no relevant expert or coordination groups exist, the sampling protocol should be shortly explained in the text, and should be available upon request for the evaluators.

Y

Refer to the Irish Eel Management Plan and subsequent compliance reports in 2012, 2015, 2018 and 2021 for details (https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/Ireland%20NationalManagementPlan191208v%5B1%5D.pdf and https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/ROI%20Eel%20Management%20Actions%202015-2017_6%20July%202018.pdf).

Also compliance with ICES datacalls. (<https://www.ices.dk/sites/pub/Publication%20Reports/Data%20calls/data.call.WGEEL.2021.zip>)

Sampling implementation

Recording of refusal rate:

NA

Monitoring of sampling progress within the sampling year: Indicate how sampling allocations are adjusted (if needed) and followed-up, what are the mechanisms in place to resolve issues and adopt mitigation measures during the sampling year?

Oversight by the National Technical Expert Group on Eel.

The total catches are what is caught – there is no sub-sampling – its census data or total catch data.

Sampling is monitored at end of season and adjusted for next season

Biological sampling rates can be adjusted as well

Data capture

Means of data capture: short description (+ photo optionally). Indicate what are the

means for collecting the data, e.g. scales, measuring board, dedicated software, ...

Field data are initially recorded by hand on field sheets or fisher's logs, and subsequently transcribed to excel spreadsheets. Data are uploaded annually to the ICES WGEEL Database.

Recruitment

Data collected from permanent ramp type elver traps, lengths using measuring board.

Silver Eel

ESB rivers (Shannon, Erne, Lee), capture by coghill and v-wing fyke nets, biometry by measuring board and balance

Otoliths are prepared by burning and cracking, following ICES WKAREA protocols, and images are captured using ImageProPlus software. Growth data is calculated by the software and transferred to excel as described in WKAREA1. https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Eel%20otolith%20ageing%20SOP.doc

Data capture documentation:

See The Stock Annexes attached to the Irish Reports to the EU in 2012, 2015, and 2018 for descriptions of the Trap and Transport programmes, estimation of silver eel escapement etc. https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/ROI%20Eel%20Management%20Actions%202015-2017_6%20July%202018.pdf

Methods are also described in:

MacNamara, R. & McCarthy, T.K. (2013). Silver eel (*Anguilla anguilla*) population dynamics and production in the River Shannon, Ireland, *Ecology of Freshwater Fish* **23** (2), 181-192.

McCarthy, T.K., Nowak, D., Grennan, J., Bateman, A., Conneely, B. & MacNamara, R. (2014). Spawner escapement of European eel (*Anguilla anguilla*) from the River Erne, Ireland. *Ecology of Freshwater Fish* **23** (1), 21-32.


ICES Manual for Eel age determination WKAREA 1 and associated Manual, WKARE 2 and associated Manual (https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Eel%20otolith%20ageing%20SOP.doc)

Quality checks documentation: Indicate with 'Y' (yes) or 'N' (no). If 'N' (no), indicate when (year) documentation will be available. Provide a link to a webpage where the documentation can be found. If no link is available, but documentation exists, provide a literature reference (author(s), year and type of publication - e.g. internal report). If no documentation on the quality checks exists, provide some details in the text box.

Visual checks are applied to the data at each stage. Once data is on spreadsheet, graphical

checks are used to identify outliers. These are either changed with cross checks with the paper records and data providers, or deleted where verification is not possible.

Below is an example of erroneous data in a graphical plot. Point 1 was weighed as 100gm but typed in as 1000gm and Point 2 was weighed as 430gm but typed in as 43.0gm.



Model outputs and stock assessment estimates are also cross-checked for spurious data.

Fishers logs and reported catch data are followed up if data is not consistent with expectations.

Data storage

National database:

Recruitment data are uploaded annually to the ICES Eel Database.

Index silver eel data and biomass estimates are uploaded into the WGEEL database every 3 years, in compliance with ICES data calls.

International database: Provide the name of international database(s) and the organisation hosting the database, if applicable. Otherwise, insert 'NA' (not applicable). Provide a link if the database is accessible through a website.

ICES WGEEL Eel Database

Quality checks and data validation documentation:

Eel Stock Annex provides some information

https://www.ices.dk/sites/pub/Publication%20Reports/Stock%20Annexes/2020/Anguilla_a_nguilla_SA.pdf

A git repository hosts the code for WGEEL for recruitment analysis and data processing and to facilitate scientific collaboration:

https://github.com/ices-eg/wg_WGEEL

"Git" is a version control system that manages and stores revisions of projects. GitHub is a Git repository hosting service. It provides a Web-based graphical interface, access control and several collaboration features, such as a wikis and basic task management tools for every project.

The relevant data (landings, restocking, mortality rates, biomass indicators) provided through the Data Call are integrated into the existing WGEEL database using a shiny application. The idea is (1) to let WGEEL experts carry out checks on the new files, (2) help national correspondents to qualify their data for quality (3) compare the new data with the existing data in the database and check for duplicates. There are two applications, one to edit data straight into the database, and display graphs to check for duplicates once data are submitted. Detailed information can be found on the website:

https://github.com/ices-eg/wg_WGEEL/tree/master/R/shiny_data_integration

The second shiny application is used to visualize and analyse the data provided. It can be found at:

http://185.135.126.249:8080/shiny_dv/

Sample storage

Storage description:

Otoliths are stored dry in envelopes in the MI or AFBINI Archives. Prepared otoliths (burn and crack) are stored mounted in cured silicone in the MI Archive. These archive databases are currently being updated (2021)

Data processing

Evaluation of data accuracy (bias and precision):

No formal evaluation, data used by WGEEL for years (since early 1980s. No issues have been identified.

See previous information on the WGEEL GITHUB.

Editing and imputation methods:

Missing weights in length weight data are calculated using log length/weight regressions, for the purposes of calculating biomass.

Sex ratios of silver eel are determined from length frequencies using the Bhattacharya Method (1967), validated using sub-samples of dissected eel

Bhattacharya, C. G. 1967. A simple method of resolution of a distribution into Gaussian components. Biometrics, 23: 115–135.

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R
1	L1	N1+	ln N1+	ln N1+	L (0)	ln N1	ln N1	N1	N2+									
1	L1	N1+	ln N1+	ln N1+	L (0)	ln N1	ln N1	N1	N2+									
2	26	0			26			0										
3	27	0			27			0										
4	28	1	0.893	0.893	28	1.715		2	0									
5	29	2	0.693	0.693	29	1.104		9	0									
6	30	3	0.197	0.197	30	0.524		9	0									
7	31	20	2.995	2.995	31	1.333		20	0									
8	32	67	4.205	4.205	32	1.142		67	0									
9	33	99	4.595	4.595	33	0.951		99	0									
10	34	80	4.382	-0.213	34	0.760		80	0									
11	35	142	4.855	0.574	35	0.555		142	0									
12	36	115	4.745	-0.211	36	0.373	4.745	115.01	0									
13	37	120	4.787	0.043	37	0.187	4.852	138.70	0									
14	38	118	4.771	-0.017	38	-0.004	4.929	138.20	0									
15	39	115	4.700	-0.070	39	-0.164	4.734	113.77	0									
16	40	74	4.304	-0.395	40	-0.385	4.349	77.39	0									
17	41	71	4.253	-0.041	41	-0.075	3.773	43.48	27.51									
18	42	69	4.234	-0.029	42	-0.767	3.005	20.19	48.81									
19	43	59	4.078	-0.157	43	-0.858	2.047	7.75	51.25									
20	44	56	4.190	0.112	44	-1.149	0.898	2.45	63.54									
21	45	50	4.382	0.182	45	-1.340	-0.442	0.54	79.35									
22	46	116	4.754	0.372	46	-1.531	-1.972	0.14	115.88									
23	47	101	4.608	0.445	47	-1.722	-0.684	0.02	180.88									
24	48	211	5.352	0.153	48	-1.913	5.697	0.50	211.00									
25	49	217	5.380	0.028	49	-2.104	-7.710	0.00	217.00									
26	50	225	5.421	0.041	50	-2.294	-10.005	0.00	225									
27	51	194	5.269	-0.153	51				194									
28	52	162	5.088	-0.189	52				162									
29	53	129	4.895	-0.228	53				129									
30	54	78	4.357	-0.503	54				78									
31	55	66	4.194	-0.262	55				66									
32	56	47	3.850	-0.244	56				47									
33	57	24	3.178	-0.872	57				24									
34	58	14	2.633	-0.539	58				14									
35	59	13	2.555	-0.074	59				13									
36	60	6	1.792	-0.773	60				6									
37	61	3	1.099	-0.893	61				3									
38	62	3	1.099	-0.893	62				3									
39	63	0	-1.099	0.000	63				0									
40	64	1	0.000	0.000	64				1									
41	65	0	0.000	0.000	65				0									
42	66	0	0.000	0.000	66				0									
43	67	1	0.000	0.000	67				1									
44	68	1	0.000	0.000	68				1									

Quality document associated to a dataset:

No. There isn't a DoI publication of the data.

Data are summarised and published in the Annual Technical Expert Group on Eel Reports (https://www.dcmaph-ireland.ie/sites/default/files/DCF_Files/TEGE_Report_2019_final.pdf) and in the triennial Stock Annexes reported to the EU (https://www.dcmaph-ireland.ie/sites/default/files/DCF_Files/ROI%20Eel%20Management%20Actions%202015-2017_6%20July%202018.pdf)

Validation of the final dataset:

Data are quality checked during capture and transcription, the annual dataset is then manually / graphically checked for correctness and consistency, the data undergoes further checks upon upload to the WGEel database. Spurious unverified data is discarded and not used in any stock assessments.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No Deviations

Sampling Design and Protocols

No Deviations

Sampling Implementation

Eel sampling in Ireland under the EUMAP involves several actors one of whom (ESB) is a national utilities company, who fund and manage this activity directly. This company changed some of its sampling arrangements in 2023, the impact of this is being discussed at the National Technical Eel Group who are working towards ensuring the Irish sampling for eel is mitigated.

Data Capture

No Deviations

Data Storage

Stored on excel and uploaded to the ICES WGEEL database

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: Diad_MI_Eel (*Anguilla anguilla*)

MS : Ireland

Region: North East Atlantic

Sampling scheme identifier: Diad_MI_Eel (*Anguilla anguilla*)

Sampling scheme type: Diadromous (Scientific)

Observation type: SciObs water body

Time period of validity: from when until when 2022-2027

Short description (max 100 words):

Species: *Anguilla Anguilla*

Sampling Scheme Type: Diadromous (Scientific)

Sampling Scheme Identifier: Diad_ESB_Eel

Commercial and recreational fisheries are closed in Ireland, so all data collection is Fishery Independent and Scientific

Programme of data collection for eel, glass (recruitment), yellow (standing stock) and silver eel stages.

a/ Data collected on silver eel: Permanent traps monitor downstream migrating silver eels on the Burrishoole River (IE_West) providing a full daily census for estimating annual production and escapement of silver eel. Numbers of fish migrating downstream, daily number, size, weight and sex ratio of emigrating silver eels (used in the Irish model for estimating silver eel escapement - IMESE), described in the Irish TEGE annual assessment reports, and the Reports to the EU – eg https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/TEGE_Report_2019_final.pdf

<https://www.dcmapp->

ireland.ie/sites/default/files/DCF_Files/ROI%20Eel%20Management%20Actions%202015-2017_6%20July%202018.pdf)

Data collection is used for reporting under obligation of the EU Eel Stock Recovery Regulation, and the ICES Data calls for eel stock assessment in ICES WGEEL.

b/ Data collected on recruiting glass eel and young yellow eels.

Fixed permanent elver ladder traps monitor upstream recruiting juvenile eel. These are situated at or closed to the tidal limit and are fixed to natural or hydropower barriers or weirs on the Liffey (IE_East) and Burrishoole (IE_West).

Elver traps provide annual index recruitment abundance data for ICES WGEEL. Data are collected in biomass (kg) or where numbers are very low, counts are made and converted to biomass. For WGEEL datacalls, numbers are converted from biomass using site specific conversion factors related to size and age of recruits.

c/ Data collected on Yellow Eel Standing Stock; Electrofishing and fyke net surveys

Electrofishing river surveys and fyke net lake surveys on the Burrishoole Catchment (IE_West) target yellow eel in selected water bodies, all fish are identified; weight and length measurements taken. Used for estimating yellow eel (river, lake and transitional water) populations. Used in time series analysis in ICES WGEEL. Data may be used in future eel stock assessment modelling employing the French Eel Density Assessment (EDA) model. Outputs will be reported under EU Regulation metrics and also in ICES Datacalls

Above described programmes contribute to the national eel monitoring programme (Eel: Council Regulation 1100/2007), which operates across different Irish agencies and parent departments

Burrishoole river is an index river for national assessment but also for the joint EIFAAC/ICES/GFCM Working Group on Eel (WGEel).

Description of the population

Population targeted:

Target Species: *Anguilla anguilla*

National, using index rivers and reporting by Eel Management Unit (EMU)

The silver eel data are collected from the total emigrating stock for the Burrishoole Index River (IE_West)

Recruit data are collected at Burrishoole (IE_West) and the Liffey (IE_East).

Yellow eel data collected in the Burrishoole in three habitats, rivers, freshwater lakes, and transitional waters (tidal lagoon lake)

Population sampled:

Recruits (glass eel and young yellow eel) : census data at whole river tidal limit

Silver Eel production (Bbest) and escapement (Bcurrent) and mortality ΣA determined on a whole river basis

Yellow eel relative abundance using time series of CPUE, length and weight

Stratification:

EU Regulation (1100/2007) requires reporting by EMU.

Burrishoole is a long-term index river and a key river in the Irish modelling for estimating silver eel output under the EU Eel Regulation EMP. https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/Ireland%20NationalManagementPlan191208v%5B1%5D.pdf

Relevant References:

Bornarel, V, et al. inc R Poole (2018). Modelling the recruitment of European eel (*Anguilla anguilla*) throughout its European range. ICES Journal of Marine Science 75(2), 541-552. <https://doi.org/10.1093/icesjms/fsx180>

Poole, W.R., Reynolds, J.D.R. & Moriarty, C. (1990). Observations on the silver eel migrations of the Burrishoole river system, Ireland. 1959 to 1988. *Int. Revue Ges Hydrobiol.* **75** (6); 807-815.

Poole W. R., Diserud, O.H., Thorstad, E.B., Durif, C., Dolan, C., Sandlund, O.T., Bergesen, K., Rogan, G., Kelly, S. & Vollestad, L.A. (2018). Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science*, 75 (5); 1627-1637; doi:10.1093/icesjms/fsy053

Sampling design and protocols

Sampling design description:

Recruitment data are continual census data at fixed trap locations

Silver Eel data are collected from the permanent traps in Burrishoole, a daily census

Yellow eel surveys are targeted at the warmer months of the year, with standard locations,

standard gear and fixed effort.

Is the sampling design compliant with the 4S principle?: Y/N/NA (NA for e.g. surveys and diadromous and recreational sampling schemes) NA

Regional coordination:

No regional co-ordination. Sampling has been reported since 2009 for silver eel, and since the 1980s for the recruitment data. No issues have been raised in ICES WGEel or in WKEPEMP 2013 or WKEMP 2018

https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2013/WKEPEMP/wkepemp_2013.pdf

https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2018/WKEMP/wkemp_2018.pdf

Link to sampling design documentation:

The overall framework for the silver eel census programme was laid out in the Irish Eel Management Plan (EMP) (https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/Ireland%20NationalManagementPlan191208v%5B1%5D.pdf), making use of historical locations with time series of data.

Burrishoole Silver eel trapping is described in Poole et al. (1990, 1998) and in the Burrishoole Traps Manual (https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Freshwater%20Traps%20Operation%20Document%2005-10-2015.pdf)

Compliance with international recommendations:

There are no specific international recommendations.

This programme follows any recommendations as laid out in the EU Regulation for the Recovery of the Eel Stock (EU 1100/2007).

Also follows recommendations from ICES WGEEL and ICES Eel Datacalls (<https://www.ices.dk/community/groups/Pages/WGEEL.aspx> <https://www.ices.dk/sites/pub/Publication%20Reports/Data%20calls/data.call.WGEEL.2021.zip>)

Link to sampling protocol documentation:

Refer to the Irish Eel Management Plan and subsequent compliance reports in 2012, 2015, 2018 and 2021 (not published yet) for details (https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/Ireland%20NationalManagementPlan191208v%5B1%5D.pdf https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/ROI%20Eel%20Management%20Actions%202015

-2017_6%20July%202018.pdf) Also refer to the Burrishoole Traps Manual, which is currently being updated (2021) (https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Freshwater%20Traps%20Operation%20Document%2005-10-2015.pdf) Compliance with international recommendations: Y Refer to the Irish Eel Management Plan and subsequent compliance reports in 2012, 2015, 2018 and 2021 for details: https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/Ireland%20NationalManagementPlan191208v%5B1%5D.pdf https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/ROI%20Eel%20Management%20Actions%202015-2017_6%20July%202018.pdf Also compliance with ICES datacalls. https://www.ices.dk/sites/pub/Publication%20Reports/Data%20calls/data.call.WGEEL.2021.zip
Sampling implementation
Recording of refusal rate: NA Monitoring of sampling progress within the sampling year: Oversight by the National Technical Expert Group on Eel. The total catches are what is caught – there is no sub-sampling – its census data or total catch data. Sampling is monitored at end of season and adjusted for next season. Biological sampling rates can be adjusted as well
Data capture
Means of data capture: Field data are initially recorded by hand on field sheets (waterproof paper), and subsequently transcribed to excel spreadsheets. Data are uploaded monthly to the Burrishoole SQL traps database, and annually to the ICES WGEEL Database. Recruitment Data collected from permanent ramp type elver traps, lengths using measuring board. Yellow Eel Data are collected as follows: Rivers using Backpack electrofishers, lakes using standard

fyke nets fished in chains of 10.

Silver Eel

Silver eels captured in permanent downstream fish traps (see https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Freshwater%20Traps%20Operation%20Document%2005-10-2015.pdf)

Otoliths are prepared by burning and cracking, following ICES WKAREA protocols, and images are captured using ImageProPlus software. Growth data is calculated by the software and transferred to excel as described in WKAREA1.

https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Eel%20otolith%20ageing%20SOP.doc

Data capture documentation:

Silver eels

(See https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Freshwater%20Traps%20Operation%20Document%2005-10-2015.pdf)


Yellow Eels (See SOP for field data); Providing a new SOP for Eel fyke netting in 2022.

Recruitment – Needs a new SOP document (2022)

Quality checks documentation:

Visual checks are applied to the data at each stage. Once data is on spreadsheet, graphical checks are used to identify outliers. These are either changed with cross checks with the paper records and data providers, or deleted where verification is not possible.

Below is an example of erroneous data in a graphical plot. Point 1 was weighed as 100gm but typed in as 1000gm and Point 2 was weighed as 430gm but typed in as 43.0gm.



Model outputs and stock assessment estimates are also cross-checked for spurious data.

Data storage

National database:

Recruitment data are uploaded annually to the ICES Eel Database. Stored locally on spreadsheets and on daily backups.

Yellow eel data are uploaded annually to the ICES Eel Database. Stored locally on spreadsheets and on daily backups.

Index silver eel data and biomass estimates are uploaded into the WGEEL database every 3 years, in compliance with ICES data calls. Silver eel data stored locally on spreadsheet and uploaded monthly to the Burrishoole Traps Database which also stores the upload sheets and QC sheets.

<http://data.marine.ie/geonetwork/srv/eng/catalog.search#/metadata/ie.marine.data:dataset.4343>

International database:

ICES WGEEL Eel Database

Quality checks and data validation documentation:

Eel Stock Annex provides some information

https://www.ices.dk/sites/pub/Publication%20Reports/Stock%20Annexes/2020/Anguilla_anguilla_SA.pdf

A git repository hosts the code for WGEEL for recruitment analysis and data processing and to facilitate scientific collaboration:

https://github.com/ices-eg/wg_WGEEL

"Git" is a version control system that manages and stores revisions of projects. GitHub is a Git repository hosting service. It provides a Web-based graphical interface, access control and several collaboration features, such as a wikis and basic task management tools for every project.


The relevant data (landings, restocking, mortality rates, biomass indicators) provided through the Data Call are integrated into the existing WGEEL database using a shiny application. The idea is (1) to let WGEEL experts carry out checks on the new files, (2) help national correspondents to qualify their data for quality (3) compare the new data with the existing data in the database and check for duplicates. There are two applications, one to edit data straight into the database, and display graphs to check for duplicates once data

are submitted. Detailed information can be found on the website: https://github.com/ices-eg/wg_WGEEL/tree/master/R/shiny_data_integration The second shiny application is used to visualize and analyse the data provided. It can be found at: http://185.135.126.249:8080/shiny_dv/
Sample storage
Storage description: Otoliths are stored dry in envelopes in the MI or AFBINI Archives. Prepared otoliths (burn and crack) are stored mounted in cured silicone in the MI Archive. These archive databases are currently being updated (2021).
Data processing
Evaluation of data accuracy (bias and precision): No formal evaluation, data used by WGEEL for years (since early 1980s. No issues have been identified. See previous information on the WGEEL GITHUB. Silver Eel data has been peer-reviewed: Poole, W.R., Reynolds, J.D.R. & Moriarty, C. (1990). Observations on the silver eel migrations of the Burrishoole river system, Ireland. 1959 to 1988. <i>Int. Revue Ges Hydrobiol.</i> 75 (6); 807-815. Poole W. R., Diserud, O.H., Thorstad, E.B., Durif, C., Dolan, C., Sandlund, O.T., Bergesen, K., Rogan, G., Kelly, S. & Vollestad, L.A. (2018). Long-term variation in numbers and biomass of silver eels being produced in two European river systems. <i>ICES Journal of Marine Science</i>, 75 (5); 1627-1637; doi:10.1093/icesjms/fsy053 Sandlund, O.T., Diserud, O. H., Poole, R., Bergesen, K., Dillane, M., Rogan, G., Durif, C., Thorstad, E. B., and Vøllestad, L. A. 2017. Timing and pattern of annual silver eel migration in two European watersheds are determined by similar cues. <i>Ecology and Evolution</i>, DOI:10.1002/ece3.3099; 11pp. Yellow Eel data is being prepared for publication in early 2022. Editing and imputation methods: Y

Missing silver eel and yellow eel weights in length weight data are calculated using log length/weight regressions, for the purposes of calculating total biomass.

Sex ratios of silver eel are determined from length frequencies using the Bhattacharya Method (1967), validated using sub-samples of dissected eel

Bhattacharya, C. G. 1967. A simple method of resolution of a distribution into Gaussian components. *Biometrics*, 23: 115–135.



Quality document associated to a dataset:

No. There isn't a DoI publication of all the data.

Silver eel count data are:

<http://data.marine.ie/geonetwork/srv/eng/catalog.search#/metadata/ie.marine.data:dataset.4343>

Data are summarised and published in the Annual Technical Expert Group on Eel Reports (https://www.dcmaph-ireland.ie/sites/default/files/DCF_Files/TEGE_Report_2019_final.pdf) and in the triennial Stock Annexes reported to the EU (https://www.dcmaph-ireland.ie/sites/default/files/DCF_Files/ROI%20Eel%20Management%20Actions%202015-2017_6%20July%202018.pdf) and in the previous peer-review publications

Validation of the final dataset:

Data are quality checked during capture and transcription, the annual dataset is then manually / graphically checked for correctness and consistency, the data undergoes further checks upon upload to the WGEel database. Spurious unverified data is discarded and not used in any stock assessments.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

Evidence of stock decline and decrease in eel size from both yellow eel surveys and silver

eel monitoring.

Sampling Design and Protocols

No Deviations

Sampling Implementation

No Deviations

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: Diad_MI_Salmon_CWT (*Salmo salar*)

MS : Ireland

Region: North East Atlantic

Sampling scheme identifier: Diad_MI_Salmon_CWT(*Salmo salar*)

Sampling scheme type: Diadromous (recreational)

Observation type: SciObs water body

Time period of validity: from when until when 2022-2027

Short description (max 100 words):

Species: *Salmo salar* / *Salmo trutta*

Sampling Scheme Type: Diadromous (Recreational)

Sampling Scheme Identifier: Diad_MI_Salmon_CWT

National Coded Wire Tagging programme data collection for salmon. Tag seaward migrating salmon smolts, detected upon river return as adults. Data include release and recovery locations, length of tagged smolt, dates and sea age.

Tagging carried out on 7 rivers, BUNDORRAGHA RIVER; BURRISHOOLE RIVER; CONG RIVER; CORRIB RIVER; ERNE RIVER; LEE RIVER and SHANNON RIVER Data used in estimation of survival/exploitation rates and straying of wild/hatchery salmon. Data are provided in data calls to ICES WGNAS. Index data from Burrishoole are used to calibrate annual models and assessments, due to the long time series available. Data also used by the National Technical Expert group on Salmon to calibrate the national assessments for setting Conservation Limits.
Description of the population Population targeted: Target Species: <i>Salmo salar</i> Stocks targeted are the ranched Salmon smolts (downstream migrating recruits) in the BUNDORRAGHA RIVER; BURRISHOOLE RIVER; CONG RIVER; CORRIB RIVER; ERNE RIVER; LEE RIVER and SHANNON RIVER Population sampled: Returning coded wire tagged adults at traps on the Bundorragha river, Burrishoole river, Cong River, Corrib river, Erne river, Lee river and Shannon river. Stratification: t's a total stock census for salmon on the Bundorragha river, Burrishoole river, Cong River, Corrib river, Erne river, Lee river and Shannon river.
Sampling design and protocols Sampling design description: Returning adults sampled at traps on rivers. Sample (core) taken if adult presents CWT (metal detection). Sample is subsequently sent to Marine Institute Newport where sample is analysed and entered into National Coded Wire Tagging database. Is the sampling design compliant with the 4S principle?: Y/N/NA (NA for e.g. surveys and diadromous and recreational sampling schemes) NA Regional coordination: No regional co-ordination.

Data are provided in annual datacall to ICES WGNAS. Marine survival data are used to calibrate annual models and assessments, for WGNAS.

Data also used by the National Technical Expert group on Salmon to calibrate the national assessments for setting Conservation Limits.

Link to sampling design documentation:

<https://www.dcmmap-ireland.ie/documents/methodologies>

<https://data.gov.ie/dataset/national-coded-wire-tagging-and-tag-recovery-programme>

Compliance with international recommendations:

There are no specific international recommendations.

Link to sampling protocol documentation:

<https://www.dcmmap-ireland.ie/documents/methodologies>

https://www.dcmmap-ireland.ie/sites/default/files/DCF_Files/docs/National%20Microtag%20Recovery%20Programme%20Student%20Protocol.docx

Compliance with international recommendations:

N, there is currently none from WGNAS.

Sampling implementation

Recording of refusal rate:

NA

Monitoring of sampling progress within the sampling year:

Oversight by the National Technical Expert Group on Salmon

The total catches are what is caught – there is no sub-sampling – its census data or total catch data.

Sampling is monitored at end of season and adjusted for next season

Data capture

Means of data capture:

Returning adults sampled at traps on rivers.

Field data are initially recorded by hand on field sheets (waterproof paper), and subsequently transcribed to excel spreadsheets.

Sample (core) taken if adult presents CWT (metal detection). Sample is subsequently sent to Marine Institute Newport where sample is analysed and serial number on microtag entered into National Coded Wire Tagging database. Data capture documentation: https://www.dcmapp-ireland.ie/documents/methodologies Quality checks documentation: Visual checks are applied to the data at each stage. Once data is on spreadsheet, graphical checks are used to identify outliers. These are either changed with cross checks with the paper records and data providers, or deleted where verification is not possible.
Data storage
National database: Microtag data are stored on the National Coded Wire tagging database. International database: ICES WGNAS salmon database Quality checks and data validation documentation: Not currently available.
Sample storage
Storage description: Cores are stored in tubes filled in ethanol in the MI, until they are prepared for analysis.
Data processing
Evaluation of data accuracy (bias and precision): No formal evaluation, data used by WGNAS for years (since early 1980s. No issues have been identified. Editing and imputation methods: N Quality document associated to a dataset: No. There isn't a DoI publication of all the data.

Data are summarised and published in the Annual National Coded Tag recovery report
<https://data.gov.ie/dataset/national-coded-wire-tagging-and-tag-recovery-programme>

Validation of the final dataset:

Data are quality checked during capture and transcription, the annual dataset is then manually / graphically checked for correctness and consistency, the data undergoes further checks upon upload to the international databases. Spurious unverified data is discarded and not used in any stock assessments.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No Deviations

Sampling Design and Protocols

No Deviations

Sampling Implementation

No Deviations

Data Capture

A lot of work done on improving data quality, both current and previous. Work ongoing to develop the MI databases structure to improve recording and quality of data.

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: Diad_MI_Salmon_SeaTrout (Salmo salar & Salmo trutta)

MS : Ireland

Region: North East Atlantic

Sampling scheme identifier: Diad_MI_Salmon_SeaTrout (*Salmo salar* & *Salmo trutta*)

Sampling scheme type: Diadromous (Scientific)
Observation type: SciObs water body
Time period of validity: from when until when 2022-2027
Short description (max 100 words): Species: <i>Salmo salar</i> / <i>Salmo trutta</i> Sampling Scheme Type: Diadromous (Scientific) Sampling Scheme Identifier: Diad_MI_Salmon_SeaTrout Programme of data collection for salmon and sea trout in the Burrishoole Index Catchment, West of Ireland. This annex covers wild salmon and wild sea trout parr, downstream migrating smolt and upstream adult returns. Also collected are data on the returns of tagged ranched salmon, numbers, survival and growth/sex. Data for the total fish census, including counts, survivals (freshwater/marine), size and fecundity, sex, have been collected in Burrishoole since 1970. Primary data are from total river permanent fish traps, with supplementary data on parr from river electrofishing surveys.
Description of the population
Population targeted: Target Species: <i>Salmo salar</i> & <i>Salmo trutta</i> Stocks targeted are the total stock (downstream migrating recruits and upstream migrating returning adults) for both species in the Burrishoole river. Population sampled: Recruits (parr and smolt): census data at whole river tidal limit, parr are site specific surveys, smolts are total production Adult salmon is the total return of both wild and ranched stock - a total census. Biological sampling covers approx. 25% of the returns of wild salmon (to minimise handling) and all the ranched fish. Wild sea trout, a total census of all returning trout (fully anadromous fish and all other non-silvered “slob” trout)

Stratification:

Burrishoole is a long-term index river and a key river in the Irish modelling setting conservation limits for salmon.

It's a total stock census for both salmon and sea trout

Relevant References:

E. de Eyto, J. White, P. Boylan, B. Clarke, D. Cotter, D. Doherty, P. Gargan, R. Kennedy, P. McGinnity, N. O'Maoiléidigh, K. O'Higgins, (2015). The fecundity of wild Irish Atlantic salmon *Salmo salar* L. and its application for stock assessment purposes,

Fisheries Research, 164; 159-169, ISSN 0165-7836

<https://doi.org/10.1016/j.fishres.2014.11.017>.

Poole, W.R., Dillane, M., deEyto, E., Rogan, G., McGinnity, P. & Whelan, K. (2006). Characteristics of the Burrishoole sea trout population: census, marine survival, enhancement and stock recruitment, 1971-2003. In: *Sea Trout: Biology, Conservation and Management* (Harris, G.S. & Milner, N.J., Eds). Proceedings of the First International Sea Trout Symposium, July 2004, Cardiff, Wales, UK. Blackwell Publishing, Oxford, pp. 279-306.

Sampling design and protocols**Sampling design description:**

Recruitment data parr are from annual electrofishing surveys carried out at standard locations.

Recruitment data (smolt) are continual census data at fixed trap locations at the tidal limit in the Burrishoole.

Adult upstream count data are collected from the permanent traps in Burrishoole, a daily census

Link to Trap manual https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Freshwater%20Traps%20Operation%20Document%2005-10-2015.pdf

Is the sampling design compliant with the 4S principle?: Y/N/NA (NA for e.g. surveys and diadromous and recreational sampling schemes) NA

Regional coordination:

No regional co-ordination. Sampling follows the schedule outlined in the ICES WKESDCF workshop report (2012) for index rivers.

Data are provided in datacalls to ICES WGNAS and in annual data collation in ICES

WGTRUTTA. Index data from Burrishoole are used to calibrate annual models and assessments, due to the long time series available.

Data also used by the National Technical Expert group on Salmon to calibrate the national assessments for setting Conservation Limits.

Link to sampling design documentation:

Link to TEGOS 2021

<https://www.fisheriesireland.ie/sites/default/files/migrated/docman/The%20Status%20of%20Irish%20Salmon%20Stocks%20in%202020%20with%20Catch%20Advice%20for%202021.pdf>

Burrishoole census trapping is described in Poole et al. (2006) and in the Burrishoole Traps Manual (https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Freshwater%20Traps%20Operation%20Document%2005-10-2015.pdf)

Poole, W.R., Dillane, M., deEyto, E., Rogan, G., McGinnity, P. & Whelan, K. (2006). Characteristics of the Burrishoole sea trout population: census, marine survival, enhancement and stock recruitment, 1971-2003. In: *Sea Trout: Biology, Conservation and Management* (Harris, G.S. & Milner, N.J., Eds). Proceedings of the First International Sea Trout Symposium, July 2004, Cardiff, Wales, UK. Blackwell Publishing, Oxford, pp. 279-306.

Compliance with international recommendations:

There are no specific international recommendations.

the data is however used by WGNAS as Burrishoole is an index river for the Lifecycle model as well as the older Bayesian Model which preceded it.

Also follows recommendations from ICES WGNAS, WGTRUTTA and ICES Datacalls
WGNAS:

<https://www.ices.dk/sites/pub/Publication%20Reports/Forms/DispForm.aspx?ID=37558>

WGTRUTTA:

<https://www.ices.dk/sites/pub/Publication%20Reports/Forms/DispForm.aspx?ID=36884>

Link to sampling protocol documentation:

Also refer to the Burrishoole Traps Manual, which is currently being updated (2021)
https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Freshwater%20Traps%20Operation%20Document%2005-10-2015.pdf

Compliance with international recommendations: Y Also compliance with ICES datacalls. https://www.ices.dk/sites/pub/Publication%20Reports/Forms/DispForm.aspx?ID=37558
Sampling implementation
Recording of refusal rate: NA Monitoring of sampling progress within the sampling year: Oversight by the National Technical Expert Group on Salmon The total catches are what is caught – there is no sub-sampling – its census data or total catch data. Sampling is monitored at end of season and adjusted for next season Biological sampling rates can be adjusted as well
Data capture
Means of data capture: Field data are initially recorded by hand on field sheets (waterproof paper), and subsequently transcribed to excel spreadsheets. Data are uploaded monthly to the Burrishoole SQL traps database, and annually to the ICES Salmon Database. Recruitment Data on parr collected from electrofishing surveys (3 catch fishing depletions, Numbers, densities), lengths using measuring board. Data on smolt collected from downstream traps(sub-daily counts), lengths using measuring board, sex by dissection. Adults Returning wild and ranched salmon adults and wild trout are counted in the upstream traps, full census (see Traps Manual https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Freshwater%20Traps%20Operation%20Document%2005-10-2015.pdf) Scales are prepared by cleaning in freshwater, and reading under a microfiche, or microscope, and images are captured using ImageProPlus software. Growth data is calculated by the software and transferred to excel as described in the Celtic Sea Trout

Manual. https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/CSTP%20Sea%20Trout%20Scale%20Manual%20Vers%201%2027-01-2011.pdf

SALSEA-Merge (2008) Workshop on Digital Scale Reading Methodology, Trondheim, Norway, 8th to 10th

September 2008. 23pp.

Data capture documentation:


Parr (See SOP for field data https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Field%20Survey%20SOPS%20for%20Juvenile%20Salmonids%20and%20eel.doc)

Smolts and adults in permanent traps – see Traps Manual https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/docs/Freshwater%20Traps%20Operation%20Document%2005-10-2015.pdf

Quality checks documentation:

Visual checks are applied to the data at each stage. Once data is on spreadsheet, graphical checks are used to identify outliers. These are either changed with cross checks with the paper records and data providers, or deleted where verification is not possible.

Below is an example of erroneous data in a graphical plot. Point 1 was weighed as 100gm but typed in as 1000gm and Point 2 was weighed as 430gm but typed in as 43.0gm.



Model outputs and stock assessment estimates are also cross-checked for spurious data.

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Data storage

National database: Electrofishing data are stored on backed up excel sheets – a database is in development. Index smolt and adult trap data are stored locally on spreadsheets and uploaded monthly to the Burrishoole Traps Database which also stores the upload sheets and QC sheets. International database: ICES WGNAS salmon database ICES WGTRUTTA database which is in development Quality checks and data validation documentation: None available.
Sample storage
Storage description: Scales are stored dry in envelopes in the MI. Prepared slides are stored mounted in the MI Archive. These archive databases are currently being updated (2021).
Data processing
Evaluation of data accuracy (bias and precision): No formal evaluation, data used by WGNAS for years (since early 1980s. No issues have been identified. Data have been peer-reviewed: E. de Eyto, J. White, P. Boylan, B. Clarke, D. Cotter, D. Doherty, P. Gargan, R. Kennedy, P. McGinnity, N. O'Maoiléidigh, K. O'Higgins, (2015). The fecundity of wild Irish Atlantic salmon <i>Salmo salar</i> L. and its application for stock assessment purposes, <i>Fisheries Research</i>, 164; 159-169, ISSN 0165-7836 https://doi.org/10.1016/j.fishres.2014.11.017. de Eyto, E., Dalton, C., Dillane, M., Jennings, E., McGinnity, P., O'Dwyer, B., Poole, R., Rogan, G., and Taylor, D. (2016). The response of North Atlantic diadromous fish to multiple stressors, including land use change: a multi-decadal study. <i>Can. J. Fish. Aquat. Sci.</i> 73(12): 1759–1769. doi:10.1139/cjfas-2015-0450. Poole, W.R., Dillane, M., deEyto, E., Rogan, G., McGinnity, P. & Whelan, K. (2006). Characteristics of the Burrishoole sea trout population: census, marine survival, enhancement and stock recruitment, 1971-2003. In: <i>Sea Trout: Biology, Conservation and Management</i> (Harris, G.S. & Milner, N.J., Eds). Proceedings of the First International Sea

Trout Symposium, July 2004, Cardiff, Wales, UK. Blackwell Publishing, Oxford, pp. 279-306.

Editing and imputation methods:

N

Quality document associated to a dataset:

No. There isn't a DoI publication of all the data.

Salmon and sea trout smolt count data are:

<http://data.marine.ie/geonetwork/srv/eng/catalog.search#/metadata/ie.marine.data:dataset.4343>

Data are summarised and published in the Annual Statistic Reports for the Burrishoole (<http://hdl.handle.net/10793/1672>) and in the previous peer-review publications

Validation of the final dataset:

Data are quality checked during capture and transcription, the annual dataset is then manually / graphically checked for correctness and consistency, the data undergoes further checks upon upload to the international databases. Spurious unverified data is discarded and not used in any stock assessments.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No Deviations

Sampling Design and Protocols

No Deviations

Sampling Implementation

No Deviations

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing
No Deviations

Sampling scheme identifier: Wild Salmon and Sea Trout Tagging Scheme Recreational

MS : IRL

Region : North-East Atlantic

Sampling scheme identifier : Wild Salmon and Sea Trout Tagging Scheme Recreational

Sampling scheme type: Diadromous (recreational)

Observation type: Self water body

Time period of validity : 2022 - 2027

Short description (max 100 words): Sampling scheme aiming at collecting annual catch quantities for *Salmo salar* in the freshwater part of their lifecycle as specified in Tables 1 and 4 of the EU MAP Delegated Decision annex and to provide data on fishing effort ;number and weight of all salmon caught separated by fisheries, location, age class with estimates also required for unreported catches; weight of ranched salmon caught; assignment to jurisdiction/region/river of origin of adult salmon; and sea age composition of returning adults; as specified by the RCG ISSG Diadromous Fishes.

Description of the population

Population targeted: *Salmo salar*

Population sampled: All nationwide recreational angling catches of *Salmo salar*.

The 5 Index rivers selected for 2022 are Owenmore, Drowes, Mulkear, Laune, Slaney and Boyne.

This may change in subsequent updated

Stratification: NA

Sampling design and protocols

Sampling design description: Data can be collated from all salmon designated rivers open to

fishing. Those river open for fishing are determined annually. This includes a maximum of 144 rivers in Ireland. Catches (with date and weight information) must be reported by anglers in logbooks under the Wild Salmon and Sea Trout Tagging Scheme and submitted to Inland Fisheries Ireland (IFI) annually. IFI collate, validate and publish this information and provide it to the Technical Expert Group on Salmon (TEGOS) (and the ICES Working Group on North Atlantic Salmon) for scientific stock assessment purposes. Scientific advice is then provided to support the regulation and management of fisheries which are designated under annual fishing Regulations.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: NA

Link to sampling design documentation:

TEGOS reports provide the sampling design documentation <https://www.fisheriesireland.ie/what-we-do/research/technical-expert-group-on-salmon-tegos> , information on the *Wild Salmon and Sea Trout Tagging Scheme* can be found at <https://store.fishinginireland.info/salmon-fishing-regulations/> and information on collection and collation of catch data can be found in the *Wild salmon and Sea Trout Statistics Reports* on the IFI website e.g. for 2019

<https://www.fisheriesireland.ie/sites/default/files/transfereed/docman/Wild%20Salmon%20and%20Sea%20Trout%20Statistics%20Report%202019.pdf>

Compliance with international recommendations: Y

Link to sampling protocol documentation:

Sampling protocol documentation can be found in TEGOS reports <https://www.fisheriesireland.ie/what-we-do/research/technical-expert-group-on-salmon-tegos> , are detailed in the *Wild Salmon and Sea Trout Tagging Scheme* <https://store.fishinginireland.info/salmon-fishing-regulations/> and detailed in the *Wild salmon and Sea Trout Statistics Reports* on the IFI website e.g. for 2019

<https://www.fisheriesireland.ie/sites/default/files/transfereed/docman/Wild%20Salmon%20and%20Sea%20Trout%20Statistics%20Report%202019.pdf>

Sampling implementation
Recording of refusal rate: NA Monitoring of sampling progress within the sampling year: Sampling allocations depend on the level of catches reported and the amount of information gathered is known after the end of the fishing season (30th September) when the information is collated. Reported catches are revised depending on logbook return rates per Fisheries District as per Small (1991) (Exploring data provided by angling for salmonids in the British Isles. In : Catch effort sampling strategies (ed. I.G. Cowx), pp 81-91. Fishing News Books, Oxford). All anglers who do not return logbooks are written to as a means of improving logbook returns and a proportion are taken to court annually and fined for non-return of logbooks.
Data capture
Means of data capture: Data are captured by anglers via logbooks under the <i>Wild Salmon and Sea Trout Tagging Scheme</i> and reported to IFI. The data is entered, stored and validated using the Bradán database maintained by IFI. The data is ultimately transferred to the National Salmon Assessment Database (NSADB) for scientific stock assessment purposes and stored on the IFI SQL server. Data capture documentation: The data is captured in angling logbooks. The overall scheme is managed by Inland Fisheries Ireland (as designated under <i>Section 69</i> of the <i>Inland Fisheries Act 2010</i>). General details on this process can be found in the follow sources: <i>Wild Salmon and Sea Trout Tagging Scheme</i> https://store.fishinginireland.info/salmon-fishing-regulations/ and <i>Wild salmon and Sea Trout Statistics Reports</i> on the IFI website e.g. for 2019 https://www.fisheriesireland.ie/sites/default/files/transfereed/docman/Wild%20Salmon%20and%20Sea%20Trout%20Statistics%20Report%202019.pdf Quality checks documentation: N (year for documentation to be available to be confirmed). Data from logbooks is entered by IFI staff in each River Basin District and collated, reviewed and quality checked by the database manager. The data is ultimately transferred to the National Salmon Assessment Database (NSADB) for scientific stock assessment purposes where final QC checks are made before assessments are undertaken.
Data storage

National database: Bradán database and National Salmon Assessment Database (NSADB) stored on the IFI SQL server. International database: NA. Quality checks and data validation documentation: There is no formal quality checks and data validation documentation available at present.
Sample storage
NA
Data processing
Evaluation of data accuracy (bias and precision): N (year for documentation to be available to be confirmed). Editing and imputation methods: N (year for documentation to be available to be confirmed). The database manager is responsible for coordinating these. Quality document associated to a dataset: No and No. Validation of the final dataset: Datasets are validated by the database manager and subsequently validated by IFI scientists before being used by the end-user. In addition, the reporting output is circulated to relevant IFI staff for review, forwarded to IFI board for approval and further sent to DECC for final review before publication on the IFI website. The data is ultimately transferred to the National Salmon Assessment Database (NSADB) for scientific stock assessment purposes where validation checks are made before assessments are undertaken.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan <u>Description of the Population</u> No Deviations <u>Sampling Design and Protocols</u>

No Deviations
<u>Sampling Implementation</u>
No Deviations
<u>Data Capture</u>
No Deviations
<u>Data Storage</u>
No Deviations
<u>Sample Storage</u>
No Deviations
<u>Data Processing</u>
No Deviations

Sampling scheme identifier: Wild Salmon and Sea Trout Tagging Scheme commercial

MS : IRL
Region : North-East Atlantic
Sampling scheme identifier : Wild Salmon and Sea Trout Tagging Scheme commercial
Sampling scheme type: Diadromous (commercial)
Observation type: Self water body
Time period of validity : 2022 - 2027
Short description (max 100 words): These fisheries are primarily in single river estuaries (only three stocks i.e. Killary, Owenmore estuary and Castlemaine are mixed-stock estuary fisheries). As such this is considered to come under sampling scheme aiming at collecting annual catch quantities for <i>Salmo salar</i> in the freshwater part of their lifecycle as specified in Tables 1 and 4 of the EU MAP Delegated Decision annex and to provide data on fishing effort ;number and weight of all salmon caught separated by fisheries, location, age class with estimates also required for unreported catches; weight of ranched salmon caught; assignment to jurisdiction/region/river of origin of adult salmon; and sea age composition of returning adults; as specified by the RCG ISSG Diadromous Fishes. There are no marine commercial or freshwater commercial fisheries for salmon in Ireland.
Description of the population

Population targeted: *Salmo salar*

Population sampled: All nationwide commercial catches of *Salmo salar* in river estuaries.

Bandon; Barrow and Pollmounty; Belclare; Blackwater, Glenshelane, Finisk; Caragh; Castlemaine; Dawros; Eany; Feale, Galey and Brick; Glenamoy; Gweebarra; Ilen; Inny; Killary; Laune and Cottoners; Lower Lee (Cork); Maine; Moy; Newport; Nore; Owenduff; Owenea and Owentocker; Owenglin; Owenmore estuary; Roughty; Sheen; Sneem; Suir, Clodiagh, Lingaun, Blackwater; Waterville.

Stratification: NA

Sampling design and protocols

Sampling design description: Data can be collated from all salmon designated commercial fisheries open to fishing. Those fisheries open for fishing are determined annually. Catches (with date and weight information) must be reported by commercial fishers in logbooks under the Wild Salmon and Sea Trout Tagging Scheme and submitted to Inland Fisheries Ireland (IFI) annually. IFI collate, validate and publish this information and provide it to the Technical Expert Group on Salmon (TEGOS) (and the ICES Working Group on North Atlantic Salmon) for scientific stock assessment purposes. Scientific advice is then provided to support the regulation and management of fisheries which are designated under annual fishing Regulations.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: NA

Link to sampling design documentation:

TEGOS reports provide the sampling design documentation <https://www.fisheriesireland.ie/what-we-do/research/technical-expert-group-on-salmon-tegos> , information on the *Wild Salmon and Sea Trout Tagging Scheme* can be found at <https://store.fishinginireland.info/salmon-fishing-regulations/> and information on collection and collation of catch data can be found in the *Wild salmon and Sea Trout Statistics Reports*

on the IFI website e.g. for 2019

<https://www.fisheriesireland.ie/sites/default/files/transfereed/docman/Wild%20Salmon%20and%20Sea%20Trout%20Statistics%20Report%202019.pdf>

Compliance with international recommendations: Y

Link to sampling protocol documentation:

Sampling protocol documentation can be found in TEGOS reports <https://www.fisheriesireland.ie/what-we-do/research/technical-expert-group-on-salmon-tegos> , are detailed in the *Wild Salmon and Sea Trout Tagging Scheme* <https://store.fishinginireland.info/salmon-fishing-regulations/> and detailed in the *Wild salmon and Sea Trout Statistics Reports* on the IFI website e.g. for 2019

<https://www.fisheriesireland.ie/sites/default/files/transfereed/docman/Wild%20Salmon%20and%20Sea%20Trout%20Statistics%20Report%202019.pdf>

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: Sampling allocations depend on the level of catches reported and the amount of information gathered is known after the end of the fishing season (30th September) when the information is collated. Return rate of logbooks is 100% as it is strictly regulated by IFI.

Data capture

Means of data capture: Data are captured by fishers via logbooks under the *Wild Salmon and Sea Trout Tagging Scheme* and reported to IFI. The data is entered, stored and validated using the Bradán database maintained by IFI. The data is ultimately transferred to the National Salmon Assessment Database (NSADB) for scientific stock assessment purposes and stored on the IFI SQL server.

Data capture documentation:

The data is captured in logbooks.

The overall scheme is managed by Inland Fisheries Ireland (as designated under *Section 69* of the *Inland Fisheries Act 2010*). General details on this process can be found in the follow sources:

Wild Salmon and Sea Trout Tagging Scheme <https://store.fishinginireland.info/salmon-fishing-regulations/> and *Wild salmon and Sea Trout Statistics Reports* on the IFI website e.g. for 2019

<https://www.fisheriesireland.ie/sites/default/files/transfereed/docman/Wild%20Salmon%20and%20Sea%20Trout%20Statistics%20Report%202019.pdf>

Quality checks documentation: N (year for documentation to be available to be confirmed).

Data from logbooks is entered by IFI staff in each River Basin District and collated, reviewed and quality checked by the database manger. The data is ultimately transferred to the National Salmon Assessment Database (NSADB) for scientific stock assessment purposes where final QC checks are made before assessments are undertaken.

Data storage

National database: Bradán database and National Salmon Assessment Database (NSADB) stored on the IFI SQL server.

International database: NA.

Quality checks and data validation documentation: There is no formal quality checks and data validation documentation available at present. Data is stored in the Bradán database and subsequently the National Salmon Assessment Database (NSADB) hosted by IFI..

Sample storage

NA

Data processing

Evaluation of data accuracy (bias and precision): N (year for documentation to be available to be confirmed).

Editing and imputation methods: N (year for documentation to be available to be confirmed). The database manager is responsible for coordinating these.

Quality document associated to a dataset: No and No.

Validation of the final dataset: Datasets are validated by the database manager and subsequently validated by IFI scientists before being used by the end-user. In addition, the reporting output is circulated to relevant non-scientific IFI staff for review, forwarded to IFI board for approval and further sent to DECC for final review before publication on the IFI website. The data is ultimately transferred to the National Salmon Assessment Database (NSADB) for scientific stock assessment purposes where validation checks are made before assessments are undertaken.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

No Deviations

Sampling Design and Protocols

No Deviations

Sampling Implementation

No sampling was possible where there was no fishery in operation and where the scientific advice was that no sustainable surplus was available for exploitation.

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: Biological sampling NSIC

MS : IRL
Region : North-East Atlantic
Sampling scheme identifier : Biological sampling NSIC
Sampling scheme type: Diadromous (scientific)
Observation type: SciObs water body
Time period of validity : 2022 - 2027
Short description (max 100 words): Length (cm), weight (kg) and age (scale sample as one-sea-winter or multi-sea-winter) are collected from a random sample of 100 adult Atlantic salmon per annum intercepted in the upstream fish trap at the National Salmonid Index Catchment River Erriff.
Description of the population
Population targeted: <i>Salmo salar</i> and <i>Salmo trutta</i> Population sampled: <i>Salmo salar</i> (River Erriff stock) Stratification: NA
Sampling design and protocols
Sampling design description: Data is collected from a random sample of 100 adult Atlantic salmon which represents approximately 5% of the total run annually. Samples are collected from May to September which encompasses the main range of the run. Is the sampling design compliant with the 4S principle?: NA Regional coordination: NA

Link to sampling design documentation:

Millane, M. and Gargan, P. (2021). National Salmonid Index Catchment Erriff Research Plan. Research & Development, Inland Fisheries Ireland pp. 23 (internal document).

Compliance with international recommendations: NA

Link to sampling protocol documentation:

Millane, M. and Gargan, P. (2021). National Salmonid Index Catchment Erriff Research Plan. Research & Development, Inland Fisheries Ireland pp. 23 (internal document).

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: Resident research staff at the National Salmonid Index Catchment monitor sampling progress during the sampling period.

Data capture

Means of data capture: Length (cm) is recorded on a measuring board. Weight is recorded on a manual weighing scales (larger fish) or an electronic balance (smaller fish). Data are initially captured on standard field books and subsequently entered into a standard database. 10-20 scale samples are taken from a fish and stored in scales envelopes for later age determination using a digital scale reader and entered into a standard database.

Data capture documentation:

Millane, M. and Gargan, P. (2021). National Salmonid Index Catchment Erriff Research Plan. Research & Development, Inland Fisheries Ireland pp. 23 (internal document).

Quality checks documentation: Y Millane, M. and Gargan, P. (2021). National Salmonid Index Catchment Erriff Research Plan. Research & Development, Inland Fisheries Ireland pp. 23 (internal document).

Data storage
National database: NA International database: NA. Quality checks and data validation documentation: Millane, M. and Gargan, P. (2021). National Salmonid Index Catchment Erriff Research Plan. Research & Development, Inland Fisheries Ireland pp. 23 (internal document).
Sample storage Scale samples are stored in the NSIC Erriff laboratory and subsequently transferred to the National Salmon Scale Archive (NSSA) in IFI HQ in Dublin for permanent storage. Access to the material is via request to IFI. The archive is hosted by IFI and holds greater than 20,000 scale samples from a total of 38 river systems in Ireland.
Data processing
Evaluation of data accuracy (bias and precision): Y Millane, M. and Gargan, P. (2021). National Salmonid Index Catchment Erriff Research Plan. Research & Development, Inland Fisheries Ireland pp. 23 (internal document). Editing and imputation methods: Y Millane, M. and Gargan, P. (2021). National Salmonid Index Catchment Erriff Research Plan. Research & Development, Inland Fisheries Ireland pp. 23 (internal document). Quality document associated to a dataset: No and No. Validation of the final dataset: 100% of the data entered from field book are reviewed for data entry errors before use by end users.
AR comment: Indicate any deviations. Do not change the text already adopted in the work

plan

Description of the Population

No Deviations

Sampling Design and Protocols

No Deviations

Sampling Implementation

No Deviations

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: Biological sampling smolts NSIC

MS : IRL
Region : North-East Atlantic
Sampling scheme identifier : Biological sampling smolts NSIC
Sampling scheme type: Diadromous (scientific)
Observation type: SciObs water body
Time period of validity : 2022 - 2027
Short description (max 100 words): Number and age composition (scale sample to determine freshwater age) are collected from a random sample of 100 adult Atlantic salmon and 100 Sea trout per annum intercepted in the Tawnyard trap in the National Salmonid Index Catchment River Erriff.
Description of the population
Population targeted: <i>Salmo salar</i> and <i>Salmo trutta</i>

Population sampled: *Salmo salar* & *Salmo trutta* (River Erriff stock)

Stratification: NA

Sampling design and protocols

Sampling design description: Data is collected per annum from a random sample of 100 Atlantic salmon and 100 sea trout in the Erriff. Samples are collected from April to May which encompasses the main range of the run. Samples will be placed in scale envelopes and later read using a light microscope. The number of salmonid smolts in the Erriff intercepted in the Tawnyard trap will be counted per annum.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: NA

Link to sampling design documentation:

Millane, M. and Gargan, P. (2021). National Salmonid Index Catchment Erriff Research Plan. Research & Development, Inland Fisheries Ireland pp. 23 (internal document).

Compliance with international recommendations: NA

Link to sampling protocol documentation:

Millane, M. and Gargan, P. (2021). National Salmonid Index Catchment Erriff Research Plan. Research & Development, Inland Fisheries Ireland pp. 23 (internal document).

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: Research staff at the National Salmonid Index Catchment monitor sampling progress during the sampling period.
Data capture
Means of data capture: Data are initially captured in standard field books and scale envelopes subsequently entered into a standard database. 10-20 scale samples are taken from a fish and stored in scales envelopes for later age determination using a light microscope and entered into a standard database. Data capture documentation: Millane, M. and Gargan, P. (2021). National Salmonid Index Catchment Erriff Research Plan. Research & Development, Inland Fisheries Ireland pp. 23 (internal document). Quality checks documentation: Y Millane, M. and Gargan, P. (2021). National Salmonid Index Catchment Erriff Research Plan. Research & Development, Inland Fisheries Ireland pp. 23 (internal document).
Data storage
National database: NA International database: NA. Quality checks and data validation documentation: Millane, M. and Gargan, P. (2021). National Salmonid Index Catchment Erriff Research Plan. Research & Development, Inland Fisheries Ireland pp. 23 (internal document).
Sample storage
Scale samples are stored in the NSIC Erriff laboratory and subsequently transferred to the National Salmon Scale Archive (NSSA) in IFI HQ in Dublin for permanent storage. Access to the material is via request to IFI. The archive is hosted by IFI and holds greater than 20,000 scale samples from a total of 38 river systems in Ireland.
Data processing

Evaluation of data accuracy (bias and precision): Y

Millane, M. and Gargan, P. (2021). National Salmonid Index Catchment Erriff Research Plan. Research & Development, Inland Fisheries Ireland pp. 23 (internal document).

Editing and imputation methods: Y

Millane, M. and Gargan, P. (2021). National Salmonid Index Catchment Erriff Research Plan. Research & Development, Inland Fisheries Ireland pp. 23 (internal document).

Quality document associated to a dataset: No and No.

Validation of the final dataset: 100% of the data entered from field book and scale envelopes are reviewed for data entry errors before use by end users.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

No Deviations

Sampling Design and Protocols

No Deviations

Sampling Implementation

No Deviations

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: Eel_elver_trap_Diadromous (scientific)

MS : Ireland
Region: North-East Atlantic
Sampling scheme identifier: Eel_elver_trap_Diadromous (scientific)
Sampling scheme type: Diadromous (scientific)
Observation type: SciObs water body
Time period of validity: 2022-2027
Short description (max 100 words): <i>e.g. sampling scheme aiming at collecting length samples from commercial landings on-shore for all species listed in Table 1 of the EU MAP Delegated Decision annex. The scheme covers mainland and all outermost regions ('RUP' in French, Portuguese, and Spanish).</i> The elver traps are located at the high water mark to capture the elvers and yellow eels migrating from the estuary into freshwater. They operate from April to August but can continue to operate into September if catches continue. The traps are fixed and sample in a standardised manner. Additional information include eel length, weight, eye and fin measurements. When necessary samples will be brought back to the laboratory to confirm 0+ status and to get an age profile of the yellow eels migrating upstream.
Description of the population
Population targeted: Specify which are the primary sampling units (PSU), e.g. all national port*days (information present in former Table 4B). For research surveys: specify the main target species from a survey perspective (as opposed to Table 1 in the Annex to the Implementing Decision) and the main survey area. The elver traps target the elver population migrating from estuary into freshwater along one bank of a river. The traps also capture older yellow eels however these are recorded separately from the recruits data.

Population sampled: Specify which part of the target population will be sampled and specify which part of the target population is unreachable for sampling or excluded for some reason to explain, *e.g. major ports being listed as auctions excluding all minor ports and no sampling during the week-ends*. For research surveys at sea describe target species in single-species surveys or ecosystem component (*e.g. demersal, pelagic*) in multispecies surveys.

The elvers that encounter the trap and climb the ramp into the holding box. The traps are fixed in place so sample the same location every year.

Stratification: Explain the logic taken to stratify the population and the number of strata generated, *e.g. population stratified in 3 geographical lots (from A to B, from B to C and from C to D). Each lot is then stratified by auction*.

For reporting to the EU under the Eel Regulation Ireland created Eel Management Units (in line with River Basin Districts). The recommendation from WKESDCF and highlighted within the dcmap directive is for each MS to monitor all eel lifestages within at least one catchment per RBD.

In Ireland the RBD or eel management units are East EMU, SouthEast EMU, SouthWest EMU, Shannon International EMU, West EMU and the NorthWest EMU. In addition there are index catchments with historical information available that is targeted in this programme to continue the longterm temporal trend.

Sampling design and protocols

Sampling design description: Describe how the sampling allocation is defined; how PSU and SSU are selected for sampling; indicate for which catch fraction the sampling scheme applies.

Sites is located at the on one river bank at the high water mark.

Is the sampling design compliant with the 4S principle?: Y/N/NA (NA for *e.g.* surveys and diadromous and recreational sampling schemes)

NA

Regional coordination: Indicate if the sampling design and protocols were developed as part of a regional or multi-lateral agreement, and if yes, refer to the agreement (table 1.3) and list all MS participating.

No regional coordination is undertaken for sampling eel. However for the transboundary North West EMU consultation occurs between agencies in Northern Ireland and Republic of Ireland in relation to combined eel surveys on occasion.

Link to sampling design documentation: Provide a link to a webpage where the

documentation can be found. If no link is available, but documentation exists, Member State shall provide a literature reference (author(s), year and type of publication - e.g. internal report). If no documentation on the sampling design exists, Member State shall provide some details in the textbox.

An elver trap SOP is available.

Compliance with international recommendations: Indicate ‘Y’ (yes) if the sampling design is in line with international recommendations, and ‘N’ if not. If no relevant expert or coordination groups exist, the sampling design should be shortly explained in the text, and should be available upon request for the evaluators.

Y

Link to sampling protocol documentation: Provide a link to a webpage where the documentation can be found. If no link is available, but documentation exists, provide a literature reference (author(s), year and type of publication - e.g. internal report). If no documentation on the sampling design exists, provide details on the sampling protocol in this textbox.

[Sampling protocols – need to put up on Ireland dcmap webpage]

Compliance with international recommendations: Member State shall state ‘Y’ (yes) if the sampling protocol is in line with international recommendations, and ‘N’ if not. If no relevant expert or coordination groups exist, the sampling protocol should be shortly explained in the text, and should be available upon request for the evaluators.

Y

The EIFAAC/ICES/GFCM working group on eel have a number of reports outlining the data requirements for international assessments and recommendations. These include but are not limited to ICES 2008, 2007, the WKESDCF report from 2012.

https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2008/WGEEL/wgeel_2008_final.pdf

<https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2007/WGEEL/2007%20EIFAC-ICES%20Report-Final-01-09-08.pdf>

<https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2012/WKESDCF/WKESDCF%20report%202012.pdf>

Sampling implementation
Recording of refusal rate: Indicate with 'Y' (yes) or 'N' (no), or 'NA' (not applicable, in case of research surveys). If 'N' (no), indicate when (year) documentation will be available. NA Monitoring of sampling progress within the sampling year: Indicate how sampling allocations are adjusted (if needed) and followed-up, what are the mechanisms in place to resolve issues and adopt mitigation measures during the sampling year? GANTT Charts are created annually to keep track of planned surveys with room for flexibility if a survey needs to be rescheduled due to poor weather or unforeseen events.
Data capture
Means of data capture: short description (+ photo optionally). Indicate what are the means for collecting the data, e.g. scales, measuring board, dedicated software, ... Data for recruits, fyke nets, electrofishing and silver eel fishing are captured using Survey123 and uploaded to the cloud for storage. A quality control procedure is then carried out on the data. If the digital capture fails on the day a paper version is captured. Equipment required include measuring board for length, scales for weight, calipers for eye and fin measurements, scanners for tag detection Data capture documentation: Provide a link to a webpage where the documentation can be found. If no link is available, but documentation exists, provide a literature reference (author(s), year and type of publication - e.g. internal report). If no documentation on data capture (e.g. measuring protocols, maturity staging, manual for the data capture means etc.) exists, provide some details in the textbox. Data capture protocols are available Quality checks documentation: Indicate with 'Y' (yes) or 'N' (no). If 'N' (no), indicate when (year) documentation will be available. Provide a link to a webpage where the documentation can be found. If no link is available, but documentation exists, provide a literature reference (author(s), year and type of publication - e.g. internal report). If no documentation on the quality checks exists, provide some details in the text box. Y

Data storage
National database: SQL database for Eels and ARCGIS geodatabase International database: WGEEL database Quality checks and data validation documentation: Y An SOP document outlines the policy and procedure for the Inputting and Quality Check of Eel Survey Data Data is checked during extractions for end-users such as ICES / European Commission.
Sample storage
Otoliths for aging are stored clean and dry at room temperature. These age structures are stored at Castlehouse before preparation and age reading is carried out. Eel specimens are frozen and stored in Castlehouse facility until processed in the wet laboratory. All necessary information is recorded and extracted and the remains disposed of in the biological waste facility.
Data processing
Evaluation of data accuracy (bias and precision): Y A number of documents relate to the inputting and quality check of data from fieldwork and laboratory work. A number of aging workshops are held with colleagues within Ireland to ensure a standardisation in method and agreement within the country. The overall aim of our qc endeavour for aging otoliths is for: - o Mean Percentage Agreement to be preferably above >90%, - o All individual % Agreement Scores on the Graph to be >80%, - o OUI Class Scores of 1 are preferable however, OUI Class 2 would be acceptable but only on older eels. We do not want any OUI Class 3 scores if possible. Class 3 specimens should be re-read with both readers to reach consensus, - o Agree difficult otolith with both readers (original and QC readers), - o If there are a lot of differences in a sample; to agree to widen QC sample; and failing that to re-age the entire site of eels. Editing and imputation methods: Y Any errors detected during the QC procedure or during the analysis and report writing is immediately corrected in the raw data files. Quality document associated to a dataset: Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors

or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g. in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

No Deviations

Sampling Design and Protocols

No Deviations


Sampling Implementation

100% achieved for the Corrib trap but under sampled for the Inagh trap.

For the Inagh trap excessive high water levels during the elver migration period meant the trap could not be serviced. The trap is located in the river and can only be monitored when conditions of low flow are experienced. Extensive rainfall for July and August impacted on the surveys. The Corrib elver trap is located in a fish ramp and is accessible under a wider range of conditions.

Data Capture

The elver data is now captured through the Survey 123forms on ArcGIS, the forms are uploaded to a geodatabase stored in the cloud. A dashboard provides summary information on the status of the traps and quantity of elvers captured to date (See image below).



Data Storage

Laboratory samples are documented using ArcGIS Survey 123 form, replacing the paper based format. This data is then uploaded to a geodatabase.

Sample Storage

No Deviations
<u>Data Processing</u>
No Deviations

Sampling scheme identifier: Eel_Silver_Diadromous (scientific)

MS : Ireland
Region: North-East Atlantic
Sampling scheme identifier: Eel_Silver_Diadromous (scientific)
Sampling scheme type: Diadromous (scientific)
Observation type: SciObs water body
Time period of validity: 2022-2027
Short description (max 100 words): <i>e.g. sampling scheme aiming at collecting length samples from commercial landings on-shore for all species listed in Table 1 of the EU MAP Delegated Decision annex. The scheme covers mainland and all outermost regions ('RUP' in French, Portuguese, and Spanish).</i> The silver eel index sites will be monitored to record the silver eel migration providing a time series data both within a year and between years. The data provides count of silver eels, mark recapture studies along with length frequency, weight biological information including length, weight, eye and fin measurements. Samples will be brought back to the laboratory for age, growth, sex, parasite prevalence and swimbladder damage indices. The escapement data is used in the IMESE model to estimate silver eel production and escapement as required under the Eel Regulation.
Description of the population
Population targeted: Specify which are the primary sampling units (PSU), e.g. all national port*days (information present in former Table 4B). For research surveys: specify the main target species from a survey perspective (as opposed to Table 1 in the Annex to the Implementing Decision) and the main survey area. The silver eel index surveys target the migrating silver eel cohort of the eel population

upstream of the fishing site. The site is fished by former eel fishers and it is fished 20-30 nights from September to December to capture the variability in eels biological parameters across the season.

Population sampled: Specify which part of the target population will be sampled and specify which part of the target population is unreachable for sampling or excluded for some reason to explain, *e.g. major ports being listed as auctions excluding all minor ports and no sampling during the week-ends*. For research surveys at sea describe target species in single-species surveys or ecosystem component (*e.g. demersal, pelagic*) in multispecies surveys.

For silver eel index sites the length profile is 30-96cm and corresponds to the migrating adult silver eel population. The sample does include eels that would be classified as yellow eel however the uncertainty around the silvering process means its difficult to say that these eels will not migrate.

Stratification: Explain the logic taken to stratify the population and the number of strata generated, *e.g. population stratified in 3 geographical lots (from A to B, from B to C and from C to D). Each lot is then stratified by auction*.

For reporting to the EU under the Eel Regulation Ireland created Eel Management Units (in line with River Basin Districts). The recommendation from WKESDCF and highlighted within the dcmep directive is for each MS to monitor all eel lifestages within at least one catchment per RBD.

In Ireland the RBD or eel management units are East EMU, SouthEast EMU, SouthWest EMU, Shannon International EMU, West EMU and the NorthWest EMU. In addition there are index catchments with historical information available that is targeted in this programme to continue the longterm temporal trend.

Sampling design and protocols

Sampling design description: Describe how the sampling allocation is defined; how PSU and SSU are selected for sampling; indicate for which catch fraction the sampling scheme applies.

The silver eel migration typically occurs around the new moon phase when rivers are in flood these nights are targeted for fishing. Outside these nights, nets are also set if a significant rise in water levels is observed as this can entice eels to migrate outside of the new moon phase. The number of nights fished varies year to year due to the environmental conditions. At a minimum 300-500 eels are required to be measured for length to capture a representative length frequency.

Is the sampling design compliant with the 4S principle?: Y/N/NA (NA for e.g. surveys and diadromous and recreational sampling schemes)

NA

Regional coordination: Indicate if the sampling design and protocols were developed as part of a regional or multi-lateral agreement, and if yes, refer to the agreement (table 1.3) and list all MS participating.

No regional coordination is undertaken for sampling eel. However for the transboundary North West EMU consultation occurs between agencies in Northern Ireland and Republic of Ireland in relation to combined eel surveys on occasion.

Link to sampling design documentation: Provide a link to a webpage where the documentation can be found. If no link is available, but documentation exists, Member State shall provide a literature reference (author(s), year and type of publication - e.g. internal report). If no documentation on the sampling design exists, Member State shall provide some details in the textbox.

Silver eel sampling design. Locations are the site of former commercial eel fisheries and are now run on a research basis. The silver eel migration is dependent on environmental conditions (new moon phase and flood conditions) over an extended period of time; August to January. Twenty – 30 nights are fished during relevant conditions during this time frame.

The sites are fished in a standardised way year to year to allow a comparison across time.

Compliance with international recommendations: Indicate ‘Y’ (yes) if the sampling design is in line with international recommendations, and ‘N’ if not. If no relevant expert or coordination groups exist, the sampling design should be shortly explained in the text, and should be available upon request for the evaluators.

Y

Link to sampling protocol documentation: Provide a link to a webpage where the documentation can be found. If no link is available, but documentation exists, provide a literature reference (author(s), year and type of publication - e.g. internal report). If no documentation on the sampling design exists, provide details on the sampling protocol in this textbox.

[Sampling protocols – need to put up on Ireland dcmap webpage]

Compliance with international recommendations: Member State shall state 'Y' (yes) if the sampling protocol is in line with international recommendations, and 'N' if not. If no relevant expert or coordination groups exist, the sampling protocol should be shortly explained in the text, and should be available upon request for the evaluators.

Y

The EIFAAC/ICES/GFCM working group on eel have a number of reports outlining the data requirements for international assessments and recommendations. These include but are not limited to ICES 2008, 2007, the WKESDCF report from 2012.

https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2008/WGEEL/wgeel_2008_final.pdf

<https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2007/WGEEL/2007%20EIFAC-ICES%20Report-Final-01-09-08.pdf>

<https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2012/WKESDCF/WKESDCF%20report%202012.pdf>

Sampling implementation

Recording of refusal rate: Indicate with 'Y' (yes) or 'N' (no), or 'NA' (not applicable, in case of research surveys). If 'N' (no), indicate when (year) documentation will be available.

NA

Monitoring of sampling progress within the sampling year: Indicate how sampling allocations are adjusted (if needed) and followed-up, what are the mechanisms in place to resolve issues and adopt mitigation measures during the sampling year?

GANTT Charts are created annually to keep track of planned surveys with room for flexibility if a survey needs to be rescheduled due to poor weather or unforeseen events.

Data capture

Means of data capture: short description (+ photo optionally). Indicate what are the means for collecting the data, e.g. scales, measuring board, dedicated software, ...

Data for recruits, fyke nets, electrofishing and silver eel fishing are captured using Survey123 and uploaded to the cloud for storage. A quality control procedure is then carried out on the

data. If the digital capture fails on the day a paper version is captured. Equipment required include measuring board for length, scales for weight, calipers for eye and fin measurements, scanners for tag detection Data capture documentation: Provide a link to a webpage where the documentation can be found. If no link is available, but documentation exists, provide a literature reference (author(s), year and type of publication - e.g. internal report). If no documentation on data capture (e.g. measuring protocols, maturity staging, manual for the data capture means etc.) exists, provide some details in the textbox. Data capture protocols are available Quality checks documentation: Indicate with 'Y' (yes) or 'N' (no). If 'N' (no), indicate when (year) documentation will be available. Provide a link to a webpage where the documentation can be found. If no link is available, but documentation exists, provide a literature reference (author(s), year and type of publication - e.g. internal report). If no documentation on the quality checks exists, provide some details in the text box. Y
Data storage National database: SQL database for Eels and ARCGIS geodatabase International database: WGEEL database Quality checks and data validation documentation: Y An SOP document outlines the policy and procedure for the Inputting and Quality Check of Eel Survey Data Data is checked during extractions for end-users such as ICES / European Commission.
Sample storage Otoliths for aging are stored clean and dry at room temperature. These age structures are stored at Castlehouse before preparation and age reading is carried out. Eel specimens are frozen and stored in Castlehouse facility until processed in the wet laboratory. All necessary information is recorded and extracted and the remains disposed of in the biological waste facility.
Data processing Evaluation of data accuracy (bias and precision): Y A number of documents relate to the inputting and quality check of data from fieldwork and laboratory work.

A number of aging workshops are held with colleagues within Ireland to ensure a standardisation in method and agreement within the country.

The overall aim of our qc endeavour for aging otoliths is for:

- o Mean Percentage Agreement to be preferably above >90%,
- o All individual % Agreement Scores on the Graph to be >80%,
- o OUI Class Scores of 1 are preferable however, OUI Class 2 would be acceptable but only on older eels. We do not want any OUI Class 3 scores if possible. Class 3 specimens should be re-read with both readers to reach consensus,
- o Agree difficult otolith with both readers (original and QC readers),
- o If there are a lot of differences in a sample; to agree to widen QC sample; and failing that to re-age the entire site of eels.

Editing and imputation methods: Y

Any errors detected during the QC procedure or during the analysis and report writing is immediately corrected in the raw data files.

Quality document associated to a dataset:

Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g. in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

Deviations as outlined below

Sampling Design and Protocols

No Deviations

Sampling Implementation

Due to limited staff resources on the Eel Monitoring Programme team in 2023, mark recaptures studies were not undertaken to assess the silver eel migrating population. It is anticipated that the staff shortages will be addressed ahead of the 2024 silver eel season.

Data Capture

Laboratory samples are documented using ArcGIS Survey 123 form, replacing the paper based format. This data is then uploaded to a geodatabase

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: Eel_Fykenet_Diadromous (scientific)

MS : Ireland

Region: North-East Atlantic

Sampling scheme identifier:

Eel_Fykenet_Diadromous (scientific)

Sampling scheme type:

Eel_Silver_Diadromous (scientific)

Observation type:

SciObs water body

Time period of validity: 2022-2027

Short description (max 100 words): *e.g. sampling scheme aiming at collecting length samples from commercial landings on-shore for all species listed in Table 1 of the EU MAP Delegated Decision annex. The scheme covers mainland and all outermost regions ('RUP' in French, Portuguese, and Spanish).*

The yellow eel fyke net surveys will provide samples from the local eel population. Fyke nets can be set in lakes, large rivers and transitional waters. This method will provide samples for count of eels, mark recapture studies, biological information including length, weight, eye and fin measurements. Samples will be brought back to the laboratory for age, growth, sex, parasite prevalence and swimbladder damage indices.

Description of the population

Population targeted: Specify which are the primary sampling units (PSU), e.g. all national port*days (information present in former Table 4B). For research surveys: specify the main target species from a survey perspective (as opposed to Table 1 in the Annex to the Implementing Decision) and the main survey area.

The fyke net surveys target the local eel population in the vicinity of the netting location in

the waterbody surveyed, e.g. lake, large river and the transitional waterbody

Population sampled: Specify which part of the target population will be sampled and specify which part of the target population is unreachable for sampling or excluded for some reason to explain, *e.g. major ports being listed as auctions excluding all minor ports and no sampling during the week-ends*. For research surveys at sea describe target species in single-species surveys or ecosystem component (*e.g. demersal, pelagic*) in multispecies surveys.

For fyke nets the population sampled is the eel population >30cm which is the typical size of eels captured in fyke nets, smaller eels can escape through the mesh of the nets and while they can be caught it is not representative of the local population.

Stratification: Explain the logic taken to stratify the population and the number of strata generated, *e.g. population stratified in 3 geographical lots (from A to B, from B to C and from C to D)*. Each lot is then stratified by auction.

For reporting to the EU under the Eel Regulation Ireland created Eel Management Units (in line with River Basin Districts). The recommendation from WKESDCF and highlighted within the DCMAP directive is for each MS to monitor all eel lifestages within at least one catchment per RBD.

In Ireland the RBD or eel management units are East EMU, SouthEast EMU, SouthWest EMU, Shannon International EMU, West EMU and the NorthWest EMU. In addition there are index catchments with historical information available that is targeted in this programme to continue the longterm temporal trend.

Sampling design and protocols

Sampling design description: Describe how the sampling allocation is defined; how PSU and SSU are selected for sampling; indicate for which catch fraction the sampling scheme applies.

Within a lake a random sampling design is employed. In lakes with historical survey information repeat sampling is employed to maintain long-term datasets.

In large Rivers a representative stretch of river is surveyed with number of nets distributed along the river length or target area. Access for boats is a consideration for ease of setting and retrieving nets. For Rivers with historical survey information repeat samples are undertaken to maintain long-term data sets.

Within Estuaries a random design along with targeted locations to compare with historical survey locations is employed.

In Lakes each chain consists of 5 fyke nets. Eight chains will be set nightly to capture the variation in the spatial distribution of eels around the lakes. This information will be consistent with data gathered under the eel monitoring programme from 2009 to present.

Is the sampling design compliant with the 4S principle?: Y/N/NA (NA for e.g. surveys and diadromous and recreational sampling schemes)

NA

Regional coordination: Indicate if the sampling design and protocols were developed as part of a regional or multi-lateral agreement, and if yes, refer to the agreement (table 1.3) and list all MS participating.

No regional coordination is undertaken for sampling eel. However for the transboundary North West EMU consultation occurs between agencies in Northern Ireland and Republic of Ireland in relation to combined eel surveys on occasion.

Link to sampling design documentation: Provide a link to a webpage where the documentation can be found. If no link is available, but documentation exists, Member State shall provide a literature reference (author(s), year and type of publication - e.g. internal report). If no documentation on the sampling design exists, Member State shall provide some details in the textbox.

In order to standardise the influence of environmental conditions and ensure the catch reflects the actual eel population as suggested by Harley et al., 2001, lakes were surveyed between June and September each year, with the surveys carried out over 2-3 visits. The nets were randomly assigned to the lake using a trap builder in the software Density 4 to remove operator bias and ensure good coverage of the area (Efford et al. 2004). The nets were set in chains of five fyke nets tied end to end and set at least 50 m apart to avoid interference between chains. The chains of nets were lifted daily to avoid gear saturation and all eels counted per net end. To standardise the difference in lake size and effort; large lakes (for example Lough Derg) were divided into upper and lower and sampled as 2 lakes and therefore had twice the effort of the smaller lakes.

Compliance with international recommendations: Indicate 'Y' (yes) if the sampling design is in line with international recommendations, and 'N' if not. If no relevant expert or coordination groups exist, the sampling design should be shortly explained in the text, and should be available upon request for the evaluators.

Y

Link to sampling protocol documentation: Provide a link to a webpage where the documentation can be found. If no link is available, but documentation exists, provide a literature reference (author(s), year and type of publication - e.g. internal report). If no documentation on the sampling design exists, provide details on the sampling protocol in this textbox.

[Sampling protocols – need to put up on Ireland DCMAP webpage]

Compliance with international recommendations: Member State shall state ‘Y’ (yes) if the sampling protocol is in line with international recommendations, and ‘N’ if not. If no relevant expert or coordination groups exist, the sampling protocol should be shortly explained in the text, and should be available upon request for the evaluators.

The EIFAAC/ICES/GFCM working group on eel have a number of reports outlining the data requirements for international assessments and recommendations. These include but are not limited to ICES 2008, 2007, the WKESDCF report from 2012.

https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2008/WGEEL/wgeel_2008_final.pdf

<https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2007/WGEEL/2007%20EIFAC-ICES%20Report-Final-01-09-08.pdf>

<https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2012/WKESDCF/WKESDCF%20report%202012.pdf>

Y –

Sampling implementation

Recording of refusal rate: Indicate with 'Y' (yes) or 'N' (no), or ‘NA’ (not applicable, in case of research surveys). If 'N' (no), indicate when (year) documentation will be available.

NA

Monitoring of sampling progress within the sampling year: Indicate how sampling allocations are adjusted (if needed) and followed-up, what are the mechanisms in place to resolve issues and adopt mitigation measures during the sampling year?

GANTT Charts are created annually to keep track of planned surveys with room for flexibility if a survey needs to be rescheduled due to poor weather or unforeseen events.

Data capture

Means of data capture: short description (+ photo optionally). Indicate what are the means for collecting the data, e.g. scales, measuring board, dedicated software, ...

Data for recruits, fyke nets, electrofishing and silver eel fishing are captured using Survey123 and uploaded to the cloud for storage. A quality control procedure is then carried out on the data. If the digital capture fails on the day a paper version is captured.

Equipment required include measuring board for length, scales for weight, calipers for eye and fin measurements, scanners for tag detection

Data capture documentation: Provide a link to a webpage where the documentation can be found. If no link is available, but documentation exists, provide a literature reference (author(s), year and type of publication - e.g. internal report). If no documentation on data capture (e.g. measuring protocols, maturity staging, manual for the data capture means etc.) exists, provide some details in the textbox.

Data capture protocols are available

Quality checks documentation: Indicate with 'Y' (yes) or 'N' (no). If 'N' (no), indicate when (year) documentation will be available. Provide a link to a webpage where the documentation can be found. If no link is available, but documentation exists, provide a literature reference (author(s), year and type of publication - e.g. internal report). If no documentation on the quality checks exists, provide some details in the text box.

Y

Control quality procedures exist for all data generated in the field and laboratory including length, weight regression analyses and identification, removal/replacement of outlier data.

Laboratory QC of eel dissection data, parasite data and swimbladder indices results are carried out with each new sample. An intensive QC methodology of all ageing and growth data is in place and constantly improved in order to gain accurate data over time.

Data storage

National database: SQL database for Eels and ARCGIS geodatabase

International database: WGEEL database

Quality checks and data validation documentation: Y

An SOP document outlines the policy and procedure for the Inputting and Quality Check of Eel Survey Data

Data is checked during extractions for end-users such as ICES / European Commission.

Sample storage**Data processing****Evaluation of data accuracy (bias and precision): Y**

A number of documents relate to the inputting and quality check of data from fieldwork and laboratory work.

A number of aging workshops are held with colleagues within Ireland to ensure a standardisation in method and agreement within the country.

The overall aim of our qc endeavour for aging otoliths is for:

- o Mean Percentage Agreement to be preferably above >90%,
- o All individual % Agreement Scores on the Graph to be >80%,
- o OUI Class Scores of 1 are preferable however, OUI Class 2 would be acceptable but only on older eels. We do not want any OUI Class 3 scores if possible. Class 3 specimens should be re-read with both readers to reach consensus,
- o Agree difficult otolith with both readers (original and QC readers),
- o If there are a lot of differences in a sample; to agree to widen QC sample; and failing that to re-age the entire site of eels.

Editing and imputation methods: Y

Any errors detected during the QC procedure or during the analysis and report writing is immediately corrected in the raw data files.

Quality document associated to a dataset:

Validation of the final dataset: Data is checked during extractions for end-users such as ICES / European Commission - the checks used will depend on the use of the data. If errors or anomalies are observed, then data is either corrected by reference to the original data sheets (e.g. in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

Due to Staffing Shortage, this sampling could not be completed, however steps have been taken to ensure it will be completed in 2024 and beyond.

<u>Sampling Design and Protocols</u>
No Deviations
<u>Sampling Implementation</u>
No Deviations
<u>Data Capture</u>
Laboratory samples are documented using ArcGIS Survey 123 form, replacing the paper based format. This data is then uploaded to a geodatabase
<u>Data Storage</u>
No Deviations
<u>Sample Storage</u>
No Deviations
<u>Data Processing</u>
No Deviations

Quality Reports Related to Table 2.4 and Text Box 2.4

Sampling scheme identifier: IMREC_CREEL

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: IMREC_CREEL
Sampling scheme type: Recreational fisheries
Observation type: SciObsOnShore
Time period of validity: 2022-Ongoing
Short description (max 100 words): The IMREC_CREEL survey of shore and small boat anglers utilises a spatio-temporal sampling method to collect catch per unit effort (CPUE) data of sea anglers around the Irish coast, whereby the unit of effort is a daily angler session.
Description of the population
Population targeted: All species caught by recreational shore and small boat anglers in Irish waters.
Population sampled: All shore and small boat anglers catch is included in the sampling programme.

Stratification:

Sampling is stratified bi-annually and recreational fisheries are grouped by geographic region (Irish Sea/Celtic Sea/ West coast of Ireland).

Sampling design and protocols**Sampling design description:**

IMREC_CREEL embraces the on-site shore and small boat angling catch surveys. For shore angling IMREC_CREEL uses a roving creel survey, with follow up call-back to estimate shore angling catch rates, which utilises a spatio-temporal sampling frame to collect catch per unit effort (CPUE) data of sea anglers around the Irish coast. This approach was applied for the pilot study (Ryan et al., 2021) due to the disparate nature of shore angling around Ireland and the multiple potential access points to the sea. PSUs consist of connected polygons which encompass all fishable sections of Irelands shoreline. The number of spatial strata and temporal strata in the sampling frame will be reduced from five to three and from four to two respectively, based on learnings from the pilot study (Ryan et al., 2021). This amendment will decrease the requirement for imputation procedures to account for missing data points without adversely affecting precision. Sampling effort will focus more on regions and times where angling effort is known to be higher.

The small boat aspect of IMREC_CREEL uses a similar spatio-temporal sampling frame where catch data are collected through a refined random-access point survey. Pilot study sampling difficulties for the small boat sector (due to multiple access points resulting in limited sampling encounters) resulted in small sample sizes and substantial catch estimate biases. As this sector is an important component of MRF catches in Ireland continued sampling refinement and development of revised techniques to monitor catch rates will be prominent in this survey.

The spatio-temporal sampling frame consists of the whole coastline split into separate sampling polygons X 365 separate sampling days (PSUs). This sampling frame embraces every angling trip on the Irish coast over a calendar year. All catch records will be noted as 'retained' or 'released'. This will allow an estimation of the rate of fish retained and returned alive per species.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: N

Link to sampling design documentation:

Ryan. D., Leonard, E., Casserly. C. M., Roche. W. (2021). Marine Recreational Angling Catches in Ireland. Pilot Study Report (2019-2021). Inland Fisheries Ireland. Dublin.

Compliance with international recommendations: Y.

The sampling programme was designed with guidance from members of the ICES WGRFS.

Ireland's report on the DCF Annual Work Plan 2020 included a report on the pilot study which was submitted to the EC STCEF for review in March 2021. The sampling programme was deemed to be satisfactory. The pilot study and the ongoing refinements will guide the current sampling programme.

<https://stecf.jrc.ec.europa.eu/documents/43805/3064868/STECF+21-09+-+Evaluation+of+AR+and+DTi.pdf/058b6ae3-ac98-4530-8a2f-be26fc42cf46?version=1.0>

Link to sampling protocol documentation:

Ryan. D., Leonard, E., Casserly. C. M., Roche. W. (2021). Marine Recreational Angling Catches in Ireland. Pilot Study Report (2019-2021). Inland Fisheries Ireland. Dublin.

Compliance with international recommendations: Y.

The sampling programme was designed with guidance from members of the ICES WGRFS. A report on the pilot study which guided the current sampling programme was submitted to the EC STCEF for review in March 2021. The sampling programme was deemed to be satisfactory.

Sampling implementation

Recording of refusal rate: Y

Refusals are recorded

Monitoring of sampling progress within the sampling year:

The sampling programme is planned to sample strata in accordance with the spatio-temporal sampling frame design. Sampling progress is continuously monitored to ensure that all strata in the sampling frame have sufficient data points to avoid imputation. If sampling coverage is lower than expected due to unforeseen issues, targeted sampling may be undertaken to gap-fill.

Data capture

Means of data capture:

Data are collected electronically on tablets via the ArcGIS Survey123© data collection tool. This resource allows instant data capture and safe storage to a centralised geodatabase. Surveyors carry measuring boards to measure lengths of captured fish when applicable.

Data capture documentation:

Reference SOPs for surveyors using the Survey123 tool are available and reviewed regularly.

Further information on processes in:

Ryan. D., Leonard, E., Casserly. C. M., Roche. W. (2021). Marine Recreational Angling Catches in Ireland. Pilot Study Report (2019-2021). Inland Fisheries Ireland. Dublin.

Quality checks documentation:

N (year for documentation to be available to be confirmed).

Data recorded during on-site interviews are uploaded to a centralised database. Data are

regularly collated, downloaded and visualised using customised R scripts to quality check for erroneous data points before assessments are undertaken.
Data storage
National database: ArcGIS geodatabase and IFI SQL server. International database: NA Quality checks and data validation documentation: Not available at present. Quality checks/reviews of MS methods are reviewed periodically by ICES WGRFS (QAT https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/EOSG/2019/WGRFS2019.pdf). When MS catch reports are submitted to end users (European Commission) standard quality checks will be in place with accompanying documentation.
Sample storage
Physical samples are not collected for this programme. All digital data collected are stored in a centralised geodatabase which is backed up hourly.
Data processing
Evaluation of data accuracy (bias and precision): N. No specific documentation is currently available. Further consultation with ICES WGRFS will determine what further processes can be put in place during 2022. CPUE estimates computed through the design-based analysis of on-site survey data are assessed for precision at analysis. See Ryan. D., Leonard, E., Casserly. C. M., Roche. W. (2021). Marine Recreational Angling Catches in Ireland. Pilot Study Report (2019-2021). Inland Fisheries Ireland, Dublin. Editing and imputation methods: N. No specific documentation is currently available. Further consultation with the ICES WGRFS will determine what further processes can be put in place during 2022. If errors are identified in the database they are removed from the dataset. The sampling programme was designed to avoid the use of imputation procedures if possible. However, if necessary, it may be appropriate to merge unsampled strata with adjacent strata with similar angling patterns. Quality document associated to a dataset: N No DOI currently exists for the dataset. No estimation process document is currently available.

Validation of the final dataset:

Data will be rechecked by the database manager prior to submission to end users. If potential errors are detected they will be crosschecked with original datasets and corrected if possible. If this is not possible, erroneous data will be removed prior to submission.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

Population targeted & sampled as described. Number of spatial strata reduced from 3 to 2 (Irish Sea & Celtic Sea regions combined and renamed “Eastcoast”).

Sampling Design and Protocols

No Deviations

Sampling Implementation

>95% of sampling schedule was achieved

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

The sampling programme has followed well established probability-based sampling methods. However, there has been no external evaluation of data accuracy. The WGRFS uses a quality assurance tool (QAT), which can assess survey quality. This survey will be assessed in the future.

Sampling scheme identifier: IMREC_SKP_DI

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: IMREC_SKP_DI
Sampling scheme type: Recreational fisheries
Observation type: SelfAtSea
Time period of validity: 2022-Ongoing
Short description (max 100 words):

An angling charter skipper catch diary (IMREC_SKP_DI) is in use which allows a sample of charter skippers to self-report fish catch, and catch and release rates at species level for each angling trip. This process will also be used to provide data to support estimations of angling effort across the Irish charter fleet.

Description of the population

Population targeted:

All species caught by charter anglers in Irish waters.

Population sampled:

All catch taken by recreational charter anglers is included in the sampling programme.

Stratification:

Sampling includes all coastlines annually and recreational angling fisheries are grouped by geographic region (Irish Sea/Celtic Sea/ West coast of Ireland).

Sampling design and protocols

Sampling design description:

A charter skipper lead catch diary was developed during the pilot study (Ryan et al, 2021) which allowed contributors to enter catch data on species, lengths (and catch and release rates) for each sampling trip. Charter angling is primarily a weather- dependent leisure activity and most effort tends to be around and during the summer months. Regional stratification was also deemed appropriate as targeted species are often dependent on region. Examination of historical voluntary charter skipper catch diaries collated by IFI (up to 2008) confirmed that catch is disparate between regions.

The total number of registered active charter skippers is currently low in Ireland (< 70) and a diary has been provided to all willing participants. Contributors are requested to record all angling trips and associated catch data. Catch is reported by vessel per trip and divided by the number of anglers on board to provide a mean catch per angler per day.

The data streams from this sampling programme will be combined with IMREC_OB_CH_SURVEY at analysis, to provide an estimate of charter angling activity and catch rates.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: N

Link to sampling protocol documentation:

Ryan. D., Leonard, E., Casserly. C. M., Roche. W. (2021). Marine Recreational Angling Catches in Ireland. Pilot Study Report (2019-2021). Inland Fisheries Ireland. Dublin.

Compliance with international recommendations: Y The sampling programme was designed through guidance from members of the ICES WGRFS. A report on the pilot study which guided the current sampling programme was submitted to the EC STCEF for review in March 2021. The sampling programme was deemed to be satisfactory.
Sampling implementation
Recording of refusal rate: NA
Monitoring of sampling progress within the sampling year: All contributors are contacted regularly during the sampling window for progress updates and identify any sampling issues. They are also requested to share data sheets if available.
Data capture
Means of data capture: Species capture and length data are initially recorded in a hardcopy, waterproof diary. Contributors are requested to take a digital image of all angling trip data and share it with the designated IFI data manager whenever possible. These data, when received, are transcribed to the IMREC centralized database application. Length data is recorded using measuring boards provided by IFI. Data capture documentation: All skippers contributing to the programme are supplied with an SOP which is regularly reviewed and updated. Further details provided in: Ryan, D., Leonard, E., Casserly, C. M., Roche, W. (2021). Marine Recreational Angling Catches in Ireland. Pilot Study Report (2019-2021). Inland Fisheries Ireland. Dublin. Quality checks documentation: N (year for documentation to be available to be confirmed). Data are regularly collated, downloaded and visualised using customised R scripts to quality check for erroneous data points before assessments are undertaken.
Data storage
National database: ArcGIS geodatabase and IFI SQL server. International database: NA Quality checks and data validation documentation: Not available at present. Quality checks/reviews of MS methods are reviewed periodically by ICES WGRFS (QAT https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/EOSG/2)

[019/WGRFS2019.pdf](#)).

When MS catch reports are submitted to end users (European Commission) standard quality checks will be in place with accompanying documentation.

Sample storage

Physical samples are not collected for this programme.

Data processing

Evaluation of data accuracy (bias and precision): N.

No specific documentation is currently available. Further consultation with ICES WGRFS will determine what further processes can be put in place during 2022.

CPUE estimates computed through the design-based analysis of on-site survey data are assessed for precision at analysis.

See Ryan. D., Leonard, E., Casserly. C. M., Roche. W. (2021). Marine Recreational Angling Catches in Ireland. Pilot Study Report (2019-2021). Inland Fisheries Ireland, Dublin.

Editing and imputation methods: N.

No specific documentation is currently available. Further consultation with the ICES WGRFS will determine what further processes can be put in place during 2022.

If errors are identified in the database they are removed from the dataset. The sampling programme was designed to avoid the use of imputation procedures if possible. However, if necessary, it may be appropriate to merge unsampled strata with adjacent strata with similar angling patterns.

Quality document associated to a dataset: N

No DOI currently exists for the dataset. No estimation process document is currently available.

Validation of the final dataset:

Data will be rechecked by the database manager prior to submission to end users. If potential errors are detected they will be crosschecked with original datasets and corrected if possible. If this is not possible, erroneous data will be removed prior to submission.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

Population targeted & sampled as described. Number of spatial strata reduced from 3 to 2 (Irish Sea & Celtic Sea regions combined and renamed “Eastcoast”).

Sampling Design and Protocols

Due to relatively low participation by charter skippers, this data stream was not combined with IMREC_OB_CH_SURVEY at analysis stage, to provide an estimate of charter angling activity and catch rates in 2023. Improved participation is expected from 2024 onwards following recent further engagement with the sector.

Sampling Implementation

As a result of contact with contributors during the sampling window in 2023, it was found that sampling coverage was insufficient to make a precise estimate of catch rates, due to difficulty in completing the diary. As a result, the diary was amended to make it more user friendly for sampling in 2024.

Data Capture

Communication with contributors found that it was unrealistic for many to record length data for species captured. This observation led to a revision of the diary, which has been implemented for 2024.

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: IMREC_OB_CH_SURVEY

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: IMREC_OB_CH_SURVEY
Sampling scheme type: Recreational fisheries
Observation type: SciObsAtSea
Time period of validity: 2022-Ongoing
Short description (max 100 words): An onsite sampling programme (IMREC_OB_CH_SURVEY) will be undertaken to randomly sample designated chartered angling trips to collect detailed species, length and weight data. This will add to data collected through the ongoing charter skipper diary (IMREC_SKP_DI). Onboard surveyors record species numbers caught, and accurately measure lengths and weights of all captured and released fish.
Description of the population
Population targeted: All species caught by charter anglers in Irish waters.

Population sampled:

All catch taken by recreational charter anglers is included in the sampling programme.

Stratification:

Sampling is stratified bi-annually and recreational fisheries are grouped by geographic region (Irish Sea/Celtic Sea/ West coast of Ireland).

Sampling design and protocols**Sampling design description:**

The sampling frame consists of a panel of participating charter skippers stratified according to their operating region. Samplers are assigned to participating charter skippers based on a stratified random sampling protocol. Samplers record angling trip data including:

- Species caught
- Whether released or retained
- Total length for the first 60 fish of each species (30 retained and 30 released fish)
- Individual weights (g) for the first 60 fish of each species (30 retained and 30 released fish)
- Total counts of all other captured fish (both released and retained) that were not measured or weighed

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: N

Link to sampling protocol documentation:

Ryan. D., Leonard, E., Casserly. C. M., Roche. W. (2021). Marine Recreational Angling Catches in Ireland. Pilot Study Report (2019-2021). Inland Fisheries Ireland. Dublin.

Compliance with international recommendations: Y.

The sampling programme was designed with guidance from members of the ICES WGRFS. Ireland's report on the DCF Annual Work Plan 2020 included a report on the pilot study which was submitted to the EC STECF for review in March 2021. The sampling programme was deemed to be satisfactory. The pilot study and the ongoing refinements will guide the current sampling programme.

<https://stecf.jrc.ec.europa.eu/documents/43805/3064868/STECF+21-09+-+Evaluation+of+AR+and+DTi.pdf/058b6ae3-ac98-4530-8a2f-be26fc42cf46?version=1.0>

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year:

Regular communication with charter skippers and the sampling team is undertaken to monitor progress. For each temporal sampling run trips are monitored to ensure that data collection is occurring across all regional strata. If this is not the case, some targeted

sampling may be required to gap-fill.
Data capture
Means of data capture: Data are initially recorded on waterproof data recording sheets. At the end of each sampling day, surveyors take a digital image of all angling trip data and share this with the designated IFI data manager immediately. The data, when received, is immediately transcribed to the dedicated IFI IMREC database application. Length data are recorded using measuring boards provided by IFI. Data capture documentation: All skippers contributing to the programme are supplied with an SOP which is regularly reviewed and updated. Further details provided in: Ryan, D., Leonard, E., Casserly, C. M., Roche, W. (2021). Marine Recreational Angling Catches in Ireland. Pilot Study Report (2019-2021). Inland Fisheries Ireland. Quality checks documentation: N (year for documentation to be available to be confirmed). Data are regularly collated, downloaded and visualised using customised R scripts to quality check for erroneous data points before assessments are undertaken.
Data storage
National database: ArcGIS geodatabase and IFI SQL server. International database: NA Quality checks and data validation documentation: Not available at present. Quality checks/reviews of MS methods are reviewed periodically by ICES WGRFS (QAT https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/EOSG/2019/WGRFS2019.pdf). When MS catch reports are submitted to end users (European Commission) standard quality checks will be in place with accompanying documentation.
Sample storage
Physical samples are not collected for this programme.
Data processing
Evaluation of data accuracy (bias and precision): N. No specific documentation is currently available. Further consultation with ICES WGRFS will determine what further processes can be put in place during 2022.

CPUE estimates computed through the design-based analysis of on-site survey data are assessed for precision at analysis.

See: Ryan. D., Leonard, E., Casserly. C. M., Roche. W. (2021). Marine Recreational Angling Catches in Ireland. Pilot Study Report (2019-2021). Inland Fisheries Ireland, Dublin.

Editing and imputation methods: N.

No specific documentation is currently available. Further consultation with the ICES WGRFS will determine what further processes can be put in place during 2022.

If errors are identified in the database they are removed from the dataset. The sampling programme was designed to avoid the use of imputation procedures if possible. However, if necessary, it may be appropriate to merge unsampled strata with adjacent strata with similar angling patterns.

Quality document associated to a dataset: N

No DOI currently exists for the dataset. No estimation process document is currently available.

Validation of the final dataset:

Data will be rechecked by the database manager prior to submission to end users. If potential errors are detected they will be crosschecked with original datasets and corrected if possible. If this is not possible, erroneous data will be removed prior to submission.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

Population targeted & sampled as described. Number of spatial strata reduced from 3 to 2 (Irish Sea & Celtic Sea regions combined and renamed “Eastcoast”).

Sampling Design and Protocols

Staff resources/regional sampler availability was the major obstacle in achieving maximum sampling coverage. Extra staff resources in 2024 will mitigate this issue. Inclement weather also caused the cancellation of planned sampling days, on occasion.

Sampling Implementation

No Deviations

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

No Deviations

Sampling scheme identifier: IMREC_ANG_DI**MS : IRL****Region:** North-East Atlantic**Sampling scheme identifier:** IMREC_ANG_DI**Sampling scheme type:** Recreational fisheries**Observation type:** SelfOnShore**Time period of validity:** 2022-Ongoing

Short description (max 100 words):

A major output of the pilot study (Ryan et al., 2021) was the creation of an online angling diary (IMREC_ANG_DI). Sea anglers in Ireland have access to this online resource to input and self-report catch data. The diary will operate in parallel with the probability based IMREC_CREEL programmes.

Description of the population**Population targeted:**

All species caught by anglers in Irish waters

Population sampled:

All recreational sea anglers and their angling catches.

Stratification:

Online diary is available to all sea anglers. As anglers must provide general fishing location and the date of their fishing trip, catch data can be stratified temporally and spatially.

Sampling design and protocols**Sampling design description:**

To collect marine recreational angling catch data on a continuous basis, an online angler catch diary was developed. The online IMREC Angler Diary (Fig.1) provides a fast and efficient recording platform for self-selecting anglers to submit catch information on fishing trips (shore, small boat or charter boat). This platform provides qualitative data on species caught, total catches, catch/release rates and length measurements from a vetted group of participating anglers across Ireland.




Fig. 1: Structure of the IMREC Online Diary

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: N

Link to sampling protocol documentation:

Ryan. D., Leonard, E., Casserly. C. M., Roche. W. (2021). Marine Recreational Angling Catches in Ireland. Pilot Study Report (2019-2021). Inland Fisheries Ireland. Dublin

Compliance with international recommendations: Y.

The sampling programme was designed with guidance from members of the ICES WGRFS. Ireland's report on the DCF Annual Work Plan 2020 included a report on the pilot study which was submitted to the EC STECF for review in March 2021. The sampling programme was deemed to be satisfactory. The pilot study and the ongoing refinements will guide the current sampling programme.

<https://stecf.jrc.ec.europa.eu/documents/43805/3064868/STECF+21-09+-+Evaluation+of+AR+and+DTi.pdf/058b6ae3-ac98-4530-8a2f-be26fc42cf46?version=1.0>

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year:

Contributors to the sampling programme are monitored and email reminders to submit data are sent periodically.

Data capture

Means of data capture:

Contributors are self-selecting and must sign up to the programme to enable them to log their angling catch data. Data are collected electronically through an online 'diary' form via the ArcHub Survey123© data collection tool. This resource allows instant data capture and safe storage to the IMREC centralised geodatabase. Contributors are requested to provide information on their catch. Information includes: species, catch retained or released, catch length (measured or estimated) and catch weight (measured or estimated).

Data capture documentation:

All anglers contributing to the programme are supplied with an SOP which is regularly reviewed and updated. This is also available on the ArcHub© diary website after contributors sign up to the programme.

Further details provided in:

Ryan, D., Leonard, E., Casserly, C. M., Roche, W. (2021). Marine Recreational Angling Catches in Ireland. Pilot Study Report (2019-2021). Inland Fisheries Ireland. Dublin.

<https://imrec-ifigis.hub.arcgis.com/pages/get-involved>

Quality checks documentation:

N (year for documentation to be available to be confirmed).

Data submitted by contributors are immediately uploaded to a centralised database. Data are regularly collated, downloaded and visualised using customised R scripts to quality check for erroneous data points before assessments are undertaken.

Data storage**National database:**

ArcGIS geodatabase and IFI SQL server.

International database: NA

Quality checks and data validation documentation:

Not available at present.

Quality checks/reviews of MS methods are reviewed periodically by ICES WGRFS (QAT <https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/EOSG/2019/WGRFS2019.pdf>).

When country catch reports assessments are submitted to end users (European Commission) standard quality checks will be in place with accompanying documentation.

Sample storage

Physical samples are not collected for this programme. All digital data collected are stored in a centralised geodatabase which is backed up daily.

Data processing

Evaluation of data accuracy (bias and precision): N.

No specific documentation is currently available. Further consultation with the ICES WGRFS will determine what further processes can be put in place during 2022.

As IMREC_ANG_DI is a self-selecting reporting interface some level of bias is expected. Data will be assessed against data collected through the on-site shore and small boat sampling programmes (IMREC_CREEL) and any bias will be quantified.

Editing and imputation methods: N.

No specific documentation is currently available. Further consultation with the ICES WGRFS will determine what further processes can be put in place during 2022.

If errors are identified, the contributor can be contacted. If the error is recalled, the correct value is inputted. Otherwise, it is removed from the dataset.

Quality document associated to a dataset:

No DOI currently exists for the dataset. No estimation process document is currently available.

Validation of the final dataset:

Data will be rechecked by the database manager prior to submission to end users. If errors are detected, they will be crosschecked with the relevant contributor and corrected if possible. If this is not possible, erroneous data will be removed prior to submission.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

No Deviations

Sampling Design and Protocols

No Deviations

Sampling Implementation

No Deviations

Data Capture

No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations

Data Processing

Although data collection is sufficient, potential bias in the dataset is not well understood yet. Therefore, these data have not been employed to estimate catch. Assessment of bias and development of methods for accounting for bias is ongoing.

Sampling scheme identifier: TUNA_CHART

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: TUNA_CHART
Sampling scheme type: Recreational fisheries
Observation type: SelfAtSea
Time period of validity: 2022-Ongoing
Short description (max 100 words): Tuna CHART is Ireland's multi-agency conventional tagging programme designed to implement, co-ordinate and oversee 'tag and release' of Atlantic Bluefin Tuna (ABFT) by anglers on board authorised Tuna CHART angling charter skipper vessels. The programme commenced in 2019 and is being undertaken to collect spatial and temporal distribution data together with some biological data. In 2018 ICCAT permitted countries in the North-East Atlantic without a Bluefin quota to authorise a limited number of charter vessels to target ABFT to collect these data.
Description of the population
Population targeted: Atlantic Bluefin Tuna taken in Irish waters. Population sampled: All ABFT taken by recreational charter anglers on authorised charter angling vessels are included in the sampling programme. Stratification: Sampling includes all coastlines annually. Angling/sampling is confined to summer and early winter period.
Sampling design and protocols
Sampling design description: Authorised skippers submit a digital report of their bluefin angling trips to Inland Fisheries Ireland (IFI) and Tuna CHART using the Tuna CHART form on Survey123, an ArcGIS application developed by IFI, on ruggedised Samsung tablets. Fields in the digital and associated paper forms (for Marine Institute) were designed to correspond to fields in the

ICCAT conventional tagging document. Skippers are obliged to submit their surveys digitally within a specific timeframe.

ABFT are targeted by anglers on board authorised charter vessels and brought alongside for tagging and measuring in the water. All tagging and relevant biometric data are recorded and all fish are released following a short recovery period.

Skippers must record all ABFT angling trips (effort) and associated catch data. ABFT catch is reported by vessel per trip per day and divided by the number of angling days per vessel per season to provide a seasonal CPUE value.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: N

Link to sampling protocol documentation:

A best practise angling method manual was developed by the Tuna CHART team in 2019 to optimise fish welfare. Manuals for angling, handling/ tagging and data recording are updated annually. Skippers receive training.

ABFT Tagging Manual for the Atlantic-Wide Research Programme for Bluefin Tuna (GBYP - 2010). https://www.iccat.int/GBYP/Docs/Tagging_Manual.pdf

Compliance with international recommendations: Y

The tagging programme was based on guidance in ABFT Tagging Manual for the Atlantic-Wide Research Programme for Bluefin Tuna (GBYP - 2010).
https://www.iccat.int/GBYP/Docs/Tagging_Manual.pdf

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year:

All skippers are obliged to submit electronic reports within a specific timeframe. Skippers

are contacted regularly during the sampling window for progress updates and to identify any sampling issues.

Data capture

Means of data capture:

ABFT capture, tagging and length data are recorded in two formats – electronically on a customized tablet supplied by the Tuna CHART programme, and on waterproof survey sheets. Environmental data are also captured. Electronic data uploads to an IFI geodatabase. Length data are recorded using a measuring tape system provided by IFI.

Data capture documentation:

All skippers are trained and are supplied with SOPs for fish handling and data capture which are reviewed annually.

Quality checks documentation:

N

Data entries are regularly checked and visualized on an ArcGIS operations dashboard. Erroneous data is identified and referred back to the originator for clarification before being approved.

Data storage

National database:

ArcGIS geodatabase and IFI SQL server.

International database: NA

Quality checks and data validation documentation:

Not available at present.

When MS catch reports are submitted to end users (European Commission, ICCAT) standard quality checks will have been undertaken.

Sample storage
Physical samples are not collected for this programme.
Data processing
Evaluation of data accuracy (bias and precision): N. Full catch census undertaken and skippers obliged to report under terms of their authorisation. Editing and imputation methods: N. No specific documentation is currently available. If errors are identified in the database they are removed from the dataset. Quality document associated to a dataset: N https://www.iccat.int/en/accesingdb.html - access to dataset via 'tagging' link Validation of the final dataset: Data will be rechecked by the database manager prior to submission to end users (ICCAT). If potential errors are detected they will be crosschecked with original datasets and corrected if possible. If this is not possible, erroneous data will be removed prior to submission.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan <u>Description of the Population</u> No Deviations <u>Sampling Design and Protocols</u> No Deviations <u>Sampling Implementation</u> No Deviations <u>Data Capture</u> No Deviations

Data Storage

No Deviations

Sample Storage

No Deviations


Data Processing

No Deviations

Quality Reports Related to Table 2.6 and Text Box 2.6**Sampling scheme identifier: IBTS_Q4**

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: IBTS_Q4
Sampling scheme type: Research survey at sea
Observation type: SciObsAtSea (Scientific observer at sea on commercial or scientific vessels)
Time period of validity: 2003 onwards
The main objective of the IBTS_Q4 is to collect data on the distribution, relative abundance and biological parameters of commercial commercially exploited demersal species in 6a south, 7b & 7g-j north. The indices currently utilised by assessment WG's are for haddock, whiting, plaice, cod, hake and sole. Survey data is also provided for white & black anglerfish, megrim, pollack, ling, blue whiting and a number of elasmobranchs as well as several pelagics (herring, horse mackerel and mackerel). Occurrence of vulnerable or sentinel invertebrate species such as corals, sea pen, fan mussel and ocean quahog is also noted. Marine litter is also sorted and recorded. Oceanographic data are collected from CTD instrument on trawl door and occasional surface to sea bed CTD transects. Sediment grabs are carried out opportunistically using a Day grab.
Description of the population
Population targeted: Main target species are haddock (<i>Melanogrammus aeglefinus</i>), whiting (<i>Merlangius merlangus</i>), cod (<i>Gadus morhua</i>), hake (<i>Merluccius merluccius</i>), anglerfish (<i>Lophius piscatorius</i> and <i>L. budegassa</i>) and megrim (<i>Lepidorhombus whiffiagonis</i> and <i>L. boscii</i>), sole (<i>Solea solea</i>), plaice (<i>Pleuronectes platessa</i>).
Population sampled: As a multi-species survey the target ecosystem component is demersal species. No sampling takes place outside the survey area or on grounds that are unsuitable for trawling.
Stratification: The stratification is based on the following considerations:

- Irish Groundfish Survey 2020



Is the sampling design compliant with the 4S principle?: NA (but note that the sampling scheme is statistically sound)

Regional coordination: The survey is also formally coordinated under WGBITS.

240

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%2015%20NeAtl%20IBTS%20Survey.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%2015%20NeAtl%20IBTS%20Survey.pdf)

Compliance with international recommendations: Y

Link to sampling protocol documentation: IBTS SISP 15 manual:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%2015%20NeAtl%20IBTS%20Survey.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%2015%20NeAtl%20IBTS%20Survey.pdf)

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: NA (the survey only takes place once per year).

Data capture

Means of data capture:

The CEFAS software FSS (Fishing Survey System) is used to enter station data and import catch data. These data are stored in a SQL database (FSS_SURVEY) on a local server.

The gear sensor data as well as bottom depth and GPS position are also automatically recorded in a SQL database (FSS_NMEA) at intervals of approximately one per second.

Catch weights, length frequency distributions and biological data are captured using the EFDAQ (Electronic Fisheries Data Acquisition) system and stored in a local database in the wet laboratory before being imported into the central SQL database (FSS_SURVEY).

Data capture documentation:

A new data capture system EFDAQ (Electronic Fisheries Data Acquisition) has been in use in the wet laboratory since 2021. This system was designed by SeaScope Fisheries Research for the Marine Institute and includes hardware such as electronic measuring boards and wands and software application to allow access, collection, visualization, quality assurance and editing of fisheries sample data. Identification and maturity staging are carried out using protocols as recommended by DATRAS and ICES working groups.

Quality checks documentation: Y

EFDAQ Catch Management (20 CatMan V1-7.pdf - not publicly available)

Quality control on sample data (e.g., individual lengths and weights, sample weights etc.) is carried out after every haul using EFDAQ application. Biological age samples such as otoliths are checked against individual fish size before boxes are stored for transport back to laboratory for analysis.

Data storage

National database: FSS (Fishing Survey System)

International database: DATRAS <https://www.ices.dk/data/data-portals/Pages/DATRAS.aspx>

Quality checks and data validation documentation:

Once a survey is complete a number of data checks are carried out on haul positions, gear geometry, catch data and internal consistency of the data. During the upload process to DATRAS a similar range of checks are carried out (<https://www.ices.dk/data/data-portals/Pages/DATRAS.aspx>).

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for analysis. These age structures are generally stored at Marine Institute premises for a period of months before age reading is carried out. Soft tissues are generally collected by request from third parties such as universities and are stored according to protocols provided. Such samples are transported to third parties within weeks of survey completion.

Age reading of IBTS_Q4 otolith samples is carried out according to internationally recognised protocols:

[https://www.ices.dk/sites/pub/Publication%20Reports/Cooperative%20Research%20Report%20\(CRR\)/CRR%20346.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/Cooperative%20Research%20Report%20(CRR)/CRR%20346.pdf)

Data processing

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

No Deviations

Sampling Design and Protocols

No Deviations

Sampling Implementation

Data Capture

No Deviations

Data Storage

No Deviations


Sample Storage

No Deviations

Data Processing


No Deviations

Figure 1. IBTS_Q4 Survey Map with Valid and Invalid stations completed in 2023



Sampling scheme identifier: IBWSS_IRL

MS: IRL

Region: North-East Atlantic
Sampling scheme identifier: IBWSS_IRL
Sampling scheme type: Research survey at sea
Observation type: SciObsAtSea (Scientific observer at sea on commercial or scientific vessels)
Time period of validity: 2004 to present
The main objective of the International blue whiting spawning stock survey is to determine the age stratified abundance and distribution of blue whiting (<i>Micromesistius poutassou</i>) using acoustic survey techniques. Biological data are collected by means of directed trawling on echotraces to determine species composition and biological characteristics of target species. Directed trawling is carried out on echotraces thought to contain mesopelagic fish species as the survey builds capacity towards reporting abundance and distribution of key fish species. Oceanographic data are collected using vertical profiles at pre-determined locations along the survey cruise track. Visual abundance surveys for marine mammals and seabirds are conducted during daylight hours.
Description of the population
Population targeted: The main target species of the survey is blue whiting (<i>Micromesistius poutassou</i>)
Population sampled: Blue whiting are targeted within a pre-defined survey boundary region, containing the core spawning grounds.
Stratification: The geographical survey area is stratified based on two key criteria; acoustic sampling effort within the stratum and scaled historic abundance (core or peripheral stratum).

Figure 1. IBWSS Survey area stratification (numbered boxes). Strata 1-3 core abundance and

4-7 peripheral, low abundance
Sampling design and protocols
Sampling design description: PSU is measured in 1 nmi (nautical mile) EDSU (Elementary distance sampling units).
Is the sampling design compliant with the 4S principle? NA
Regional coordination: IBWSS is coordinated through ICES WGIPS.
Link to sampling design documentation: The Manual for International Pelagic Surveys (IPS) describes the methods used for survey design, analysis and reporting of survey data (SISP #9) and is described in detail in the latest IBWSS survey report (http://hdl.handle.net/10793/1689)
Compliance with international recommendations: Y
Link to sampling protocol documentation: The Manual for International Pelagic Surveys (IPS) describes the methods used for survey design, analysis and reporting of survey data (SISP #9) and is described in detail in the latest IBWSS survey report (http://hdl.handle.net/10793/1689)
Compliance with international recommendations: Y
Sampling implementation
Recording of refusal rate: NA
Monitoring of sampling progress within the sampling year: NA (Annual survey)
Data capture
Means of data capture: Acoustic data are recorded via a Simrad EK60 scientific echosounder and processed using a proprietary software (Echoview V12). Biological data are collected and stored within an SQLite database and held nationally. Aggregated acoustic and biological data are uploaded to the open access ICES Trawl Acoustic repository post survey (https://www.ices.dk/data/data-portals/Pages/acoustic.aspx).
Data capture documentation: The Manual for International Pelagic Surveys (IPS) describes the methods used for survey design, analysis and reporting of survey data (SISP #9)
Quality checks documentation: Y (The Manual for International Pelagic Surveys (IPS) describes the methods used for survey design, analysis and reporting of survey data (SISP #9))
Data storage
National database: Acoustic data repository with data stored separately for each survey/year
International database: ICES acoustic trawl survey database https://www.ices.dk/data/data-portals/Pages/acoustic.aspx
Quality checks and data validation documentation: Data undergo checks and validation during submission to ICES. The ICES controlled vocabularies can be found at http://vocab.ices.dk/?theme=4

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

Biological samples (otoliths) are aged onboard the ship (herring & blue whiting) for species requiring additional processing prior to age reading (horse mackerel, boarfish & mackerel) samples are dry stored for transportation to the Marine Institute.

Age reading of IBWSS samples is carried out according to internationally recognised protocols:

[https://www.ices.dk/sites/pub/Publication%20Reports/Cooperative%20Research%20Report%20\(CRR\)/CRR%20346.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/Cooperative%20Research%20Report%20(CRR)/CRR%20346.pdf)

Data processing

Evaluation of data accuracy (bias and precision): Acoustic biomass and abundance from survey data is calculated using the open source software StoX (<https://doi.org/10.1111/2041-210X.13250>). Within StoX, the RStoX package has been developed to calculate the coefficient of variation (CV) of survey estimates. CV across the survey time series is described in the latest IBWSS survey report (<http://hdl.handle.net/10793/1689>)

Editing and imputation methods: Y within the StoX analysis framework. Survey estimates are reviewed annually at the survey coordination group ICES WGIPS.

Quality document associated to a dataset: The publishing of DOIs relating to survey data uploaded to the ICES data portal is under development and will be implemented as part of the Transparent Assessment Framework within ICES (<http://ices.dk/marine-data/assessment-tools/Pages/transparent-assessment-framework.aspx>)

Validation of the final dataset: Data upload to the ICES portal (<https://www.ices.dk/data/data-portals/Pages/acoustic.aspx>) is dependent on meeting defined metadata standards described in the vocabulary (<http://vocab.ices.dk/?theme=4>)

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

No Deviation

Sampling Design and Protocols

No Deviation

Sampling Implementation

Due to equipment failure on the Celtic Explorer, fishing trawls were not possible for the duration of the Irish leg of the International Blue Whiting Acoustic Survey 2023. Acoustic data from the survey was combined with acoustic data from the other nations taking part in this internationally coordinated survey as per standard protocols. Biological data from the nearest hauls taken by the other vessels were used for the calculation of the age-stratified blue whiting biomass for the entire area. There was therefore no impact on the quality of the combined spawning stock biomass index.

Data Capture

No Deviation

Data Storage

No Deviation


Sample Storage

No Deviation

Data Processing

No Deviation

Figure 2. IBWSS Survey Map 2023



Sampling scheme identifier: MEGS_IRL

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: MEGS_IRL
Sampling scheme type: Research survey at sea
Observation type: SciObsAtSea (Scientific observer at sea on commercial or scientific vessels)
Time period of validity: 1992 onwards
The main objective of the MEGS_IRL sampling is to provide egg counts and histology data for mackerel (<i>Scomber scombrus</i>) and horse mackerel (<i>Trachurus trachurus</i>), as well as CTD data, to let WGMEGS calculate an SSB for North-east Atlantic mackerel, and an egg production estimate for horse mackerel. Eggs are collected, identified and staged from plankton tows carried out using a GULF VII plankton sampler every ICES half statistical rectangle in ICES areas 5, 6, 7 and 12. Histology samples are collected from opportunistic fishing hauls. Oceanographic data are collected from every plankton station using a CTD mounted on the GULF frame. Secondary objectives are to collect egg count data of other species, such as hake (<i>Merluccius merluccius</i>).
Description of the population
Population targeted: The main target species are mackerel (<i>Scomber scombrus</i>) and horse mackerel (<i>Trachurus trachurus</i>). Population sampled: The main species targeted from the plankton sampling are mackerel (<i>Scomber scombrus</i>) and horse mackerel (<i>Trachurus trachurus</i>). Secondary species identified include hake (<i>Merluccius merluccius</i>) and ling (<i>Molva Molva</i>). Stratification: The stratification is based on the following considerations: • Plankton samples are collected from every ICES half statistical rectangle, along a series of transects, in the survey area allocated to Ireland by WGMEGS. (Fig 1) This location can vary from survey to survey. The survey is adaptive so the decision on when to finish a transect is at the discretion of the scientist-in-charge. • Histology samples, as well as length, weight and maturity data, and otoliths, are collected from female mackerel and horse mackerel caught in opportunistic trawl

hauls.




Fig 1: Plankton station locations for MEGS_IRL for both March and June surveys 2022

Sampling design and protocols

Sampling design description: Individual hauls are the PSU. Plankton samples are collected from each ICES half statistical rectangle, along a series of transects. Histology samples, as well as associated biological data are collected from female mackerel and horse mackerel caught in opportunistic trawl hauls. Plankton and histology samples are processed according to the MEGS SISP 5 and 6 manuals.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: MEGS_IE is formally coordinated under WGMEGS.

Link to sampling design documentation: ICES SISP 5 and 6 manuals:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%205%20-%20WGMEGS%20Manual%20for%20AEPM%20and%20DEPM.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%205%20-%20WGMEGS%20Manual%20for%20AEPM%20and%20DEPM.pdf)

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%206%20Manual%20for%20the%20mackerel%20and%20horse%20mackerel%20egg%20surveys,%20smapping%20at%20sea_Jan%202019.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%206%20Manual%20for%20the%20mackerel%20and%20horse%20mackerel%20egg%20surveys,%20smapping%20at%20sea_Jan%202019.pdf)

Compliance with international recommendations: Y

Link to sampling protocol documentation: ICES SISP 5 and 6 manuals:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%206%20Manual%20for%20the%20mackerel%20and%20horse%20mackerel%20egg%20surveys,%20smapping%20at%20sea_Jan%202019.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%206%20Manual%20for%20the%20mackerel%20and%20horse%20mackerel%20egg%20surveys,%20smapping%20at%20sea_Jan%202019.pdf)

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%205%20-%20WGMEGS%20Manual%20for%20AEPM%20and%20DEPM.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%205%20-%20WGMEGS%20Manual%20for%20AEPM%20and%20DEPM.pdf)

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: NA

Data capture

Means of data capture:

Egg counts and fishing biological data are initially recorded on paper and then transferred to spreadsheet as soon as possible after sampling. A database to hold this data is currently under development. CTD data is collected and stored electronically.

Data capture documentation:

Plankton and histology data collected on survey are compiled into spreadsheets designed by WGMEGS. Egg identification and maturity staging are carried out using protocols as recommended by WGMEGS.

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%205%20-%20WGMEGS%20Manual%20for%20AEPM%20and%20DEPM.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%205%20-%20WGMEGS%20Manual%20for%20AEPM%20and%20DEPM.pdf)

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%206%20Manual%20for%20the%20mackerel%20and%20horse%20mackerel%20egg%20surveys,%20smapping%20at%20sea_Jan%202019.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%206%20Manual%20for%20the%20mackerel%20and%20horse%20mackerel%20egg%20surveys,%20smapping%20at%20sea_Jan%202019.pdf)

Quality checks documentation: Y Survey data is compiled by a WGMEGS coordinator. Total survey data is compiled from all Institutes and is run through an R script https://github.com/GersomCostas/Teggprod to check for errors prior to being used to provide the SSB assessment. Biological age samples such as otoliths are checked against individual fish size before boxes are stored for transport back to laboratory for analysis.
Data storage
National database: FEAS_MEGS (in development) International database: ICES Egg and larval database; https://www.ices.dk/data/data-portals/Pages/Eggs-and-larvae.aspx Quality checks and data validation documentation: Once a survey is complete several data checks are carried out on haul positions and CTD data. During the upload process to the ICES egg and larval database a similar range of checks are carried out. The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).
Sample storage
Plankton samples and extracted eggs are stored onsite in the Marine Institute in 4% buffered formalin. Histology samples are kept for 18 months until the data has been accepted and validated by WGMEGS. These histology samples are then disposed of. Age reading is of MEGS samples is carried out according to internationally recognised protocols: https://www.ices.dk/sites/pub/Publication%20Reports/Cooperative%20Research%20Report%20(CRR)/CRR%20346.pdf
Data processing
Evaluation of data accuracy (bias and precision): N/Y There is no procedure in place to estimate bias. Interim results are provided to the relevant stock assessment working group (WGWIDE) in the year of the survey. Final results, including precision of abundance and biomass estimates for main target species, are provided in the WGMEGS report published the year

after the survey has taken place, and are subsequently presented to WGWIDE.

<https://www.ices.dk/sites/pub/Publication%20Reports/Forms/DispForm.aspx?ID=37365>

Editing and imputation methods: NA

Quality document associated to a dataset:

No DOI is currently created for the dataset. The data is uploaded to the ICES database the year following the survey and is publicly available from there.

Validation of the final dataset:

Datasets are validated prior to upload to the egg and larval database. This ICES database also has additional validation checks which are applied during data upload.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Description of the Population

No MEGS Survey in 2023, next scheduled for 2025

Sampling Design and Protocols

No MEGS Survey in 2023, next scheduled for 2025

Sampling Implementation

No MEGS Survey in 2023, next scheduled for 2025

Data Capture

No MEGS Survey in 2023, next scheduled for 2025

Data Storage

No MEGS Survey in 2023, next scheduled for 2025

Sample Storage

No MEGS Survey in 2023, next scheduled for 2025

Data Processing

No MEGS Survey in 2023, next scheduled for 2025

Sampling scheme identifier: WESPAS_IRL

MS: IRL

Region: North-East Atlantic

Sampling scheme identifier: WESPAS_IRL

Sampling scheme type: Research survey at sea

Observation type: SciObsAtSea (Scientific observer at sea on commercial or scientific

vessels)

Time period of validity: 2011 to present

The primary aim of the WESPAS survey is to determine the age stratified abundance and distribution of herring (*Clupea harengus*), boarfish (*Capros aper*) and horse mackerel (*Trachurus trachurus*) using acoustic survey techniques. Biological data are collected by means of directed trawling on echotracers to determine species composition and biological characteristics of target species. Oceanographic data are collected using vertical profiles at pre-determined locations along the survey cruise track. Zooplankton sampling is conducted at hydrographic station and used to determine the dry weight biomass across the survey area. Visual abundance surveys for marine mammals and seabirds are conducted during daylight hours.

Description of the population

Population targeted: The main target species of the survey are herring (*Clupea harengus*), boarfish (*Capros aper*) and horse mackerel (*Trachurus trachurus*).

Population sampled: Target species are sampled on the summer feeding grounds (herring) and spawning grounds (boarfish and horse mackerel).

Stratification: The geographical survey area is stratified based on two key criteria; acoustic sampling effort within the stratum and scaled historic abundance (core or peripheral stratum).




Figure 1. WESPAS Survey area stratification and trawl stations (grey circles) 2021.

Sampling design and protocols

Sampling design description: PSU is measured in 1 nmi (nautical mile) EDSU (Elementary distance sampling units).

Is the sampling design compliant with the 4S principle? NA

Regional coordination: WESPAS is coordinated through ICES WGIPS. Link to sampling design documentation: The Manual for International Pelagic Surveys (IPS) describes the methods used for survey design, analysis and reporting of survey data (<u>SISP #9</u>) and is described in detail in the latest WESPAS survey report (http://hdl.handle.net/10793/1659) Compliance with international recommendations: Y Link to sampling protocol documentation: The Manual for International Pelagic Surveys (IPS) describes the methods used for survey design, analysis and reporting of survey data (<u>SISP #9</u>) and is described in detail in the latest WESPAS survey report (http://hdl.handle.net/10793/1659) Compliance with international recommendations: Y Sampling implementation Recording of refusal rate: NA Monitoring of sampling progress within the sampling year: NA (Annual survey) Data capture Means of data capture: Acoustic data are recorded via a Simrad EK60 scientific echosounder and processed using a proprietary software (Echoview V12). Biological data are collected and stored within a SQLite database and held nationally. Aggregated acoustic and biological data are uploaded to the open access ICES Trawl Acoustic repository post survey (https://www.ices.dk/data/data-portals/Pages/acoustic.aspx). Data capture documentation: The Manual for International Pelagic Surveys (IPS) describes the methods used for survey design, analysis and reporting of survey data (<u>SISP #9</u>) Quality checks documentation: Y (The Manual for International Pelagic Surveys (IPS) describes the methods used for survey design, analysis and reporting of survey data (<u>SISP #9</u>))
Data storage National database: Acoustic data repository with data stored separately for each survey/year International database: ICES acoustic trawl survey database https://www.ices.dk/data/data-portals/Pages/acoustic.aspx Quality checks and data validation documentation: Data undergo checks and validation during submission to ICES. The ICES controlled vocabularies can be found at http://vocab.ices.dk/?theme=4 The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF). Sample storage

Biological samples (otoliths) are aged onboard the ship (herring & blue whiting) for species requiring additional processing prior to age reading (horse mackerel, boarfish & mackerel) samples are dry stored for transportation to the Marine Institute.

Age reading of WESPAS samples is carried out according to internationally recognised protocols:

[https://www.ices.dk/sites/pub/Publication%20Reports/Cooperative%20Research%20Report%20\(CRR\)/CRR%20346.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/Cooperative%20Research%20Report%20(CRR)/CRR%20346.pdf)

Data processing

Evaluation of data accuracy (bias and precision): Acoustic biomass and abundance from survey data is calculated using the open-source software StoX (<https://doi.org/10.1111/2041-210X.13250>). Within StoX, the RStoX package has been developed to calculate the coefficient of variation (CV) of survey estimates. CV across the survey time series is described in the latest IBWSS survey report (<http://hdl.handle.net/10793/1659>)

Editing and imputation methods: Y within the StoX analysis framework. Survey estimates are reviewed annually at the survey coordination group ICES WGIPS.

Quality document associated to a dataset: The publishing of DOIs relating to survey data uploaded to the ICES data portal is under development and will be implemented as part of the Transparent Assessment Framework within ICES (<http://ices.dk/marine-data/assessment-tools/Pages/transparent-assessment-framework.aspx>)

Validation of the final dataset: Data upload to the ICES portal (<https://www.ices.dk/data/data-portals/Pages/acoustic.aspx>) is dependent on meeting defined metadata standards described in the vocabulary (<http://vocab.ices.dk/?theme=4>)

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

No Deviation

Sampling Design and Protocols

No Deviation

Sampling Implementation

No Deviation

Data Capture

No Deviation

Data Storage

No Deviation

Sample Storage

No Deviation

Data Processing


No Deviation

Figure 3. WESPAS Survey Map 2023

Sampling scheme identifier: *Nephrops* UWTV Survey (UWTV 16 – 17, UWTV 19, UWTV 20 – 22)

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: <i>Nephrops</i> UWTV Survey (UWTV16-17, UWTV19,

UWTV20-22)
Sampling scheme type: Research survey at sea
Observation type: SciObsAtSea (Scientific observer at sea on commercial or scientific vessels)
Time period of validity: 2002 – 2027
The main objective of the <i>Nephrops</i> UWTV Survey sampling scheme is to obtain quality assured burrow abundance estimates for Norway lobster (<i>Nephrops norvegicus</i>) in Functional Units (FU): 16, 17, 19, 20-21 combined, and 22 in area 7. Secondary objectives are to record observations of trawl marks, fish and <i>Nephrops</i> activity. Occurrence of vulnerable or sentinel invertebrate species such as soft corals and sea pens is also noted. Marine litter is recorded. Oceanographic data are collected from a sledge mounted CTD instrument. Beam trawl tows are carried out on FU 17 “Aran” and FU 22 “Smalls” grounds only when UWTV operations have been fully achieved. Sediment grabs are carried out opportunistically using a Day grab and contribute to the Irish national “INFOMAR” seabed mapping programme and are used during ICES FU assessment Benchmark processes by contributing to the definition of the spatial area of FU <i>Nephrops</i> grounds.
Description of the population
Population targeted: Main target species is Norway lobster (<i>Nephrops norvegicus</i>). Population sampled: Main target species is Norway lobster (<i>Nephrops norvegicus</i>). No sampling takes place outside survey areas. Stratification: The stratification is based on the following considerations: - Clearly defined <i>Nephrops</i> grounds (see map below) were identified as separate strata; an area defined by sediment data and high fishing intensity surrounded by low fishing intensity signify that the bottom type and ecology on the fishing ground is different from that of the surrounding area. <i>Nephrops</i> FUs are treated separately because they comprise different assessment areas.



Sampling design and protocols

Sampling design description: Individual video transects are the Primary sample Unit (PSU), these are selected from random locations inside each stratum. Each transect is then processed according to the ICES Manual for *Nephrops* Underwater TV Surveys TIMES 65 manual <https://doi.org/10.17895/ices.pub.8014>.

Is the sampling design compliant with the 4S principle?: NA (but note that the sampling scheme is statistically sound)

Regional coordination: NA. The survey is formally coordinated under ICES WGNPS. International staff exchange is facilitated when possible on UWTV surveys to allow for protocol and technical expertise development and international standardisation.

Link to sampling protocol documentation: ICES Manual for *Nephrops* Underwater TV Surveys TIMES 65: [https://www.ices.dk/sites/pub/Publication%20Reports/Techniques%20in%20Marine%20Environmental%20Sciences%20\(TIMES\)/TIMES%2065.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/Techniques%20in%20Marine%20Environmental%20Sciences%20(TIMES)/TIMES%2065.pdf)

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: NA (the survey only takes place once per year).

Data capture
Means of data capture: Video transect metadata data are stored in a local database in the research vessel dry laboratory before being imported into the central SQL database (UWTV_Surveys) on return to shore. The GPS and USBL position and depth are automatically recorded at intervals of approximately one every three seconds and stored in a local database in the research vessel dry laboratory before being imported into the central SQL database (UWTV_Surveys) on return to shore. Burrow counts and presence/absence data are captured using an HD Image annotation “R-Shiny” app system and stored in a local database in the research vessel dry laboratory before being imported into the central SQL database (UWTV_Surveys) on return to shore. (Note that owing to the Covid-19 pandemic and restrictions on the numbers of scientists aboard research vessels, in 2020 and 2021 it was necessary to undertake this operation on-shore. It is the intention of the programme to return these operations to being undertaken at sea during surveys). Following beam trawl tows (carried out on FU 17 “Aran” and FU 22 “Smalls” grounds – see above), <i>Nephrops</i> catch weights, length frequency distributions and biological data are captured using digital electronic callipers and marine scales connected to a tablet and stored in a local database in the research vessel wet laboratory, before being imported into the central SQL database (FSS_SURVEY) on return to shore. Fish (weights only) and benthic catch data (weights and counts) are stored on paper copy and transferred to electronic spreadsheets before uploaded to the survey network. Data capture documentation: A new data capture system HD Image annotation “R-Shiny” app has been in use since 2019. This system was designed in-house by the Marine Institute (Aristegui, M., 2020) and allows access, collection, visualization, quality assurance and editing of image and position data. Burrow identification and technical processes are carried out using protocols as recommended by ICES WGNPS. Quality checks documentation: Y Quality control on position data is carried out after every video transect using in-house developed scripts. Burrow count data are verified according to international standards.
Data storage
National database: FEAS_UWTV_Surveys All data are maintained on secure servers. International database: International database is under development.

Quality checks and data validation documentation:

Once a survey is complete a number of data checks are carried out on transect positions, count data and internal consistency of the data.

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

No biological samples are stored.

Nephrops burrow identification and counting is carried out according to internationally recognised protocols:

[https://www.ices.dk/sites/pub/Publication%20Reports/Cooperative%20Research%20Report%20\(CRR\)/CRR340.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/Cooperative%20Research%20Report%20(CRR)/CRR340.pdf)

<https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/EOSG/2018/WKNEPS/WKNEPS%20report%202018.pdf>

Data processing**Evaluation of data accuracy (bias and precision):** N/Y

There is no procedure in place to estimate bias.

Precision of abundance estimates for the main target species are provided in Table 1 of the annual survey reports and are provided to the relevant stock assessment working group (ICES WGCSE and WGNEPS). The target level of precision (an overall coefficient of variance or standard error of less than 20%) is the international standard.

Editing and imputation methods: NA**Quality document associated to a dataset:**

Procedure for producing the estimations of abundance for the target species is included in annual survey reports. No DOI is created.

Annual survey reports are available for each *Nephrops* FU:

FU 16: <http://hdl.handle.net/10793/1655>

FU 17: <http://hdl.handle.net/10793/1656>

FU19: <http://hdl.handle.net/10793/1654>

FU 20-21 combined: <http://hdl.handle.net/10793/1657>

FU 22: <http://hdl.handle.net/10793/1658>

Validation of the final dataset:

Marine Institute *Nephrops* UWTV survey data and products are included in the Data

Management Quality Management Framework (DM-QMF) by the (UNESCO) International Oceanographic Commissions (IODE) - International Oceanographic Data and Information Exchange programme framework since 2019.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

No deviation.

Sampling Design and Protocols

No deviation.

Sampling Implementation

In 2023 additional video transect sampling was carried out on the FU 15 western Irish Sea *Nephrops* grounds in collaboration with AFBI – Belfast, Northern Ireland.

Data Capture

No deviation.

Data Storage

No deviation.


Sample Storage

No biological samples are stored

Data Processing

No deviation.

Figure 4. UWTV Survey Map 2023 (All Functional Units).



Sampling scheme identifier: CSHAS_IRL

MS: IRL
Region: North-East Atlantic
Sampling scheme identifier: CSHAS_IRL
Sampling scheme type: Research survey at sea
Observation type: SciObsAtSea (Scientific observer at sea on commercial or scientific vessels)
Time period of validity: 2004 to present
The primary aim of the CSHAS survey is to determine the age stratified abundance and distribution of herring (<i>Clupea harengus</i>) and sprat (<i>Sprattus sprattus</i>) using acoustic survey techniques. Biological data are collected by means of directed trawling on echotracers to determine species composition and biological characteristics of target species. Oceanographic data are collected using vertical profiles at pre-determined locations along the survey cruise track. Zooplankton sampling is conducted at hydrographic station and used to determine the dry weight biomass across the survey area. Visual abundance surveys for marine mammals and seabirds are conducted during daylight hours.
Description of the population
Population targeted: The main target species of the survey are herring (<i>Clupea harengus</i>) and sprat (<i>Sprattus sprattus</i>).

Population sampled: Target species are sampled on the spawning/pre-spawning grounds (herring) and feeding grounds (sprat).

Stratification: The geographical survey area is stratified based on two key criteria; acoustic sampling effort within the stratum and scaled historic abundance (core or peripheral stratum).




Figure 1. CSHAS_IRL Survey area stratification (Pass 1 & Pass 2) and trawl stations (blue circles) 2021.

Sampling design and protocols

Sampling design description: PSU is measured in 1 nmi (nautical mile) EDSU (Elementary distance sampling units).

Is the sampling design compliant with the 4S principle? NA

Regional coordination: CSHAS_IRL is coordinated through ICES WGIPS.

Link to sampling design documentation: The Manual for International Pelagic Surveys (IPS) describes the methods used for survey design, analysis and reporting of survey data (SISP #9) and is described in detail in the latest CSHAS survey report (<http://hdl.handle.net/10793/1664>).

Compliance with international recommendations: Y

Link to sampling protocol documentation: The Manual for International Pelagic Surveys (IPS) describes the methods used for survey design, analysis and reporting of survey data (SISP #9) and is described in detail in the latest CSHAS survey report (<http://hdl.handle.net/10793/1664>).

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: NA (Annual survey)

Data capture
Means of data capture: Acoustic data are recorded via a Simrad EK60 scientific echosounder and processed using a proprietary software (Echoview V12). Biological data are collected and stored within a SQLite database and held nationally. Aggregated acoustic and biological data are uploaded to the open access ICES Trawl Acoustic repository post survey (https://www.ices.dk/data/data-portals/Pages/acoustic.aspx). Data capture documentation: The Manual for International Pelagic Surveys (IPS) describes the methods used for survey design, analysis and reporting of survey data (SISP #9) Quality checks documentation: Y (The Manual for International Pelagic Surveys (IPS) describes the methods used for survey design, analysis and reporting of survey data (SISP #9))
Data storage
National database: Acoustic data repository with data stored separately for each survey/year International database: ICES acoustic trawl survey database https://www.ices.dk/data/data-portals/Pages/acoustic.aspx Quality checks and data validation documentation: Data undergo checks and validation during submission to ICES. The ICES controlled vocabularies can be found at http://vocab.ices.dk/?theme=4 The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).
Sample storage
Biological samples (otoliths) are aged onboard the ship (herring & blue whiting) for species requiring additional processing prior to age reading (horse mackerel, boarfish & mackerel) samples are dry stored for transportation to the Marine Institute. Age reading of CSHAS samples is carried out according to internationally recognised protocols: https://www.ices.dk/sites/pub/Publication%20Reports/Cooperative%20Research%20Report%20(CRR)/CRR%20346.pdf
Data processing
Evaluation of data accuracy (bias and precision): Acoustic biomass and abundance from survey data is calculated using the open-source software StoX (https://doi.org/10.1111/2041-210X.13250). Within StoX, the RStoX package has been developed to calculate the coefficient of variation (CV) of survey estimates. CV across the survey time series is described in the latest IBWSS survey report (http://hdl.handle.net/10793/1664). Editing and imputation methods: Y within the StoX analysis framework. Survey estimates are reviewed annually at the survey coordination group ICES WGIPS. Quality document associated to a dataset: The publishing of DOIs relating to survey data uploaded to the ICES data portal is under development and will be implemented as part of the Transparent Assessment Framework within ICES (http://ices.dk/marine-data/assessment-

[tools/Pages/transparent-assessment-framework.aspx](#))

Validation of the final dataset: Data upload to the ICES portal (<https://www.ices.dk/data/data-portals/Pages/acoustic.aspx>) is dependent on meeting defined metadata standards described in the vocabulary (<http://vocab.ices.dk/?theme=4>)

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

No deviation.

Sampling Design and Protocols

No deviation.

Sampling Implementation

No deviation.

Data Capture

No deviation.

Data Storage

No deviation.


Sample Storage

No biological samples are stored

Data Processing

No deviation.

Figure 5. CSHAS Survey Map 2023 showing achieved survey track and hauls.



Sampling scheme identifier: IAMS_ IRL

MS : IRL
Region: North-East Atlantic
Sampling scheme identifier: IAMS_ IRL
Sampling scheme type: Research survey at sea
Observation type: SciObsAtSea (Scientific observer at sea on commercial or scientific vessels)
Time period of validity: 2016 onwards
The main objective of the IAMS_ IRL sampling scheme is to obtain biomass and abundance indices for anglerfish (<i>Lophius piscatorius</i> and <i>L. budegassa</i>) and megrim (<i>Lepidorhombus whiffiagonis</i> and <i>L. boscii</i>) in areas 6a (south of 58°N) and 7 (west of 8°W). Secondary objectives are to collect data on the distribution, relative abundance and biology of other commercially exploited species. Occurrence of vulnerable or sentinel invertebrate species such as corals, sea pen, fan mussel and ocean quahog is also noted. Marine litter is also sorted and recorded. Oceanographic data are collected from CTD instrument on trawl door and occasional surface to seabed CTD transects. Sediment grabs are carried out opportunistically using a Day grab.

Description of the population					
Population targeted:	Main target species are anglerfish and megrim (<i>Lophius piscatorius</i> and <i>L. budegassa</i>) (<i>Lepidorhombus whiffiagonis</i> and <i>L. boscii</i>).				
Population sampled:	Main target species are anglerfish and megrim (<i>Lophius piscatorius</i> and <i>L. budegassa</i>) (<i>Lepidorhombus whiffiagonis</i> and <i>L. boscii</i>). No sampling takes place outside the survey area or on grounds that are unsuitable for trawling.				
Stratification: The stratification is based on the following considerations: • Depth: 0-200m; 200-500m; and 500-1,000m • Clearly defined fishing grounds were identified as separate strata; an area with high fishing intensity surrounded by low fishing intensity signify that the bottom type and ecology on the fishing ground is different from that of the surrounding area. • Catch rates of the target species were also taken into account in determining the boundaries of the strata. • Rocky bottom types are excluded from the survey area. • Regions 6a and 7bcjk are treated separately because they comprise different assessment and TAC areas.					
Sampling design and protocols					
Sampling design description: Individual hauls are the PSU, these are selected from random locations inside each stratum. The catch is then processed according to the IBTS SISP 15 manual.					
Is the sampling design compliant with the 4S principle?: NA (but note that the sampling scheme is statistically sound)					
Regional coordination: IAMS_IE is informally coordinated with the Scottish Anglerfish and Megrim Survey (SIAMISS). The survey is also formally coordinated under WGBITS.					
Link to sampling design documentation: IBTS SISP 15 manual: https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%2015%20NeAtl%20IBTS%20Survey.pdf					
Compliance with international recommendations: Y					
Link to sampling protocol documentation: IBTS SISP 15 manual: https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%2015%20NeAtl%20IBTS%20Survey.pdf					
Compliance with international recommendations: Y					
Sampling implementation					
Recording of refusal rate: NA					
Monitoring of sampling progress within the sampling year: NA (the survey only takes place once per year).					

Data capture	
Means of data capture: The CEFAS software FSS (Fishing Survey System) is used to enter station data and import catch data. These data are stored in a SQL database (FSS_SURVEY) on a local server. The gear sensor data as well as bottom depth and GPS position are also automatically recorded in a SQL database (FSS_NMEA) at intervals of approximately one per second. Catch weights, length frequency distributions and biological data are captured using the EFDAQ (Electronic Fisheries Data Acquisition) system and stored in a local database in wet laboratory before being imported into the central SQL database (FSS_SURVEY). Data capture documentation: A new data capture system EFDAQ (Electronic Fisheries Data Acquisition) has been in use in the wet laboratory since 2021. This system was designed by SeaScope Fisheries Research for the Marine Institute and includes hardware such as electronic measuring boards and wands and software application to allow access, collection, visualization, quality assurance and editing of fisheries sample data. Identification and maturity staging are carried out using protocols as recommended by DATRAS and ICES working groups. Quality checks documentation: Y EFDAQ Catch Management (20 CatMan V1-7.pdf - not publicly available) Quality control on sample data (e.g. individual lengths and weights, sample weights etc.) is carried out after every haul using EFDAQ application. Biological age samples such as otoliths are checked against individual fish size before boxes are stored for transport back to laboratory for analysis.	
Data storage	
National database: FSS (Fishing Survey System) International database: DATRAS https://www.ices.dk/data/data-portals/Pages/DATRAS.aspx Quality checks and data validation documentation: Once a survey is complete a number of data checks are carried out on haul positions, gear geometry, catch data and internal consistency of the data. During the upload process to DATRAS a similar range of checks are carried out (https://www.ices.dk/data/data-portals/Pages/DATRAS.aspx). The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).	
Sample storage	

Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for analysis. These age structures are generally stored at Marine Institute premises for a period of months before age reading is carried out. Soft tissues are generally collected by request from third parties such as universities and are stored according to protocols provided. Such samples are transported to third parties within weeks of survey completion.

Age reading of IAMS samples is carried out according to internationally recognised protocols:

[https://www.ices.dk/sites/pub/Publication%20Reports/Cooperative%20Research%20Report%20\(CRR\)/CRR%20346.pdf](https://www.ices.dk/sites/pub/Publication%20Reports/Cooperative%20Research%20Report%20(CRR)/CRR%20346.pdf)

Data processing

Evaluation of data accuracy (bias and precision): N/Y

There is no procedure in place to estimate bias.

Precision of abundance and biomass estimates for main target species are provided in Table 6 of the annual survey report: e.g. <https://oar.marine.ie/handle/10793/1691> and are provided to the relevant stock assessment working group (WGBIE).

Editing and imputation methods: NA – no imputation takes place (with the exception of gear parameters that could not be observed – these are imputed using a model based on observed values).

Quality document associated to a dataset:

Procedure for producing the estimations of abundance and biomass for main species is included in annual survey report (<https://oar.marine.ie/handle/10793/1691>). No DOI is created.

Validation of the final dataset: Datasets are validated prior to DATRAS upload. For more information see: <https://datras.ices.dk/Data%20submission/Default.aspx>

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

No deviation.

Sampling Design and Protocols

No deviation.

Sampling Implementation

No deviation.

Data Capture

No deviation.

Data Storage

No deviation.

Sample Storage

No biological samples are stored

Data Processing

No deviation.

Figure 6. IAMS Survey Map 2023 showing achieved hauls.

Sampling scheme identifier: RCIS_IRL

MS : IRL


Region : North-East Atlantic

Sampling scheme identifier : RCIS_IRL


Sampling scheme type: Research Survey at sea
Observation type: SciObsAtSea
Time period of validity : 2017 onwards
The main objective of the Razor Clam Irish Sea (RCIS_IRL) surveys is to estimate the biomass of razor clams and to provide advice for <i>Ensis siliqua</i> in Area 7a. Secondary objectives are to collect data on the distribution and abundance of other bivalve species caught as by-catch during the surveys.
Description of the population
Population targeted: The swept area of all dredge hauls for the main target species, <i>Ensis siliqua</i> in the northern and southern Irish Sea Population sampled: 800 dredge hauls are sampled throughout the extent of the razor clam beds in the North Irish Sea and approximately 120 dredge hauls in the southern Irish Sea in water depths of 4-14 meters. Stratification: The North and South Irish Sea surveys follow a design which uses an iVMS grid to allocate survey effort. Inshore VMS activity is seen as a proxy for the abundance of razor clams. Four iVMS effort strata of the same surface area were defined and 200 stations were randomly assigned within each strata, to ensure an even distribution of randomly assigned grid cells across the range of iVMS effort.



iVMS data showing distribution of Razor Clam fishery in the north Irish Sea inside 10m depth contour and the extent of the survey area of the 2021 North Irish Sea Razor clam survey.



iVMS data showing distribution of Razor Clam fishery in the South Irish Sea



Extent of the survey area of the 2020 South Irish Sea razor clam surveys off Curracloe and Rosslare.

Sampling design and protocols

Sampling design description: The PSU is dredge haul. The location of each dredge haul is selected randomly within four iVMS effort strata. Biomass of each dredge haul is estimated as the product of density and mean individual weight calculated from the size distribution at the station and a weight-length relationship. The following protocols are followed:

If total catch is of manageable size sort the catch completely. All bivalves are kept and sorted by species. Record weight and/or count of all Razor clams. Record the count of all other bivalves. If quantities of by-catch bivalves are too high subsample the catch by discarding a portion of the catch once all razors are removed (i.e. 1/2 or 1/3). Record the portion discarded and count all bivalves by species. If more subsampling is required because of high numbers of certain species record total weight of the sample and then the subsample weight and count of the bivalves in question. Measure all or a subsample of Razor clams (If only measuring a sub-sample of razor clams record a weight of the sub-sample).

If total catch/bulk is not of manageable size there are two options.

1) In order to adequately sample large grade razors which may be present in low numbers but which contribute a lot to overall weight of razors in the catch sort through the catch/bulk retaining all **large** commercial sized razor clams. Record the weight and /or number of these commercial razor clam and measure all or a subsample of them. If only measuring a subsample record the count of the full sample and the weight of the subsample. Level the remaining catch/bulk on the table and discard a portion of it (i.e. 1/2 or 1/3). Record proportion discarded. Sort through the remaining removing all bivalves. Record weight (or count if there are no scales on board) of all razor clams. Measure all/subsample of these razor clams. Record count of all other bivalves. If quantities of other bivalves are too high, subsample the catch again (i.e. 1/2 or 1/3). Record the proportion discarded and then sort the bivalves and count the number of each retained. If quantities of some by-caught bivalve species are still very high (i.e. *Pharus legumen* and *Acanthocardia sp.*), then sub-sample them by recording the weight of the sample and then record the weight and count of the subsample. Measure a sample/subsample of the other razor clam.

2) Where there is no evidence that large razors are present in low numbers Level the catch/bulk on the table and discard a portion of it (i.e. 1/2). Record proportion discarded. Sort through the remaining keeping all bivalves. Record weight (or count if there are no scales on board) of all razor clams. Measure all large commercial size razor clams. Be sure to record grade on datasheet. Measure a sample/subsample of the other smaller razor clams. Be sure to record grade on datasheet. Record count of all other bivalves. If quantities of other bivalves are too high, subsample the catch again (i.e. 1/2 or 1/3). Record the proportion discarded and then sort the bivalves and count the number of each retained. If quantities of some by-caught bivalve species are still very high (i.e. *Pharus legumen* and *Acanthocardia sp.*), then sub-sample them by recording the weight of the full sample and then record the weight and count of the subsample.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: Sampling design and protocols were not developed as part of a regional or multi-lateral agreement.

Link to sampling design documentation: <http://hdl.handle.net/10793/1688>

Compliance with international recommendations: N

The survey is a stratified random sampling design. Fishing pressure, estimated from high frequency VMS data, is used to stratify the survey. Spatial autocorrelation is accounted for in

the estimation using a geostatistical method developed in house. These methods are based on recent ICES and other training courses.

Link to sampling protocol documentation: <http://hdl.handle.net/10793/1688>

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: NA (surveys only take place once per year).

Data capture

Means of data capture: Razor clam lengths and weights are measured using measuring boards and scales and recorded on to hardcopy datasheets. The data is then entered in to excel spreadsheets and uploaded to an internal SQL shellfish database (FEAS_InshoreFisheries).

Data capture documentation: Razor clams are measured to the nearest millimetre below. A total weight of all or a subsample of razor clams per dredge haul is recorded. An SOP for sampling razor clams is held in Paradigm 3 (a document management system) and are reviewed and updated regularly.

Quality checks documentation: N

Quality checks are carried out on the biological data recorded during the survey using an R script prior to assessment being undertaken.

Data storage

National database: FEAS_InshoreFisheries

International database: NA

Quality checks and data validation documentation: Once the data has been recorded and collated for assessment data checks are carried out on the haul positions and biological data using an R script prior to assessment being undertaken.

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage:

Storage description: Some samples of the target species are collected to enable individual length and weight measurements to be recorded. These samples are kept in water on board where possible and once measured ashore these samples are returned to water on board the vessel prior to being replaced on the razor beds being surveyed.

Data processing

Evaluation of data accuracy (bias and precision): N

Surveys are stratified random design or on a systematic grid. Sampling efficiency (catchability of the hydraulic dredge sampling gear) is assumed to be 100%. There may be some underestimation bias if catchability is <1. Sampling procedures on board the survey vessel are either probabilistic random or census at a given station. Precision of biomass estimates is provided in the output of a geostatistical model.

Editing and imputation methods: Y

Where errors are identified then the preferred action is to correct the errors in the database – this might involve reference to the original datasheets if the error has occurred during transcription.

Quality document associated to a dataset: N

No DOI is currently created for the dataset, however the dataset can be requested.

Validation of the final dataset: Data checks are undertaken using an R-script prior to an assessment being run. If errors or anomalies are observed the data is either corrected by reference to the original datasheets (e.g. in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

No Deviation

Sampling Design and Protocols

No Deviation

Sampling Implementation

95% of the target dredge sampling was achieved but the population was considered to be adequately covered.

Data Capture

No Deviation

Data Storage

No Deviation

Sample Storage

No Deviation


Data Processing

No Deviation

Figure 7. RCIS Survey Map 2023 North Survey Zones.




Figure 7.1. RCIS Survey Map 2023 North Biomass Map.



Sampling scheme identifier: CNIS_IRL

MS : IRL
Region : North-East Atlantic
Sampling scheme identifier : CNIS_IRL
Sampling scheme type: Research Survey at sea
Observation type: SciObsAtSea
Time period of validity : 2007 onwards
The main objective of the Cockle North Irish Sea (CNIS_IRL) surveys is to estimate the biomass of cockles and to provide advice for <i>Cerastoderma edule</i> in Area 7a. Secondary objectives are to collect data on the distribution and abundance of other bivalve species such as <i>Angulus tenuis</i> and <i>Macoma balthica</i> caught during the surveys, along with recording the depth of the redox potential discontinuity layer and abundance of <i>Arenicola marina</i> sand castings at each sampling station. These data also support Habitats Directive Article 6 assessments.
Description of the population
Population targeted: The main cockle (<i>Cerastoderma edule</i>) bed in Dundalk Bay.
Population sampled: The target species is <i>Cerastoderma edule</i> in the intertidal zone of Dundalk Bay, Northwest Irish Sea.
Stratification: A 500x500 m survey grid is mapped over the intertidal sand flat. Each grid cell is divided into 400 sub-cells of 25 m ² in area and a quadrat (0.25 m ²) and raked sample (2 m ²) are collected from three randomly selected sub-cells. Since the survey began some more easterly grid cells were added to the northern survey area to ensure the eastern extent of the cockle bed was being surveyed.



Stratified random survey grid (500 x 500 m) for cockles in Dundalk Bay SAC/SPA, North Irish Sea.

Sampling design and protocols

Sampling design description: The PSU is the averaged area of a 0.25 m² quadrat and a 2 m² raked area. The location of each of the three sampling stations within each survey grid cell is selected randomly based on the 25 m² grid. Biomass of cockles at each sampling station is estimated as the product of density and mean individual weight calculated from the size distribution at the station and a weight-length relationship.

The sediment within each quadrat sample is dug out to a depth of approximately 30 cm and sieved through a 4 mm mesh. Rake samples were collected by raking over the 2 m² area to a depth of approximately 5 cm. Raking was included to increase the change of encountering larger cockles over 22 mm shell width (commercial size), which occur in lower abundance. All cockles are retained for size measurements. A sub-sample of the cockles retained are weighed and aged.

Numbers of cockles per sample are standardised to density per square meter. Densities at each station are averaged cross quadrat and rake samples prior to interpolation.

Counts of all *Angulus tenuis* and *Macoma balthica* are recorded from each sample along with a count of the *Arenicola marina* sand castings. The RPD layer depth is also measured where possible.

Is the sampling design compliant with the 4S principle?: NA Regional coordination: Sampling design and protocols were not developed as part of a regional or multi-lateral agreement. Link to sampling design documentation: http://hdl.handle.net/10793/1688 Compliance with international recommendations: N The sampling design for this survey is based on 3 randomly selected stations per 500 x 500 m² which spans the extent of Dundalk Bay. A quadrat (0.25 m²) and rake (2 m²) sample are collected from each sampling location. The sediment within each quadrat sample is dug out to a depth of approximately 30 cm and sieved through a 4 mm mesh. Rake samples were collected by raking over the 2 m² area to a depth of approximately 5 cm. Raking was included to increase the chance of encountering larger cockles over 22 mm shell width (commercial size), which occur in lower abundance. All cockles are retained for size measurements. A sub-sample of the cockles retained are weighed and aged. Numbers of cockles per sample are standardised to density per square meter. Densities at each station are averaged across quadrat and rake samples prior to interpolation. Link to sampling protocol documentation: http://hdl.handle.net/10793/1688
Sampling implementation Recording of refusal rate: NA Monitoring of sampling progress within the sampling year: NA (surveys only take place once per year).
Data capture Means of data capture: Cockle shell widths are measured using electronic callipers and individual weights are recorded using a precision scales (0.01g). All measurements and weights are recorded electronically in to an excel spreadsheet prior to being uploaded to an internal SQL shellfish database (FEAS_InshoreFisheries). Data capture documentation: Cockles are measured to the nearest millimetre below. Individual weights of cockles are recorded to 0.01 of a gram. An SOP for sampling (collecting, measuring, weighing and recording data) are held in Paradigm 3 (a document management system) and are reviewed and updated regularly. Quality checks documentation: N Quality checks are carried out on the biological data recorded during the survey using an R script prior to assessment being undertaken.
Data storage National database: FEAS_InshoreFisheries International database: NA Quality checks and data validation documentation: Once the data has been recorded and

collated for assessment data checks are carried out on the sample positions and biological data using an R script prior to assessment being undertaken.

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage

Cockle samples are collected to enable individual shell width and weight measurements and age data to be recorded. These samples are kept in labelled zip lock bags and stored chilled in freezer boxes. Once measured these samples are returned to the sand flats.

Data processing

Evaluation of data accuracy (bias and precision): N

The survey is a stratified random design. Sampling efficiency at point survey stations is 100%; all cockles at the sampling point are detected counted and measured. Probability of false zero or undercounting bias is absent. Precision of biomass estimates is provided in the output of a geostatistical model.

Editing and imputation methods: Y

Where errors are identified then the preferred action is to correct the errors in the database – this might involve reference to the original datasheets if the error has occurred during transcription.

Quality document associated to a dataset: N

No DOI is currently created for the dataset, however the dataset can be requested.

Validation of the final dataset: Data checks are undertaken using an R-script prior to an assessment being run. If errors or anomalies are observed the data is either corrected by reference to the original datasheets (e.g. in the case of input error) or excluded from that particular use.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Description of the Population

No Deviation

Sampling Design and Protocols

No Deviation

Sampling Implementation

Some areas were not accessible due to being located below the low tide mark during the survey and the varying movements of streams running through the sandflats from year to year. However, the survey area was covered adequately even though samples weren't collected from all the planned sampling points.

Data Capture

No Deviation

Data Storage

No Deviation

Sample Storage

No Deviation

Data Processing

No Deviation

Figure 8. CNIS Survey Map 2023 Survey Zones.





Figure 8.1. CNIS Survey Map 2023 Biomass Map.



Sampling scheme identifier: OWI_IRL

MS: IRL
Region: North-East Atlantic
Sampling scheme identifier: OWI_IRL
Sampling scheme type: Research Survey at sea
Observation type: SciObsAtSea
Time period of validity: 2010 onwards
The main objective of the Oyster West Ireland (OWI_IRL) surveys is to estimate the biomass of the native oyster and to provide advice for <i>Ostrea edulis</i> in Area 7. And to estimate the density and biomass of both the native (<i>Ostrea edulis</i>) and pacific (<i>Magallana gigas</i>) species in Area 6.
Description of the population
Population targeted: The swept area of all dredge hauls for the main target species, <i>Ostrea edulis</i> in Area 7 and both <i>Ostrea edulis</i> and <i>Magallana gigas</i> in Area 6.
Population sampled: The target species a is the native oyster, <i>Ostrea edulis</i> in Area 7 and <i>Ostrea edulis</i> and <i>Magallana gigas</i> in Area 6.
Stratification: Sampling locations are chosen randomly within a survey grid for most of the oyster surveys. Grid cells can vary in size from 100 x 100 m to 250 x 250 m depending on the extent of the oyster bed. A number of oyster beds occur as separate stocks in Bays around along the west and northwest coasts.
Sampling design and protocols
Sampling design description: The PSU is a dredge haul. Dredge designs vary locally, and

these locally preferred designs are used in the surveys. The location of each dredge haul is selected randomly within a survey grid. Biomass is estimated using a geostatistical model accounting for the spatial autocorrelation in the survey data.

Is the sampling design compliant with the 4S principle? NA

Regional coordination: Sampling design and protocols were not developed as part of a regional or multi-lateral agreement.

Link to sampling design documentation: <http://hdl.handle.net/10793/1688>

Compliance with international recommendations: N

Surveys are stratified random design or on a systematic grid. Sampling efficiency (catchability of the oyster dredge sampling gear) is assumed to be 32%. New estimates have recently been obtained by comparison with methods where catchability is known to be 1. Estimates are raised according to dredge efficiency. Sampling procedures on board the survey vessel are either probabilistic random or census at a given station.

Link to sampling protocol documentation: <http://hdl.handle.net/10793/1688>

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: NA (surveys only take place once per year).

Data capture

Means of data capture: Oyster lengths are measured using callipers and total sample weights are achieved using marine compensated scales. Both are recorded on to hardcopy datasheets. The data is then entered in to excel spreadsheets and uploaded to an internal SQL shellfish database (FEAS_InshoreFisheries).

Data capture documentation: Oysters are measured to the nearest millimetre below. A total weight of all or a subsample of oysters per dredge haul is recorded.

Quality checks documentation: N

Quality checks are carried out on the biological data recorded during the survey using an R script prior to assessment being undertaken.

Data storage

National database: FEAS_InshoreFisheries

International database: NA

Quality checks and data validation documentation: Once the data has been recorded and collated for assessment data checks are carried out on the haul positions and biological data using an R script prior to assessment being undertaken.

The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage
Storage description: Some samples of the target species are collected to enable individual length and weight measurements to be recorded. These samples are kept in water on board where possible and once measured ashore these samples are returned to water on board the vessel prior to being replaced on the oyster beds.
Data processing
Evaluation of data accuracy (bias and precision): N Sampling efficiency (catchability of the oyster dredge sampling gear) is assumed to be 32%. New estimates have recently been obtained by comparison with methods where catchability is known to be 1. Estimates are raised according to dredge efficiency. Sampling procedures on board the survey vessel are either probabilistic random or census at a given station. Precision of biomass estimates is provided in the output of a geostatistical model. Editing and imputation methods: Y Where errors are identified then the preferred action is to correct the errors in the database – this might involve reference to the original datasheets if the error has occurred during transcription. Quality document associated to a dataset: N No DOI is currently created for the dataset, however the dataset can be requested. Validation of the final dataset: Data checks are undertaken using an R-script prior to an assessment being run. If errors or anomalies are observed the data is either corrected by reference to the original datasheets (e.g. in the case of input error) or excluded from that particular use.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan <u>Description of the Population</u> No Deviation <u>Sampling Design and Protocols</u> No Deviation <u>Sampling Implementation</u> No Deviation <u>Data Capture</u> No Deviation <u>Data Storage</u> No Deviation <u>Sample Storage</u> No Deviation <u>Data Processing</u> No Deviation

Figure 9. OWI Survey Map 2023 Tralee Bay Survey Zones.




Figure 9.1. OWI Survey Map 2023 Inner Tralee Bay Survey Zones



Figure 9.2. OWI Survey Map 2023 Inner Tralee Bay Biomass Map *Ostrea edulis*




Figure 9.3. OWI Survey Map 2023 Clew Bay South Survey Zones




Figure 9.4. OWI Survey Map 2023 Clew Bay South Abundance *Ostrea edulis*

Figure 9.5. OWI Survey Map 2023 Inner Galway Bay Survey Zones



Figure 9.6. OWI Survey Map 2023 Inner Galway Bay Biomass Map *Ostrea edulis*




Figure 9.7. OWI Survey Map 2023 Cill Chiaráin Bay Survey Zones





Figure 9.8. OWI Survey Map 2023 Cill Chiaráin Bay Biomass Map *Ostrea edulis*



Sampling scheme identifier: ICEos_IRL

MS : IRL

Region: North-East Atlantic

Sampling scheme identifier: ICEcoS IRL

Sampling scheme type: Research survey at sea

Observation type: SciObsAtSea (Scientific observer at sea on commercial or scientific vessels)

Time period of validity: 2023 onwards

The main objective of the ICEco Survey is to provide a new time series of abundance and distribution for key stocks within recruitment/nursery areas along the south and west coast of Ireland. Including but not limited to cod, whiting, haddock, plaice, sole, lemon sole, turbot, flounder, John dory, skates, rays and other elasmobranchs. This standardised inshore survey is timed to observe juvenile fish abundances, and associated ecosystem information.

Occurrence of vulnerable or sentinel invertebrate species such as corals, sea pen, fan mussel and ocean quahog is also noted. Marine litter is also sorted and recorded. Oceanographic data are collected from CTD instrument on trawl door.

Description of the population: Main target species are haddock (*Melanogrammus aeglefinus*), whiting (*Merlangius merlangus*), cod (*Gadus morhua*), pollack (*Pollachius pollachius*), sole (*Solea solea*), plaice (*Pleuronectes platessa*).

Population sampled: As a multi-species survey the target ecosystem component is demersal

species. No sampling takes place outside the survey area or on grounds that are unsuitable for trawling.

Stratification: The stratification is yet to be finalised, however the Surveys will be undertaken on 10-15m commercial vessels chartered for 5 days in 8 distinct areas, with at least 3 tows would be conducted each day and the total number of stations would be in the order of 120 survey tows.




Figure 1 The eight approximate survey areas are shown with the red rectangles.

Sampling design and protocols


Sampling design description: Individual hauls are the PSU, these are selected from random locations inside each stratum. The catch will be processed based on a protocol which is under development.

Is the sampling design compliant with the 4S principle?: NA (but note that the sampling scheme is statistically sound) Regional coordination: National Survey Link to sampling design documentation: Under development Compliance with international recommendations: Y Link to sampling protocol documentation: Under development
Sampling implementation Recording of refusal rate: NA Monitoring of sampling progress within the sampling year: NA (the survey only takes place once per year).
Data capture Means of data capture: Initially by hand – measuring boards and paper. These data are stored in a SQL database (FSS_SURVEY) on a local server. The gear sensor data as well as bottom depth and GPS position are also automatically recorded in a SQL database (FSS_NMEA) at intervals of approximately one per second. In future Catch weights, length frequency distributions and biological data will be captured using the EFDAQ (Electronic Fisheries Data Acquisition) system and stored in a local database in the wet laboratory before being imported into the central SQL database (FSS_SURVEY). Data capture documentation: Under development Quality checks documentation: Under development
Data storage National database: FSS (Fishing Survey System) International database: Not yet applicable Quality checks and data validation documentation: Once a survey is complete a number of data checks are carried out on haul positions, gear geometry, catch data and internal consistency of the data. The data management of this data collection activity is incorporated into the Marine Institute's IODE accredited Data Management – Quality Management Framework (DM-QMF).

Sample storage
Samples for aging such as otoliths are stored clean and dry at room temperature before transport to Marine Institute laboratories for analysis. These age structures are generally stored at Marine Institute premises for a period of months before age reading is carried out. Soft tissues are generally collected by request from third parties such as universities and are stored according to protocols provided. Such samples are transported to third parties within weeks of survey completion.
Data processing
Evaluation of data accuracy (bias and precision): N There is no procedure in place to estimate bias. Editing and imputation methods: NA – no imputation take place (with the exception of gear parameters that could not be observed – these are imputed using a model based on observed values). Quality document associated to a dataset: Procedure for producing the estimations of abundance and biomass for main species is included in annual survey report. Validation of the final dataset: Final dataset will be checked for quality.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan <u>Description of the Population</u> No Deviation <u>Sampling Design and Protocols</u> No Deviation <u>Sampling Implementation</u> Some days at sea were lost due to bad weather, but 100 stations were achieved in 2023. <u>Data Capture</u> No Deviation <u>Data Storage</u> No Deviation <u>Sample Storage</u> No Deviation <u>Data Processing</u>

No Deviation

Figure 10. ICEcoS achieved Sampling in 2023



Sampling scheme identifier: SS_IRL

MS : IRL

Region: North-East Atlantic

Sampling scheme identifier: SS_IRL

Sampling scheme type: Research survey at sea

Observation type: SciObsAtSea (Scientific observer at sea on commercial or scientific vessels)

Time period of validity: 2023 onwards

The main objective of the SS_IRL sampling scheme is to develop a research survey times

series for scallop (*Pecten maximus* and *Aequipecten opercularis*) in the western Irish Sea and north Celtic Sea. Data on distribution, relative abundance and spatially defined size and age structure will be generated.


Description of the population

Population targeted: Scallop populations in western Irish Sea (ICES VIIa) and northern Celtic Sea (VIIg)

Population sampled: Scallop populations in western Irish Sea (ICES VIIa) and northern Celtic Sea (VIIg) Distribution of sampling in the initial year will also take into account Industry expert knowledge. VMS of scallop fishing vessels in previous years will be reviewed and used to define the probable distribution of scallop populations in the area. Data for surveys in 2001-2005 in the Celtic Sea will be reviewed.

Stratification: Will be based on sediment layers sourced from Emodnet or acoustic backscatter layers from Irelands national seabed survey (INFOMAR). Scallop abundance is correlated with multibeam acoustic backscatter which reflects differences in sediment grain sizes in sand and gravel fractions.

Likely distribution of the survey is outlined below based on underlying VMS data for vessels targeting scallop in 3 areas and current information on activity of vessels under 12m which do not report VMS data.



Sampling design and protocols
Sampling design description: The PSU is a dredge haul. The location of each dredge haul will be selected randomly within sediment strata. Relative biomass will be estimated using a geostatistical model accounting for the spatial autocorrelation in the survey data and which incorporates sediment data as a co-variate. Catchability estimates for scallop dredges can be used to derive absolute biomass. Is the sampling design compliant with the 4S principle?: N/A Regional coordination: N/A Link to sampling design documentation: New protocols will be developed for this survey in 2023 Compliance with international recommendations: Design of scallop surveys have been discussed at ICES WGScallop. The proposed survey will follow WGScallop recommendations and the survey data will be reported to WGScallop. Link to sampling protocol documentation: New documentation for this survey will be developed in 2023
Sampling implementation
Recording of refusal rate: NA Monitoring of sampling progress within the sampling year: NA (the survey only takes place once per year).
Data capture
Means of data capture: Survey station tow data will be recorded using digital methods (GPS survey units onboard) defining the length of the tow. Number of dredges and dredge width * tow length provides swept area estimates. Catch (numbers and weight), size of scallop (all or sample depending on catch volume) and age (sample) will be recorded for each station. Data capture documentation: Protocol for the survey will be documented in 2023 Quality checks documentation: Yes this will be described in the data capture documentation
Data storage
National database: Data will be stored in Inshore Fisheries_SQL database at Marine Institute International database: Data will be provided to relevant data calls Quality checks and data validation documentation: R scripts will be used to identify data outliers and to visualise data prior to upload to database

Sample storage
A sample of scallop shells will be retained for age validation at ageing workshops run by WGScallop
Data processing
Evaluation of data accuracy (bias and precision): Confidence intervals are reported for final estimates of relative abundance
Editing and imputation methods: Various checks (visualisation) are done during data input and prior to data upload. Survey track data is screened (R-script) for GPS outliers. Perceived errors in data and checked against original datasheets.
Quality document associated to a dataset:
Validation of the final dataset: Data checks are undertaken using an R-script prior to an assessment being run. If errors or anomalies are observed the data is either corrected by reference to the original datasheets (e.g. in the case of input error) or excluded from that particular use
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan
<u>Description of the Population</u> No Deviation
<u>Sampling Design and Protocols</u> No Deviation
<u>Sampling Implementation</u> Certain Scallop beds in the North Irish Sea were inaccessible due to static gear on the ground. Also no favourable combination of tides, weather conditions and vessel availability were achieved in 2023 to undertake the South Irish Sea and Celtic Sea Scallop surveys, which have been re-scheduled to be undertaken in 2024. Therefore, the number of days at sea and number of sampling activities were not fully defined in 2023.
<u>Data Capture</u> No Deviation
<u>Data Storage</u> No Deviation
<u>Sample Storage</u> No Deviation
<u>Data Processing</u> No Deviation

Figure 11. Scallop Survey Bray Inshore Survey Zones



Figure 11.1. Scallop Survey Bray Inshore Biomass Map *Pecten maximus*




Figure 11.2. Scallop Survey Bay Offshore Survey Zones




Figure 11.3. Scallop Survey Bay Offshore Biomass Map *Pecten maximus*





Figure 11.4. Scallop Survey Dundalk Bay Offshore Survey Zones



Figure 11.5. Scallop Survey Dundalk Bay Offshore Biomass Map *Pecten maximus*



ANNEX 1.2 - QUALITY REPORT FOR SOCIOECONOMIC DATA SAMPLING SCHEME

The quality report fulfils Article 6 (3) (d) of the Regulation (EU) 2017/1004. This document is intended to specify data to be collected under chapter II, points 3, 5, 6, and 7 of the Delegated Decision annex: Socioeconomic data on fisheries, aquaculture and any complementary data collection of fishing activity and fish processing.

Use this document to describe quality aspects of the data collection process (design, sampling implementation, data capture, data storage and data processing etc.). The annex should be filled for each sampling scheme. Where applicable, use the handbook on sampling design (Deliverable 2.1 from MARE/2016/22 SECFISH study), available on the DCF website.

Provide information under each point in all sections. Do not delete any text from the template.

Quality Report Related to Table 5.2 and Text Box 5.2

Sector Name(s): Fisheries - Non-Probability Sample Survey

Survey Specifications
<i>Sector name refers to socio economic data on fisheries, aquaculture and any complementary data collection of fishing activity and processing as given in the EU MAP Delegated Decision annex.</i> <i>Sampling scheme refers to survey technique: by census, by sampling, random or non-random, other (with explanation). If sampling, then outline sampling design.</i> <i>Variables refer to Tables 7, 9 and 10 of the EU MAP Delegated Decision annex. Supra region refers to Table 2 of the EU MAP Implementing Decision annex. If the sampling scheme is the same in all supra regions put 'All Supra regions'.</i>
Sector name(s): Fisheries
Sampling scheme: Non-Probability Sample Survey
Variables: Consumption of fixed capital, Days at sea, Employment by age, Employment by employment status, Employment by gender, Employment by level of education, Employment by nationality, Energy consumption, Energy costs, FTEs by gender, Full-time equivalent (FTE), Gross debt, Gross value of landings, Income from leasing out quota or other fishing rights, Investments in tangible assets (net purchase of assets), Lease/rental payments for quota or other fishing rights, Mean age of vessels, Mean LOA of vessels, Number of fishing enterprises/units, Number of vessels, Operating subsidies, Other income, Other non-variable costs, Other variable costs, Paid labour, Personnel costs, Repair and maintenance costs, Subsidies on investments, Total hours worked per year (optional), Total value of assets, Total vessel power, Total vessel tonnage, Unpaid labour,

Unpaid labour by gender, Value of physical capital, Value of quota and other fishing rights, Value of unpaid labour
Supra region(s): Baltic Sea; North Sea; Eastern Arctic; NAFO; extended North Western waters (ICES areas 5, 6 and 7) and extended South Western waters (ICES areas 10, 12 and 14)
Survey planning
Provide a short description of the population the sampling scheme applies to; e.g. <i>‘less active vessels using passive gears’</i>. The target population is the “commercial fishing fleet” as recorded in the EU Fleet Register on the last day of the reference year. Fleet Segmentation: The segmentation of the fleet, will follow the guidelines in Table 8 of Commission Delegated Decision (EU) 2021/1167 and is used to stratify the collection of all, non-transversal, economic parameters. The following data sources will be used to segment the fleet: • EU Fleet Register on the 31st December for the reference year. • EU log-book activity records for vessels active in the reference year (>10 meters); • Sentinel Vessel Programme Effort Data • Recorded fishing activity from previous economic surveys. Individual vessels are assigned to fleet segments by overall length (LOA) class and the main fishing method engaged in by the vessel, in the previous calendar year. In cases where there is a risk of natural persons and/or legal entities being identified clustering may be applied to report economic variables in order to ensure statistical confidentiality. Such a clustering scheme shall be consistent over time. The source of information used to distinguish the sampling frame from the target population, will be based on EU logbook data as follows: • Active Vessels: For vessels greater than 10 meters in overall length, only those with

at least one entry in the EU log-book, in the reference year, will be deemed active. This analysis will take place once the log-book data are available for a particular reference year, which is approximately 3 months after the end of the calendar year (March $n-1$);

- For vessels less than 10 meters in overall length, an estimate of inactivity will be conducted each year using all available sources, including: previous survey responses, the National Inshore Sentinel programme, sales notes data and the fleet register.

Required sampling intensities have been estimated using statistical analysis of the previous year's survey data. The analysis determines required sample size n , based on the mean of a finite population, to achieve a given level of precision (e.g., a CV of 25% on the sample mean).

Applying the function, we can see that for very low CV, all vessels need to be sampled and that the required sample number increases with the standard deviation of the segment. However, due to the finite population function you can never sample more than the full population (census). Some segments have a planned sample rate of 0% as the number of active vessel in the segment are very low ($n \leq 5$).

Survey design and strategy

List data sources; e.g. interviews, registers, log books, sales notes, VMS, financial accounts etc.

Describe how the sample sizes were determined.

Describe survey methods and distribution; e.g. questionnaire forms by post, by email, on website, by phone etc. access to other datasets etc.

Describe the role of auxiliary information, if any, in the strategy: e.g. for validation, cross referencing, fall back data source etc.

The data sources used to collect economic and social data from fleet segments are:

- Sales notes data for landing income for vessels under 10m.
- Logbook data for effort and landing income for vessels over 10m.
- Voluntary questionnaire information returned by vessel owners targeted in the annual economic survey for all economic and social variables.
- Face-to-face/phone interviews with vessel owners to clarify any issues arising with economic and social variables from questionnaire.
- Mandatory economic and social questionnaire information returned by vessel owners applying for EU/National grant aid,
- Data from vessel owners from a national Sentinel Vessel Programme (to collect both transversal and non-transversal economic and social data from vessels in the small scale fisheries where log-book declarations are not mandatory). See next Annex for details.

The population shall be all active and inactive vessels registered in the Union Fishing Fleet Register as defined in Commission Regulation (EC) No 26/2004 (2) on 31 December of the reporting year and vessels that do not appear on the Register at that date but have fished at least one day during the reporting year.

The data sources for the national implementation for the fleet target population are:

- EU Fleet register;
- EU Log-book data.

Estimation design

Describe method of calculating population estimate from sample.

Describe method of calculating derived data: e.g. imputed values.

Describe treatment of nonresponse.

Recognising the implications and influences imposed by the voluntary nature of the annual survey on the probability sample survey design standard appropriate raising techniques will be used, to derive final estimates for each variable collected. This methodology was reviewed in 2018, which resulted in a report to assess and improve the raising estimations. Various methods of raising are possible and this report set to establish a theoretical and empirical basis for the decision as to how best to raise sampled economic data to the fleet level.

The mean squared error (MSE) encapsulates the bias and variance of an estimator. The MSE was used as the basis for comparing raising performance. We first derive theoretical expectations on which raising method would work best when there is or is not a relationship with fishing effort. Raising methods were then tested on the real data via re-sampling and appraisal of the ability of various raising methods to recover the true sum. A suite of specifically developed visualisation code assists in appraising the distribution of the data, in particular with identifying outlying values that can overly influence the raised sum.

From the theoretical analyses there were two major conclusions for raising sample data:

1. Where a variable is independent of effort it is best to raise the average to the segment level as the inclusion of unrelated effort adds additional variance to the estimator.
2. Where a variable is proportional to effort, raising based on effort will provide a better estimator as long as the residual variance of the relationship between the economic variable and effort is comparatively small and the strength of the proportionality constant comparatively large. In other words if there is a strong relationship with effort, raise by effort, if not raise by the average. A ‘strong’ relationship is defined as having less bias associated with raised estimating using this theory.

A harmonised FTE will be estimated for each of the fleet segments. For vessels >10 meters in length (LOA), operational data from log-book submissions will be used in the estimation of fishing time on a trip-by-trip basis. In addition, there are several questions on the annual economic survey forms that deal specifically with hours worked and the nature of the engagement (full-time, part-time, casual). Questions regarding gender breakdown and age profiles, education and nationality have been added to the annual survey.

Error checks

Describe potential errors and how and where in the process these are detected, avoided or eliminated e.g., data; duplication, double counting, respondent error, upload error, processing error etc.

The issue of consistency of data coming from different data sources is recognized as being of significant importance. The introduction of bias in this area, is under continual assessment and is currently being addressed by restricting acceptance of data to a small number of official data streams (i.e. data items consistent with fields in annual company returns (provided via accountants), EU logbook data and Sales notes data).

Although error associated with bias and variability will effectively be introduced if observed returns do not match those expected, these descriptors will be reported where possible and with appropriate caveats.

Data storage and documentation
<i>Describe how the data is stored.</i> <i>Provide link to webpage where additional methodological documentation can be found, if any.</i> The data from the electronic forms are stored as .xml files and imported into a database. Data received through postal surveys, or phone surveys are entered into the electronic forms and submitted as .xml files to the same database. The data is stored on a secure server which is only accessible by EU MAP staff. Published documentation of methodology of all EUMAP data-collection is found at: https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/DCF_Methodology_Economic_IRL_2020.pdf The MS wants to move away a paper-based system with manual data entry towards an online data entry system where data is stored in a database as there are still some paper based surveys circulated to the industry. The first phase of development in late 2021 will be for the fisheries industry whereby survey forms will be available online and these will connect with a database back end. This will enable the digital capturing, storing and reporting of data. The EU - MAP system will facilitate the entry of data through an online web portal and through a data entry and integration layer which will allow for access to data in current and future systems. The EU - MAP system will include the development of a scalable database storage and support reporting through a BI module. The main objective of this system to support the mandatory EU - MAP reporting.
Revision
Describe the frequency of the methodology review e.g., revision of; segmentation, survey method per segment, per variable etc. Segmentations are review annually to ensure that they are maintain confidentiality. The current national work programme is moving to an on-line platform for collecting all EU MAP data. This will be operational in 2022.
Confidentiality

Are procedures for confidential data handling in place and documented? Are protocols to enforce confidentiality between DCF partners in place and documented? Are protocols to enforce confidentiality with external users in place and documented? Are there any issues with publication of data due to confidentiality reasons? Provide an explanation. <i>Are procedures for confidential data handling in place and documented?</i> The frame population are made aware of BIM's <u>privacy policy</u> and <u>data protection policies and procedures</u>. <i>Are protocols to enforce confidentiality between DCF partners in place and documented?</i> There is a Data Sharing Agreement between the two main agencies carrying out the DCF work, the Marine Institute and Bord Iascaigh Mhara (BIM). <i>Are protocols to enforce confidentiality with external users in place and documented?</i> Data protocols to enforce confidentiality are followed strictly. In cases where there is a risk of natural persons and/or legal entities being identified clustering is applied to report economic variables to ensure statistical confidentiality. For internal business units and some closely related agencies, formal data-sharing agreements between the data controller and data processor must be in place before an appropriate level of data-sharing can be permitted. Are there any issues with publication of data due to confidentiality reasons? Provide an explanation NA	
AR comment: Indicate any deviations. Do not change the text already adopted in the work	

plan

Sampling Scheme

No deviation from WP in 2023.

Survey Planning

No deviation from WP in 2023.

Survey Design and Strategy

No deviation from WP in 2023.

Estimation Design

No deviation from WP in 2023.

Error Checks

No deviation from WP in 2023.

Data Storage and Documentation

No deviation from WP in 2023.

Revision

No deviation from WP in 2023.

Confidentiality

No deviation from WP in 2023

Quality Report Related to Table 5.2 and Text Box 5.2

Sector Name(s): Fisheries – Sentinel Vessel Programme

Survey Specifications

Sector name refers to socio economic data on fisheries, aquaculture and any complementary data collection of fishing activity and processing as given in the EU MAP Delegated Decision annex.

Sampling scheme refers to survey technique: by census, by sampling, random or non-random, other (with explanation). If sampling then outline sampling design.

Variables refer to Tables 7, 9 and 10 of the EU MAP Delegated Decision annex. Supra region refers to Table 2 of the EU MAP Implementing Decision annex. If the sampling scheme is the same in all supra regions put 'All Supra regions'.

Sector name(s): Fisheries – Sentinel Vessel Programme

Sampling scheme: Non-Probability Sample Survey
Variables: • Days at sea • Fishing days • Live Weight of landings total and per species (to augment data from Sales Notes) • Average prices per species (to augment data from Sales Notes)
Supra region(s): Baltic Sea; North Sea; Eastern Arctic; NAFO; extended North Western waters (ICES areas 5, 6 and 7) and extended South Western waters (ICES areas 10, 12 and 14)
Survey planning The Member State will continue to collect transversal data, on a daily basis, from vessels < 12 meters in length (LOA) in a national, Sentinel Vessel Programme (SVP). This is justified on the basis that it is not currently possible to define quantitative targets for a sampling programme for transversal parameters within metiers containing an inshore component; specifically for vessels <10 metres LOA and where official declarations of their landings are not required. In accordance with Article 11(4) of Council Regulation (EC) No 199/2008 – defining the eligibility of self-sampling aboard Community fishing vessels – vessel owners participating in the sentinel programme will record their daily landings, effort and price data in a ‘sentinel record book’, specifically designed to capture these data, for a full calendar year. Additionally, and to enhance the cost-effectiveness of this programme; • Non-transversal economic parameter will also be collected, again on a daily basis, by vessels < 12 meters (LOA) participating in the sentinel programme, these are included in Table 5.2; • Non-operational economic parameters will also be collected at the end of the reference year. These data will be collected using a survey and an exit interview from the sentinel programme;

- Biological (length composition) and discard information will be collected on a weekly basis by vessels participating in the sentinel programme to complement the data on biological variables in these metiers.
- Daily landings and price figures are collected

Survey design and strategy

The sample population for the sentinel programme is designed to represent the diverse inshore fisheries sector in Ireland. Specific fisheries are targeted that represent gear usage, target species, and geographical location. The list of participants is reviewed annually to ensure that it remains representative of the small scale fisheries.

Logbooks are provided to participants every January. The logbooks record, vessel details and economic data annually. Daily logbook entries record target species, effort, landings, and economic data (fish prices, fuel prices). Weekly logbook entries record biological data of the catches. Logbooks are collected at the end of the year and data is entered and checked.

Those vessels accepted into the sentinel programme are remunerated to the amount of €1000 per vessel per annum. Remuneration is based on an average of 180 days at sea per year, and an estimated 30 - 40 minutes to record the data in the supplied sentinel record book. When the additional time commitment required from each participating vessel owner for direct contact with survey personnel, to complete the exit interview required at the end of the data collection period and to provide non-operation data are included, the total commitment amounts to 115 hours per year. Using the national minimum wage in Ireland (currently €8.65 per hour for adults over the age of 18) the total annual cost of 115 hours at €8.65 per hour is €1,000. This remuneration is deemed the minimum necessary to attract eligible fishermen into the programme and to ensure accurate and reliable data are recorded.

Estimation design

Operational data from log-book submissions will be used in the estimation of fishing time on a trip-by-trip basis to estimate Days at sea and Fishing days.

Error checks
Describe potential errors and how and where in the process these are detected, avoided or eliminated e.g., data; duplication, double counting, respondent error, upload error, processing error etc. The data collected under the SVP are entered by MI analysts. Data entry checks and validation occurs at data entry and the data are visually checked for outliers. Each book is validated by a second person
Data storage and documentation
<i>Describe how the data is stored.</i> <i>Provide link to webpage where additional methodological documentation can be found, if any.</i> SVP books are distributed to personnel in the regional laboratories and headquarters where they are held in locked cupboards. After three years they are archived offsite in secure storage areas. The SVP books are transcribed into Excel Spreadsheets before being uploaded into a central SQL Server Inshore Fisheries database. The data is subject to a series of integrity checks during upload – any errors are flagged to the user. If errors do occur the user must resolve them before re-uploading the spreadsheet. The Excel spreadsheets are stored within a structured network folder system, which reside on a secure a Microsoft Windows Network maintained by the IT Operations Department within the Marine Institute. The SQL Server Inshore Fisheries database containing the uploaded SVP data resides on a Marine Institute secure production centralised server. Access to the network folders system and/or database is controlled by membership to a specific Windows Work Group that is maintained by IT Operations. If a new member of staff requires access to the network folder system containing the SVP Excel Spreadsheets or the database, an approval process is place, which determines whether the member of staff should be granted access. If/when a member a staff member leaves the Marine Institute access to the specific windows group is revoked by IT Operations A new graphical user interface is currently in development which will allow SVP data to be entered directly into the SQL Server Inshore Fisheries Database. Access to the new interface is controlled by the same Windows Group Membership required for the database.
Revision
Describe the frequency of the methodology review e.g., revision of; segmentation, survey

method per segment, per variable etc. Segmentations are review annually to ensure that they are maintain confidentiality.
Confidentiality
Are procedures for confidential data handling in place and documented? Yes, access to the physical SVP logbooks, the transcribed spreadsheets, and the database is controlled via a defined procedure. Are protocols to enforce confidentiality between DCF partners in place and documented? Are protocols to enforce confidentiality with external users in place and documented? Are there any issues with publication of data due to confidentiality reasons? Provide an explanation.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan <u>Sampling Scheme</u> No deviation from WP in 2023. <u>Survey Planning</u> No deviation from WP in 2023. <u>Survey Design and Strategy</u> A review of the participants at the end of 2022 highlighted a trend of reducing numbers in recent years. In 2023, the number of participating vessels was increased by 11 vessels (17%) based on gear usage, target species, and geographical location to ensure the Sentinel Vessel Programme remains representative of the SSCF in Ireland. Following feedback from participants, the remuneration figure was increased from €1000 to €1250 per vessel per annum. <u>Estimation Design</u> No deviation from WP in 2023.

Error Checks

No deviation from WP in 2023.

Data Storage and Documentation

No deviation from WP in 2023.

Revision

No deviation from WP in 2023.

Confidentiality

No deviation from WP in 2023.

Quality Report Related to Table 6.1 and Text Box 6.1

Sector Name(s): Aquaculture Census

Survey Specifications

Sector name refers to socio economic data on fisheries, aquaculture and any complementary data collection of fishing activity and processing as given in the EU MAP Delegated Decision annex.

Sampling scheme refers to survey technique: by census, by sampling, random or non-random, other (with explanation). If sampling, then outline sampling design.

Variables refer to Tables 7, 9 and 10 of the EU MAP Delegated Decision annex. Supra region refers to Table 2 of the EU MAP Implementing Decision annex. If the sampling scheme is the same in all supra regions put 'All Supra regions'.

Sector name(s): Aquaculture

Sampling scheme: Census

Variables:

- Fish feed used
- Full-time equivalent (FTE)
- Gross sales per species
- Livestock used
- Number of enterprises by size category
- Operating subsidies

• Paid labour • Raw material: feed costs • Raw material: livestock costs • Subsidies on investments • Weight of sales per species • Employment by employment status • Employment by gender • FTEs by gender
Supra region(s): Baltic Sea; North Sea; Eastern Arctic; NAFO; extended North Western waters (ICES areas 5, 6 and 7) and extended South Western waters (ICES areas 10, 12 and 14)
Survey planning
Provide a short description of the population the sampling scheme applies to; e.g. <i>‘less active vessels using passive gears’</i>. The population covered by this census survey are all licenced businesses and their production units, known to be producing or endeavouring to produce products by aquaculture and who employ themselves and/or staff to do so. In 2021 these totalled 264 businesses, operating 308 production units and are made up predominantly of farmed salmon, oyster mussel, and most recently micro-algal segments, located along the coast, mostly the west coast, with a small number of land-based finfish units inland. The marine units are mostly inshore, using on-bottom, off-bottom or suspended cultures while penned salmon culture occurs in more exposed locations. All production meets certification standards, with little, or no, medicine or other chemical input. Finfish is dominated by salmon production while shellfish output is almost entirely consisting of bivalve molluscs. The majority of enterprises are micro-businesses. No thresholds are applied. If necessary, in order to protect individual business data, micro-segments below 4 production units in membership, are incorporated into suitable larger segments in order to enable total national value estimates to be reported.
Survey design and strategy
Estimation design
Describe method of calculating population estimate from sample.

Describe method of calculating derived data: e.g. imputed values.

Describe treatment of nonresponse.

The response rate from the census survey, on average, is 78-82%.

Non-respondent performance can be estimated by a combination of methods. The Irish Seafood Development Agency's (BIM) regional aquaculture staff can estimate production from frequent site visits to the businesses in question. The Government Department responsible for aquaculture licensing can provide area data which in turn indicates production capacity while the unit sales value of adjacent compliant producers of the same product can be used for output value estimation purposes. Historical averages, if available, together with capacity data can also be used for this kind of imputation. Sister agencies supply aggregated data that may be used in place of in-house estimates if deemed to be of better quality. Finally, the most recent historical data received can be used in the case of enterprises known by local officers to have steady outputs, with their endorsement. Imputed data is obtained for some social variables. This is done using the empirical value of employment status given for some companies with more than one production unit and culture. The turnover per production unit is known. The turnover contribution per single FTE for the business as a whole is calculated and the empirical breakdown of production status given for the business as a whole, is then used to estimate employment status per production unit.

Non-Respondents are given repeated reminders to participate in the annual survey and will receive up to three phone calls until a date at the end of March when preparations to report on the annual survey must begin.

Error checks

Describe potential errors and how and where in the process these are detected, avoided or eliminated e.g., data; duplication, double counting, respondent error, upload error, processing error etc.

Double counting can occur within multi-production units of the salmon and oyster segments, especially if a given unit engages in just part of the production cycle. An on-growing unit may send its product to an in-house finishing unit, in the case of oyster production or to a harvesting station in the case of salmon production. Specific turnovers may be generated at each stage, but the volume could be counted twice. Close engagement with the producer reduces, if not eliminates this issue.

Phone surveys are a great source of quality data, but incorrect figures may be given or

accurate data mis-recorded. Errors of decimal placing, or poor hand-writing can result in false data spikes after uploading. These become apparent in graphics set up specifically to find these outliers in the database and the cause of the spike can be investigated back to the data supplier if necessary. The survey questionnaire form is an electronic form, and this reduced errors and as such is advocated as the most accurate way to receive data over postal surveys. The formatting of these electronic forms are set to avoid ambiguous entries.

Data storage and documentation

Describe how the data is stored.

Provide link to webpage where additional methodological documentation can be found, if any.

The data from the electronic forms are stored as .xml files and imported into a database. Data received through postal surveys, or phone surveys are manually entered into the same database. The data is stored on a secure server which is only accessible by EU MAP staff.

Published documentation of methodology of all EUMAP data-collection is found at:

https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/DCF_Methodology_Economic_IRL_2020.pdf

Aquaculture methodology is described from page 11.

Revision

Describe the frequency of the methodology review e.g., revision of; segmentation, survey method per segment, per variable etc.

This is done annually or as the need arises. The smaller segments require close attention as these may have to be amalgamated with others if numbers within a segment are in decline. There may also be new growth segments that need to be accounted for in the sampling design and or segment aggregations. For example, seaweed segmentation is under continuing review due to segment growth predictions and for reporting purposes from 2023 onward, the national micro-algal (or seaweed) segment has developed to the point with sufficient new companies now at a commercial phase that data can be disaggregated into the stand-alone segment 'Longline Seaweeds.'

Confidentiality

Are procedures for confidential data handling in place and documented?

The frame population are made aware of BIM's privacy policy and data protection policies and procedures.

Are protocols to enforce confidentiality between DCF partners in place and documented?

There is a Data Sharing Agreement between the two main agencies carrying out the DCF work, the Marine Institute and Bord Iascaigh Mhara (BIM).

Are protocols to enforce confidentiality with external users in place and documented?

Data protocols to enforce confidentiality are followed strictly. In cases where there is a risk of natural persons and/or legal entities being identified clustering in the form of segment amalgamations is applied to report economic variables to ensure statistical confidentiality. For internal business units and some closely related agencies, formal data-sharing agreements between the data controller and data processor must be in place before an appropriate level of data-sharing can be permitted.

Are there any issues with publication of data due to confidentiality reasons? Provide an explanation

Production and reporting of data from the salmon on-growing segment is restricted by the small number of operators involved and the fact that one very much dominates the segment. There is close engagement between BIM, as the data controller, and the concerned parties manage the appropriate level of reporting of this segment and other similar segments of small populations.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Sampling Scheme

No deviation from WP in 2023.

Survey Planning

No deviation from WP in 2023.

Survey Design and Strategy

No deviation from WP in 2023.

Estimation Design

No deviation from WP in 2023.

Error Checks

No deviation from WP in 2023.

Data Storage and Documentation

No deviation from WP in 2023.

Revision

No deviation from WP in 2023.

Confidentiality

No deviation from WP in 2023.

Quality Report Related to Table 6.1 and Text Box 6.1

Sector Name(s): Aquaculture NPS PSS

Survey Specifications

Sector name refers to socio economic data on fisheries, aquaculture and any complementary data collection of fishing activity and processing as given in the EU MAP Delegated Decision annex.

Sampling scheme refers to survey technique: by census, by sampling, random or non-random, other (with explanation). If sampling then outline sampling design.

Variables refer to Tables 7, 9 and 10 of the EU MAP Delegated Decision annex. Supra region refers to Table 2 of the EU MAP Implementing Decision annex. If the sampling scheme is the same in all supra regions put 'All Supra regions'.

Sector name(s): Aquaculture
Sampling scheme: NPS and PSS
Variables: Consumption of fixed capital Energy costs Financial expenditures Financial income Number of hours worked by employees and unpaid workers (optional) Other income Other operating costs Personnel costs Repair and maintenance costs Unpaid labour Value of unpaid labour Employment by age Employment by level of education Employment by nationality Unpaid labour by gender Gross debt Investments in tangible assets (net purchase of assets) Total value of assets
Supra region(s): All
Survey planning
Provide a short description of the population the sampling scheme applies to; e.g. <i>‘less active vessels using passive gears’</i> . The population covered by this NPS Sample survey are of all licenced businesses and their

production units, known to be producing or endeavouring to produce products by aquaculture and who employ themselves and / or staff to do so. A total of 264 businesses, operating 308 production units in 2021 are made up predominantly of Farmed Salmon, oyster mussel, and most recently, micro-algal (or seaweed) segments. The majority of these are located along the west coast with a small number of land-based finfish units inland. The marine units are mostly inshore, using on-bottom, off bottom or suspended cultures while penned salmon culture occurs in more exposed locations. All production is to certified standards, with little or no medicine or other chemical input. Finfish is dominated by salmon production while shellfish output is almost entirely of bivalve molluscs. The majority of enterprises are micro-businesses. The population is covered every 4-5 years by the sample questionnaire which pursues a rotating 25% profile of the population. The online survey, covering up to 33% of the population annually, includes annually sampled, indicator companies, maintained for their pivotal role within their segments.

Survey design and strategy

List data sources; e.g. interviews, registers, log books, sales notes, VMS, financial accounts etc.

Describe how the sample sizes were determined.

Describe survey methods and distribution; e.g. questionnaire forms by post, by email, on website, by phone etc. access to other datasets etc.

Describe the role of auxiliary information, if any, in the strategy: e.g. for validation, cross referencing, fall back data source etc.

There are two main data sources for the variables listed above; the producers themselves, via annual sample questionnaire and abridged financial accounts of the year n-1 that become accessible through specialist websites.

Sampling for these variables was decided over inclusion of them in the established census survey, to reduce the additional burden imposed on clients and to protect the developing quality of census returns.

Prior to the survey year 2022 (of year 2021), a rotating sample was chosen over a random survey due to the level of variation in terms of both population and statistical unit size within and between segments to maintain balanced sampling in this way. A sample size of 25% of the overall sector population was chosen as sufficiently big enough to provide a 20% response, a viable annual sample size of the aquaculture population generally,

notwithstanding the variation in segment population sizes. By this sample size, clients would be approached only once every four years for the more sensitive data.

The 25% sample groups were created from the time of the first upload of 2008 data. The population was ordered by decreasing turnover size at that time and assigned a number 1-4 in repeating sequence down this decreasing turnover column. Thus four profile sampling groups were created and have been maintained until and including the survey of 2020.

Notwithstanding the reasoning behind NPS used above, maintaining the coherence of fixed sample groups response to the sample questionnaire proved increasingly difficult. The Pandemic Unemployment Payment (Covid-19 government assistance scheme), run during the winter of 2020-2021 improved the response rate to the sample survey, Survey participation has also been linked to the EMFAF grant application process from 2021. The opportunity has been provided therefore to move to an administratively easier random sampling regime for the data gathered by sample and via questionnaire. Data sampled from online abridged accounts is still conducted by a non-probability approach in that indicator companies are targeted for their size relative to their segment. Continued variations exist in terms of segment population and statistical unit sizes within segments. Fixed sample group coherence is not an issue for this data source, unlike with questionnaires.

The questionnaire is activated on the BIM website and all producers within the appropriate 25% are invited through the Customer Relationship Management (CRM) system to participate. The same questionnaire can be distributed by email or post and surveys can be conducted by phone if preferred. Non-respondents are pursued by a combination of all these, particularly by phone. Data from online abridged accounts and from in-house datasets are used to validate, cross-reference or to cover certain data shortfalls from the sample questionnaire.

Estimation design

Describe method of calculating population estimate from sample.

Describe method of calculating derived data: e.g. imputed values.

Describe treatment of nonresponse.

National turnover per segment is used to assign a percentage value to the turnover reported or estimated for each Production Unit (PU) of that segment. This proportion is then used to assign the same proportional value to costs and financial data obtained for each PU. For each such variable therefore, each sample return value also has a value proportional to the national value for that variable. The sum of the sample value for each variable is divided by

the sum of assigned proportional values of each PU responding, then multiplied by 100 to raise the sample value to a population estimate for that variable.

$$(\text{Variable Sample sum} / \text{sum of \% s of national turnover of each responding PU turnover}) * 100$$

A disadvantage of this methodology is that, for very small samples obtained for a given variable, the figure calculated is likely to be an over or under-estimate, due to varying segment statistical unit size. Estimation procedures are periodically reviewed.

Variables derived are: 'Imputed value of unpaid labour', certain social variables and for finfish 'Mortality'

Method for calculating 'FTE':

Full Time: >30 Hrs/week or > 40 weeks

Part Time: 10-30 Hrs /week or 13-39 weeks * 40 Hrs

Casual: > 10 Hrs /week or < 13 weeks * 40 Hrs

'Imputed value of unpaid labour':

This is estimated for each sampled business, then for each segment sample, then estimated for the national segment.

Minimum expected value for 'wages and salaries' for the segment is calculated by:

Segment FTE * national minimum wage

Actual 'wages and salaries' value for the segment is obtained by survey.

The two values are compared

If 'Actual value' $\geq$ minimum expected, then no unpaid labour value

If 'Actual value' < minimum expected, then the difference = 'imputed value of unpaid labour'.

Imputed data is obtained for some other social variables. This is done using the empirical value of employment status given for some companies with more than one production unit and culture. The turnover per production unit is known. The turnover contribution per single FTE for the business as a whole is calculated and the empirical breakdown derived from company-level data provided, is used to estimate employment status per production unit.

Non-Respondents are given repeated reminders to participate in the annual survey and will receive several phone calls until a date at the end of March when preparations to report on the annual survey must begin. Late returns are accommodated as much as possible and data updates are always applied as one supplies on the basis of the best available data in hand at a given moment in time.

Error checks

Describe potential errors and how and where in the process these are detected, avoided or eliminated e.g., data: duplication, double counting, respondent error, upload error, processing error etc.

Insufficient sample size can lead to over or under-estimation, depending on the size of the reporting unit, relative to other segment units. Production units making up the segment population vary in size and contribution to the segment's turnover. Errors are detected by viewing time series data for each variable and detecting data spikes that require analysis of cause. Phone surveys are a great source of quality data but incorrect figures may be given or accurate data mis-recorded. Errors of decimal placing, or poor handwriting can result in false data spikes after uploading which become apparent in graphics set up specifically to find such in the database and the cause of the spike can be investigated back to the data supplier if necessary. The use of electronic questionnaire forms is advocated as the most effective way to receive accurate data. Greater engagement with the client or with their contacts within this agency, develops a better understanding of their operation scale and trend, reducing the incidence of uploading data incompatible with operational capability.

Data storage and documentation

Describe how the data is stored.

Initially data is uploaded to Excel storage files and is in the initial stages of integration to a Data warehouse environment. Reporting to date has been at:

www.bim.ie/publications/aquaculture

Provide link to webpage where additional methodological documentation can be found, if any.

Published documentation of methodology of all EUMAP data-collection is found at:

https://www.dcmapp-ireland.ie/sites/default/files/DCF_Files/DCF_Methodology_Economic_IRL_2020.pdf

Aquaculture methodology is described from page 11.

Revision

Describe the frequency of the methodology review e.g., revision of; segmentation, survey method per segment, per variable etc.

This is done annually or as the need arises. The smaller segments require close attention as these may have to be amalgamated with others if in decline or may be new growth segments. Seaweed (i.e., macroalgae) segmentation is under continuing review due to segment growth predictions and in recent years has been too small to report as a dedicated segment. The national macro-algal segment has now developed to the point with sufficient new companies now at a commercial phase that data can be disaggregated into the stand-alone segment 'Longline Macroalgae' and data will be collected for this new segment from 2023 onwards.

In contrast, the land-based shellfish units have declined in output and unit number below the point enabling responsible reporting of them as a separate segment. For data collection from 2023 onwards, land-based shellfish units will be incorporated into the most appropriate larger shellfish segment 'Oyster Other methods' (Oyster on trestles) as the units involved are mainly hatcheries and nurseries that supply this segment.

As a result of these changes to macro algae and land-based shellfish, the 'multispecies' segment is now defunct and will not be used for data collection from 2023 onwards.

Confidentiality

Are procedures for confidential data handling in place and documented?

The frame population are made aware of BIM's privacy policy and data protection policies and procedures.

Are protocols to enforce confidentiality between DCF partners in place and documented?

There is a Data Sharing Agreement between the two main agencies carrying out the DCF work, the Marine Institute and Bord Iascaigh Mhara (BIM).

Are protocols to enforce confidentiality with external users in place and documented?

Data protocols to enforce confidentiality are followed strictly. In cases where there is a risk of natural persons and/or legal entities being identified clustering in the form of appropriate segment amalgamation is applied to report economic variables to ensure statistical confidentiality. For internal business units and some closely related agencies, formal data-sharing agreements between the data controller and data processor must be in place before an appropriate level of data-sharing can be permitted.

Are there any issues with publication of data due to confidentiality reasons? Provide an explanation

Production of data from the salmon on-growing segment is restricted due to the small number of operators involved and the fact that one very much dominates the segment. There is close engagement between BIM, as the data controller, and the concerned parties to manage the appropriate level of reporting of this and similar segments of low populations.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan

Sampling Scheme

No deviation from WP in 2023.

Survey Planning

No deviation from WP in 2023.

Survey Design and Strategy

The original 25% population sample approach documented in the WP has been revised to 33% reflecting the overall greater response rate now attainable each year.

Estimation Design

No deviation from WP in 2023.

Error Checks

No deviation from WP in 2023.

Data Storage and Documentation

No deviation from WP in 2023.

Revision

No deviation from WP in 2023.

Confidentiality

No deviation from WP in 2023.